Radiology Dossier



ملاحظات عامة

- السلايد التالي فيه الcolor code مهم تشوفه وانتبه في شابترات ممكن المسؤول عنها زاد شوي عليها رح يكون مكتوب قبل الشابتر لو في ملاحظات شوفها
- حاولنا نلم المهم في كل المادة وكل دكتور وكل موضوع رح تلاقوا فيه اختلاف شوي بطريقة السرد بما يتناسب معه
- شاملة؟ شوف عيني مستحيل أقلك اه وبعدها يجيك سؤال فلكي أو نكون ناسيين سلايد برمائي متخبي بين +100 سلايد الي في محاضرة ويجي منه شيء وتدعي او تسب علي، ان شاء الله كل الأفكار المطلوبة حطيناها وشملناها وحاولنا كل حرف يكون موجود لكن كونه عمل بشري ممكن ننسى، في حال حد لقى معلومة فلكية جديدة مهمة يبلغنا نضيفها
- هذا العمل مساعدة طلابية من الطلاب للطلاب بهدف لم الملاحظات والشرح الي مرات بكون مش مكتوب لكن مهم ومطلوب
- حاولوا لو في دكتور بخليكم تعملوا سمينارات ما تتعدوا المعلومات والصور الي في الدوسية مشان الامتحان ممكن يجي من السمينارات تاعتكم
- الجماعة الي بتطبع لا تطبعوا الدوسية ما رح تزبط معكم، أولا طويلة، ثانيا كونه الخلفية سودة ما رح تزبط، ثالثا لو طبعتها ما رح تبين الصور منيح ادرسوها الكتروني افضل بكثير
- أخيرا أعزائي الطلبة هذا العمل حرثنا وتعبنا فيه بهدف توفير مادة مكتوبة وشاملة قدر الأمكان ونتمنا منكم المحافظة عليه حتى تستفيدوا منه وغيركم يستفيد ونتجنب الحركات السنفورية، مثلا ألاقي واحد خاش على الدكتور يسأله دوسية اللجنة بتكفي وبالي الطلاب بامتحان من بعد موازي فيه القلب جاي مكان الكبد وقتها انتم الي رح تتضرروا مش نحن بس برضو منك لله عيني

الطب والجراحة لجنة

في حال كان عندك ملاحظات أو حاب تقييم الدوسية أو في مشكلة
أو خطئ رابط التقييم ↓

اضغط هنا لتقييم الدوسية

الطب والجراحة
لجنة

Color Code

- Color code for this chapter
  - Red: Important
  - Red star at the upper Rt. Corner: Important slide
  - Yellow, Olive, and blue: مشان الترتيب
  - Green: معلومات زيادة مش مطلوبة منك أوتها ليلة الامتحان بس شوفها لو قاعد تدرس أول بأول

Chapters

- Chapter 1: General Review of Modalities of Imaging
- Chapter 2: Neuroradiology
- Chapter 3: Chest Imaging
- Chapter 4: Abdomen Imaging
- Chapter 5: Skeletal System Imaging
- Chapter 6: Urinary Tract Imaging
- Chapter 7: Pediatrics
- **Nuclear medicine are not included don't forget to studies the slides**

الطب والجراحة لجنة

Chapter 1
General Review of Modalities of Imaging

Done By: Mahmoud Barakat

Source:

DR. Hana Quidsieh slides 2021/2022; Mutah University

Table of Content


1. Plain X-ray film
2. IVU (IVP) intravenous pyelography or urography
3. Fluoroscopy
4. CT Scan
5. MRI
6. Ultrasound
7. Contrast medias

الهدف من هذا الفصل هو التعرف بشكل عام على طرق التصوير أما التفصيل يكون بالفصول التالية

الطب والجراحة لجنة

Section1: Plain X-ray film

X-ray is an ionizing radiation consisting of electro photons (has direction and energy) coming from x-ray tube → hit the target area of the body (chest, leg, hand,…) → change in the direction and or energy of the photon → passes through the body → hit the film



Plain X-ray film

Advantages

1. Easily performed
2. Available in almost all radiology centers
3. Not coasty
4. First modality of imaging in many radiopathologies
5. Shows bone, metallic object with no artifacts …..

Disadvantages

1. Almost Not allowed in pregnant .
2. Radiation exposure (but smaller dose than CT scan)
3. Limited diagnostic information in any radiological cases.


الطب والجراحة لجنة


Examples of Plain X-rays

1. Chest Xray; Routinely PA (Posteroanterior) view
2. Abdomen Xray; Routinely Erect and Supine views
3. KUB; Area of Kidney, Ureter, Bladder
4. Wrist Xray
5. Knee Xray
6. Cervical Spine Xray


الطب والجراحة لجنة


Examples of Plain X-rays













الطب والجراحة لجنة

Section2: IVU (IVP)

Intravenous Pyelography or Urography

- It is study for the pelvis of kidneys, ureters and urinary bladder


Procedure:

1. We start with KUB (Control film)
2. And then give the patient contrast media I.V. (intravenously)
3. Then do X-ray at different time (immediate, 5min , 10min,…) and continue as each case required.

IVU (IVP) Examples



5 MIN FILM



15 MIN FILM

Section3: Fluoroscopy


It is a dynamic X-ray (video like) with contrast media given to the patient

EXAMPLES:

- Barium swallow (esophagus)
- Barium meal (stomach)
- Barium follow though (small bowel)
- Barium enema (large bowel)
- Hysterosalpingography (uterus)
- Urethrography (urethra)
- MCUG (micturition cystourethrogram) urinary bladder

Barium Swallow

Taking images while the patient is swallowing the oral contrast media
upper (lateral and ap):



Barium Swallow

Taking images while the patient is swallowing the oral contrast media
Lower level (ap and lateral):



Barium Meal:


TAKING IMAGES WITH DIFFERENT VIEWS WHILE THE ORAL CONTRAST IN THE STOMACH








Follow Through



We give oral contrast barium, then we take X-ray films on
Different times for example:
Every 20 minutes in the first hour
Every 30 minutes in the second hour
Every 60 minutes till reaching the terminal ileum
Then we do compression view under Fluoroscope guidance to examine terminal ileum

الطب والجراحة لجنة

Compression view barium follow through



Barium Enema:




Through rectal tube we introduce
barium contrast under fluoroscopy guidance to large bowel only

MCUG (Micturition Cyst Urethrogram)

- Usually used to detect VUR (vesical ureteric reflux)
- We introduce NICN (non ionized contrast media) through foleys catheter to the urinary bladder.
- The contrast media should fill the urinary bladder without retrograde passage to the ureters
- If there is influx of contrast media to ureters it is called VUR (vesico ureteric reflux)

الطب والجراحة لجنة

MCUG



NORMAL
MCUG



VUR IN MCUG
ABNORMAL

الطب والجراحة
لجنة


Urethrogram

We introduce NICM through foleys catheter (it's balloon in the tip of the penis) to see if there is any stricture or rupture in the urethra



Hysterosalpingography

Introduce NICM through catheter or long canula to the uterus , mainly to detect any blockage of fallopian tubes



Section4: CT Scan

It is multiple X-rays beam that penetrate the scanned area and received by detectors and then analyzed by computer

Advantages

1. Rapid scan
2. First choice for trauma cases ,and brain insult
3. Best method for calcification and fractures

Disadvantages

1. High exposure dose
2. Costy
3. Less diagnostic information than MRI
4. Not allowed for pregnants

الطب والجراحة لجنة

CT Scan Windowing

- CT data sets have a very high dynamic range (extending from -1000 HU to 1000 HU) which must be reduced for display or printing
- This is typically done via a process of "windowing", which maps a range (the "window") of pixel values to a grayscale ramp.

ال CT-Scan بيعطي مجال واسع من الفرق في ال radiopacity بمجال بيصل من -1000 ل1000 بالتالي ما في كومبيوتر بتحمل هذا الكم الهائل من الفروقات وما بقدر يعرضهم بصورة بتحوي كل هذا المدى مرة وحدة فيتم تقسيم المدى على شكل نوافذ بتحدد قيمة عليا وقمة سفلى للنافذة وبعرض الكومبيوتر أي شيء ال radiopacity له ضمن القيم هذه (العليا والصغرى) بالرمادي حسب اختلاف الكتل بينها وأي شيء أعلى من القيمة العليا ببين أبيض وأي شيء أقل من القيمة الصغرى ببين أسود

HU means Hounsfield unit

The Hounsfield scale is a quantitative scale for describing radiodensity. It is frequently used in CT scans, where its value is also termed CT number.

الطب والجراحة لجنة

CT Scan Windowing

- CT data sets have a very high dynamic range (extending from -1000 HU to 1000 HU) which must be reduced for display or printing
- This is typically done via a process of "windowing", which maps a range (the "window") of pixel values to a grayscale ramp.

ال CT-Scan بيعطي مجال واسع من الفرق في ال radiopacity بمجال بيصل من -1000 ل1000 بالتالي لو اعتبرنا -1000 أسود وال+1000 أبيض والقيم الي بينهم درجات من الرمادي ما رح نقدر نميز الأنسجة القريبة على بعض رح تعطي تقريبا نفس اللون فمناخذ window مثلا بين ال-100 و+100 (أغلب الأنسجة أصلا كثافتهم بتكون بين هذول الرقمين) فلما يكون الأي شيء أقل من -100 الي هو بشكل عام هواء بس و أي شيء أكبر من 100 أبيض الي هو العضم والتكلسات بشكل عام وباقي الأشياء ببين الاختلاف بينها بشكل واضح (توزع درجات الرمادي بين ال100 سالب وموجب أوضح من انك توزعهم بين 1000)


HU means Hounsfield unit

The Hounsfield scale is a quantitative scale for describing radiodensity. It is frequently used in CT scans, where its value is also termed CT number.

الطب والجراحة لجنة

CT Scan Windowing

- For example, CT images of the brain are commonly viewed with a window extending from 0 HU to 80 HU. Pixel values of 0 and lower, are displayed as black; values of 80 and higher are displayed as white; values within the window are displayed as a grey intensity proportional to position within the window.
- The window used for display must be matched to the X-ray density of the object of interest, in order to optimize the visible detail. E.g.:



Multiplanar reformation or reconstruction (MPR)

- involves the process of converting data from an imaging modality acquired in a certain plane, usually axial, into another plane. It is most commonly performed with thin-slice data from volumetric CT in the axial plane, but it may be accomplished with scanning in any plane and whichever modality capable of cross-sectional imaging, including magnetic resonance imaging (MRI), PET and SPECT.
- The acquired data, for example from the axial plane, can then be converted to non-axial planes such as coronal, sagittal or oblique.
- بالعربي نحن منأخذ صور متعددة من الX-ray وراء بعض لو صفتناهم فوق بعض بعطونا مجسم 3D منه منقدر نشوف الصور على محاور غير الaxial مثلا زي الcoronal والsaggital

- In addition, with the aid of various software, several manipulations of the data can be made. One of these methods, called curved planar reformation (CPR), involves tracing a structure, usually a blood vessel, and generating a planar (two-dimensional) image that transects the structure along its short axis. This is very useful in CT/MR angiography (CTA).

الطب والجراحة
لجنة


Section5: MRI

- MRI is a large very strong magnetic field
- It is not allowed to enter any ferromagnetic object to MRI room at all
- You must take good history from the patient with his documented sign that he has no “MRI non compatible” prosthesis or pacemaker (documented)

- **Unlike CT where we describe "density", images are described by signal intensity ("hyper-" bright, "hypo-" dark)**

MRI

- In MRI we scan the patient with three different planes:
  Axial, Coronal and Sagittal
- And in many different sequences, so it takes long time, examples of sequences in axial plane. (further discussed in neuroradiology part)



MRI

Advantages

1. No radiation exposure, however, pregnants in the first trimester are not allowed to have MRI because of lack of enough safety research
2. High diagnostic information
3. Very sensitive for early brain ischemia diagnosis

Disadvantages

1. It is relatively long time for scanning 15 min – 1 hour
2. Not allowed for patient with (non-MRI computable prosthesis)
3. Not optimum for calcification .
4. The machine has long closed tube that may trigger claustrophobia for some patients
5. The machine has very loud noise.
6. Costy

الطب والجراحة لجنة

MRI ACCIDENTS















Section6: Ultrasound

Examples: Kidney ultrasound, thyroid ultrasound, liver ultrasound





Ultrasound

Advantages

1. No harmful radiation exposure
2. Available
3. Not coasty
4. Best method for hydronephrosis and gall bladder stone

Disadvantages

1. Operator dependent
2. Limitation technical factors by opacity ,uncooperative patient, excessive gases,…..

الطب والجراحة لجنة

Section7: Contrast medias

It is material given orally to opacify bowel or iv to opacify vessels or some lands of tumor

1. Oral
   A. Barium sulfate
   B. NICM (Non-Ionized Contrast Media)
2. IV
   A. NICM (Non-Ionized Contrast Media)
   B. Gadolinium
3. Other

Contrast medias

Oral Contrast Media

A. Barium sulfate

- used for swallow ,meal, follow through and enema , and in diluted form for abdomen CT-scan
- If it enter the peritoneal cavity, it may cause severe peritonitis, so it is not used when there is suspicion of perforation or leak.

B. NICM (Non-Ionized Contrast Media)

- Used as oral contrast for CT-scan abdomen to opacify OWEL
- And when there is suspicion of perforation or leak

الطب والجراحة لجنة

Contrast medias

I.V CONTRAST

A. NICM (Non-Ionized Contrast Media)
   - History of allergy must be taken carefully ,if there is a history of allergy like asthma or penicillin use another image modality or prepare the patient with oral or iv corticosteroid
   - Check the kidney function test
   - It is used in CT-scan and IVP

B. Gadolinium used for MRI

Others

- Intrauterine contrast in hysterosalpingogram we use NICM
- In urethrogram and MCUG we use NICM

الطب والجراحة لجنة

Chapter 2
Neuroradiology

Done By: Mahmoud Barakat

Sources:

DR. Hana Quidsieh slides 2021/2022; Mutah University

Radiopaedia.org

Table of Content

1. **Imaging modalities**
   - CT Head
   - MRI brain
   - Digital subtraction angiography
2. **Neuroanatomy**
   - Falx cerebri & Tentorium cerebri
   - Lobes & Sulci
   - Corpus callosum & Internal capsule
   - Basal Ganglia
   - Brainstem & Cerebellum
   - Ventricular System
   - Cisterns
   - Blood Supply
   - Quiz
3. **Pathology**
   - Traumatic brain injuries
   - Strokes
   - Intracranial Masses
   - Miscellaneous Topics
4. **Key Findings**
   - Midline shift
   - Mass effect
   - Hydrocephalus
   - Brain mass lesion
   - Cerebral oedema
   - Intracranial haemorrhage
   - Extra-axial collection
5. **Approach**
6. **Clinical Cases**

Section1: Imaging modalities

Imaging modalities

- **CT Head**
  - Refers to a computed tomography examination of the brain and surrounding cranial structures. It is most commonly performed as a non-contrast study, but the addition of a contrast-enhanced phase is performed for some indications.
- **MRI Brain**
  - It is the main method to investigate conditions such as multiple sclerosis and headaches and used to characterize strokes and space-occupying lesions.
- **Ultrasound carotids*** (معلومة إضافية)
  - **US (ultrasound) carotids** is a standard test performed as part of the assessment of the cranial arterial blood supply.
  - **Indications:** assessment of carotid stenosis


الطب والجراحة لجنة

CT Head (CT Brain)

Indications of brain CT:

1. Head injury
2. Head infection (in a combination with contrast media)
3. Brain hemorrhage (sensitive)
4. Stroke we may need CTA (study of vessel with contrast media).
5. Increase intracranial pressure (hydrocephalus)
6. Others; dementia, tumor, and seizure.


CT Head - Windows



BRAIN window	BONE window	SUBDURAL window
W:80 L:40	W:2500 L:480	W:350 L:90

الطب والجراحة لجنة


CT Head - Densities description




CSF-3HU
Grey matter-38HU
White matter-30HU
Fresh blood-70-80HU

الطب والجراحة لجنة

CT Head Densities description




CT Head - Densities description



Remember this is a brain window thus the densities are descried in relativity to brain (soft tissue) density, also remember that any density out of the window range are just masked out with white for hyper dense and black for hypodense thus you can't relay on this window to truly view them

الطب والجراحة لجنة


بالCT لما تكبس على أي مكان بالصورة او تحدد مكان الجهاز بعطيك معلومات عنه مثل الaverage HU



CT Head - Contrast Media



Without contrast (non enhanced) plain



With contrast (enhanced) plain
the subarachnoid spaces are
hyperdense are hyperdense

Purpose of Contrast Media

- The purpose of contrast in the setting of head imaging is to evaluate the physiological and pathological processes that alter the permeability of the blood-brain barrier that causes abnormal contrast enhancement.
- Physiologically, the blood-brain barrier prevents the leakage of contrast material into the interstitium in the brain, spinal cord and the proximal nerves.
- Alteration of the blood-brain barrier is caused by physiological and pathological processes including:

- angiogenesis: new blood vessel formation
- inflammation
- ischemia
- increased pressure


الطب والجراحة لجنة

Indication of Contrast Media

1. Tumors:
   A. Brain metastases: variable enhancement of the lesion post-contrast
   B. Meningioma: solid intense enhancement of the lesion post-contrast
   C. Necrotic neoplasm (e.g., glioblastoma) or abscess: outer enhancing ring surrounding a necrotic center
   D. Lymphoma: enhancement of lesion post-contrast
2. Brain abscesses:
   - double rim sign: hypodense outer rim and a hyperdense inner rim
   - single rim of hyperdense or Isodense
   - fluid/pus: hypoattenuating center
3. Meningitis and meningoencephalitis: Leptomeningeal enhancement
4. Multiple sclerosis: contrast enhancement of plaques
5. Ependymitis and ventriculitis: thin linear enhancement of the margins of the ventricles

الطب والجراحة لجنة

Examples of enhancement



Mets enhancement



Abscess:
Ring Enhancement

MCA Aneurysm in CTA



Coronal reconstructions

الطب والجراحة لجنة


CTA

CT angiography is a computed tomography technique used for angiography — the visualization of arteries and veins — throughout the human body. Using contrast injected into the blood vessels, images are created to look for blockages, aneurysms (dilations of walls), dissections (tearing of walls), and stenosis (narrowing of vessel). CTA can be used to visualize the vessels of the heart, the aorta and other large blood vessels, the lungs, the kidneys, the head and neck, and the arms and legs



الطب والجراحة
لجنة

CTA



MRI Brian

- Indications
  - Confirmation of stroke
  - Assessment of intracranial tumor
  - Chronic headache
  - Seizure disorder
- Important pathology
  - Stroke
  - Intracranial tumors
  - Multiple sclerosis

- MRI Brian Sequences
  1. T1-weighted
  2. T2-weighted
  3. FLAIR
  4. GRADIENT ECHO
  5. DWI and ADC (diffusion-weighted imaging and apparent diffusion coefficient)

الطب والجراحة لجنة


T1-weighted

✓ **Recognition**

- Grey matter is darker than the white matter
- Fat is bright
- Fluid (in CSF and orbits) is dark
- New blood is bright

✓ **Useful for**

- Provides the most anatomically relevant images
- Vascular changes +C
- Disruption of BBB +C



الطب والجراحة لجنة

T2-weighted


✓ **Standard sequence**

✓ **Recognition**

- White matter is darker than grey
- Fluid is bright
- Signal voide blood flow


✓ **Useful for**

- Most lesions
- But cannot distinguish edema from CSF




الطب والجراحة لجنة

T1-weighted Vs T2-weighted



T1-weighted



T2-weighted

الطب والجراحة
لجنة

FLAIR sequence (fluid attenuation inversion recovery)


✓ Commonly used sequence

✓ **Recognition**

- Similar to T2, but the free-flowing fluid (CSF) is darker or "suppressed"


✓ **Useful for**

- Areas of oedema or inflammation
- Identify plaques in multiple sclerosis (especially periventricular)
- Can improve grey-white differentiation



الطب والجراحة لجنة


T2-weighted Vs FLAIR



GRADIENT ECHO

GRE

Gradient Echo (SWI, T2*)



RECOGNITION

- Paramagnetic substances are dark
  - Blood
  - Calcium
  - Other metals

USEFUL FOR

- Early Hemorrhage
- Old Hemorrhage

GRADIENT ECHO



DWI and ADC

- ✓ Uses the diffusion of water molecules to generate contrast in MR images. Show how easily water moves around
- ✓ Molecular diffusion in tissues is not random, but reflects interactions with many obstacles, such as macromolecules, fibers, and membranes. Water molecule diffusion patterns can therefore reveal microscopic details about tissue architecture, either normal or in a diseased state.
- ✓ Restricted diffusion occurs in stroke, abscesses and cellular tumors
- ✓ADC map, is an MRI image that more specifically shows diffusion than conventional DWI, by eliminating the T2 weighting that is otherwise inherent to conventional DWI.

الطب والجراحة لجنة

Diffusion MRI (DWI)

DWI

Diffusion Weighted Imaging (ADC)



RECOGNITION


- Fluid restriction is bright (cytotoxic edema)
- Must correlate with ADC
  - Fluid restriction is dark
  - Rule out “T2 Shine Through”

USEFUL FOR

- Ischemia
- Abscess

الطب والجراحة لجنة

DWI and ADC



MRI Brian Sequences Summary

- **T1-weighted**
  - ✓ Provides the most anatomically-relevant images because:
    fluid (in CSF and orbits) is dark & grey matter is darker than the white matter
- **T2-weighted**
  - ✓ Useful in most lesions but cannot distinguish edema from CSF
- **FLAIR**
  - ✓ Similar to T2, but the fluid is darker or "suppressed"
  - ✓ Useful for areas of oedema or inflammation
  - ✓ Used to identify plaques in multiple sclerosis (especially periventricular)
- **DWI**
  - ✓ These "blocky" images show how easily water moves around
  - ✓ Restricted diffusion occurs in stroke, abscesses and cellular tumors

الطب والجراحة لجنة

MRA


- A magnetic resonance angiogram (MRA) is a type of magnetic resonance imaging (MRI) scan that uses a magnetic field and pulses of radio wave energy to provide pictures of blood vessels inside the body.
- In many cases MRA can provide information that can't be obtained from an X-ray, ultrasound, or computed tomography (CT) scan.




الطب والجراحة لجنة

Digital subtraction angiography

- **Digital subtraction angiography (DSA)** is a fluoroscopic technique used extensively in interventional radiology for visualizing blood vessels.
- Radiopaque structures such as bones are eliminated ("subtracted") digitally from the image, thus allowing for an accurate depiction of the blood vessels.




Digital subtraction angiography





Digital subtraction angiography





Section2: Neuroanatomy

Neuroanatomy

- In this section we will try to be as minimalist as possible as this is not a neurology dossier, we will discuss the important anatomical concepts regarding neuroradiology and will try to provide you with some extra tools to help you to understand these concepts.
- Remember the following slides are meant to help you understand some concepts and are not meant to be learnt by heart

Click on the buttons bellow to view on web browser

Anatomy 2019: Brain Slides contributed by: Assoc Prof Craig Hacking	Radiology Cafe Head and spine anatomy
Anatomy 2019: Ventricles and cisterns Slides contributed by: Dr Andrew Dixon	
Anatomy 2019: Craniocervical vessels Slides contributed by: Assoc Prof Craig Hacking	

الطب والجراحة لجنة


Falx cerebri & Tentorium cerebri

Section2: Neuroanatomy

Falx cerebri

Gross anatomy


- The falx cerebri is a double-fold of dura mater that descends through the interhemispheric fissure in the midline of the brain to separate the two cerebral hemispheres.
- The falx cerebri is relatively thin anteriorly where it attaches to the crista galli of the ethmoid bone but is broader posteriorly where it attaches to the superior surface of the tentorium cerebelli inferiorly.
- It attaches superiorly to the midline of the cranium and extends posteriorly to attach to the internal occipital protuberance.



Falx cerebri

Gross anatomy

- Relations
  - superior margin contains the superior sagittal sinus
  - free edge between its attachments inferiorly closely related to the superior surface of the corpus callosum and contains the inferior sagittal sinus
  - straight sinus is included in the line of attachment between the falx cerebri and the tentorium cerebelli




Falx cerebri – CT-Scan

Radiographic features


CT

- anterior midline linear density near the vertex (red line)
- triangular density inferiorly and posteriorly on axial sections
- partially calcified in 7% of individuals



(CT) Axial

Falx cerebri – CT-Scan




Red dotted line: Falx cerebri
Yellow dotted line: Tentorium cerebri

Falx cerebri – MRI

Radiographic features

MRI


- thin membrane on T1W and T2W images
- calcifications visible on T1W imaging as hyperintensities and hypointensities on T2W imaging



MRI Sagittal T2

الطب والجراحة لجنة

Falx cerebri – MRI



Axial T2


Axial T1

Red dotted line: Falx cerebri

Tentorium cerebri

Gross anatomy


- The tentorium cerebelli is attached to the falx cerebri at its midline, and this attachment contains the straight sinus. This attachment is more superior than its anterolateral and posterolateral attachments giving it a "tented" appearance.
- The anterior portion contains the tentorial notch (or tentorial incisure/incisura) which is a U-shaped (concave) free edge and extends anteriorly to attach to the anterior clinoid processes and forms the lateral part of the cavernous sinus. The deep tentorial notch surrounds the tentorial hiatus, which allows communication between the supratentorial and infratentorial comartments.



Tentorium cerebri

Gross anatomy; cont.

- The anterolateral margin attaches to the superior border of the petrous part of the temporal bone. This margin lies inferior to the tentorial notch and extends anteriorly to attach to the posterior clinoid process and contains the superior petrosal sinus.
- The posterolateral margin separates into the two layers of dura and attaches to the edges of the transverse sulci of the occipital bone. As it extends anteriorly it attaches to the posteroinferior angles of the parietal bone. At its most posterior aspect, it is attached to the internal occipital protuberance and posterolaterally contains the transverse sinus.



Tentorium cerebri

Radiographic features

CT

✓non-contrast

- appreciated best on coronal and sagittal views as a linear structure separating the occipital lobe from the cerebellum (or supratentorial compartment from the infratentorial compartment)
- on coronal views is visible as being continuous with the falx cerebri

✓contrast-enhanced

- on axial views tentorial margins best appreciated with IV contrast as uniform linear densities


الطب والجراحة لجنة

Tentorium cerebri – CT-Scan



Red dotted line: Tentorium cerebri

Tentorium cerebri – MRI (T1-weighted)



Clinical importance of falx cerebri & tentorium cerebri

Falx cerebri

- Cingulate gyrus can herniate under the falx cerebri resulting in subfalcine (cingulate) herniation which may result in compression of the anterior cerebral artery, which causes infarction

Tentorium cerebri

- Transtentorial herniations (e.g., uncal herniation, downward (central) transtentorial herniation) is a potentially lethal complication of epidural hematoma
- CNS tumors are classified according to its location in respect to the tentorium
  - In children, primary CNS tumors are usually infra-tentorial
  - In adults, primary CNS tumors are usually supra-tentorial

الطب والجراحة لجنة

Lobes & Sulci

Section2: Neuroanatomy

Lobes and Main Sulci

Lobes


1. Frontal Lobe
2. Parietal Lobe
3. Temporal Lobe
4. Occipital Lobe
5. Insular Lobe


Main Sulci


1. Medial longitudinal fissure
2. Center sulcus
3. Lateral sulcus
4. Parieto-occipital sulcus
5. Calcarine sulcus
6. Cingulate sulcus


الطب والجراحة لجنة


Lobes & Main Sulci














Lobes


Lobes
Frontal lobe
Central s.
Parietal lobe
Parieto-occipital s.
Occipital lobe
Adapted from illustration from "Sobotta's Textbook and Atlas of Human Anatomy" 1908, now in the public domain.
Radiopaedia.org



Frontal lobe
Parietal lobe
Insular cortex
Occipital lobe
Temporal lobe
Adapted from illustration from "Sobotta's Textbook and Atlas of Human Anatomy" 1908, now in the public domain.
Radiopaedia.org



Cingulate gyrus
Frontal lobe
Parietal lobe
Corpus callosum
Occipital lobe
Temporal lobe
Adapted from illustration from "Sobotta's Textbook and Atlas of Human Anatomy" 1908, now in the public domain.
Radiopaedia.org

Frontal lobe

Gross anatomy

- The frontal lobe resides largely in the anterior cranial fossa, lying on the orbital plate of the frontal bone.
- Its most anterior part is known as the frontal pole and extends posteriorly to the central (Rolandic) sulcus which separates it from the parietal lobe.
- Posteroinferiorly it is separated from the temporal lobe by the lateral sulcus (Sylvian fissure) and the insular cortex which is hidden deep to the lateral sulcus.
- The interhemispheric fissure separates its medial surface from the contralateral frontal lobe.







Parietal lobe

Gross anatomy

- The parietal lobe extends from the central sulcus anteriorly, which separates it from the frontal lobe, to the parieto-occipital fissure posteriorly, which separates it from the occipital lobe.
- The lateral sulcus corresponds to its inferolateral boundary, separating it from the temporal lobe.
- Medially, it is confined by the medial longitudinal fissure which divides both cerebral hemispheres.







Temporal lobe

Gross anatomy

- The lobe extends superiorly to the Sylvian fissure, and posteriorly to an imaginary line; the lateral parietotemporal line, which separates the temporal lobe from the inferior parietal lobule of the parietal lobe superiorly and the occipital lobe inferiorly.
- The middle cranial fossa forms its anterior and inferior boundaries.







Occipital lobe

Gross anatomy

- The boundaries of the occipital lobe have been arbitrarily (اعتباطيا) defined, giving it a triangular shape. It is separated from the parietal and temporal lobes on the medial surface by the parieto-occipital sulcus and on the lateral side by the lateral parietotemporal line, a near vertical imaginary line extending from the preoccipital notch of the temporal lobe superiorly to the parieto-occipital sulcus.
- Medially, it is confined by the medial longitudinal fissure which divides both cerebral hemispheres.
- The tentorium cerebelli forms its inferior border and the occipital bone its posterior margin.







Insular cortex

Gross anatomy

- The insular cortex lies deep to the lateral sulcus (Sylvian fissure), which separates the frontal and parietal lobes dorsally from the temporal lobe ventrally.
- The overlying cortical areas formed by the frontal, parietal, and temporal lobes are known as opercula (meaning "lids").





Main Sulci

- Medial longitudinal fissure is the deep groove within the midline separating both cerebral hemispheres and containing the falx cerebri.
- Center sulcus separates the frontal lobe from the parietal lobe, and more specifically separates the primary motor cortex anteriorly from the primary somatosensory cortex posteriorly
- Cingulate sulcus is situated directly superior to the cingulate gyrus, which is formed by the medial surface of the frontal lobes that is directly above the corpus callosum.
- Marginal sulcus is the extension of the cingulate sulcus posterosuperiorly, separating the paracentral lobule from the precuneus of the parietal lobe on the medial surface of the cerebral hemispheres.


الطب والجراحة لجنة

Main Sulci cont.

- Lateral sulcus separates the frontal and parietal lobes superiorly from the temporal lobe inferiorly. The insula is located immediately deep to the Sylvian fissure.
- Parieto-occipital sulcus is an oblique sulcus which demarcates the occipital lobe from the parietal lobe on the medial surface of the cerebral hemisphere. The parieto-occipital sulcus joins the calcarine sulcus to form a pattern similar to the letter Y which has been placed sideways. Specifically, it delineates the precuneus superiorly from the cuneus inferiorly.
- Calcarine sulcus is located on the medial surface of the occipital lobe and divides the visual cortex (a.k.a. calcarine cortex) into two


الطب والجراحة لجنة

Main Sulci



Adapted from illustration from "Sobotta's Textbook and Atlas of Human Anatomy" 1908, now in the public domain.

Frank Gaillard
Radiopaedia.org




Adapted from illustration from "Sobotta's Textbook and Atlas of Human Anatomy" 1908, now in the public domain.


Frank Gaillard
Radiopaedia.org

الطب والجراحة لجنة

Sylvian Fissure (lateral sulcus)



Sylvian Fissure (lateral sulcus)




Red dotted line: Sylvian Fissure (lateral sulcus)

Center & Cingulate sulci

How to locate the center sulcus on the axial view ?

Method 1: by locating the marginal sulcus which is a mustache like structure coming out of the middle line posteriorly; the center sulcus is directly anterior to it




Central sulcus and Marginal sulcus

الطب والجراحة لجنة

Center & Cingulate sulci

How to locate the center sulcus on the axial view ?

Method 2: by locating the superior frontal sulcus which is a long sulcus runs posteriorly from the frontal pole, parallel to the interhemispheric fissure, back to the precentral sulcus which it intersects in an “Y”, “L” or “T” junction. The center sulcus is just posterior to the precentral sulcus




Central sulcus, Precentral sulcus and
Superior Frontal sulcus

الطب والجراحة
لجنة

Center & Cingulate sulci

How to locate the center sulcus on the axial view ?

Method 3: by identifying the sigmoidal hook sign (Sigmoidal hook sign or omega sign is a term sometimes used to denote the appearance of the hand motor area (hand knob), projecting posteriorly from the precentral gyrus. This is a helpful feature in identifying the central sulcus.)




The term is however not terribly well known and is therefore probably not very helpful and is probably best avoided.

Center & Cingulate sulci

How to locate the center sulcus on the sagittal view ?

First, we locate the marginal sulcus which is the posterosuperior projection of the cingulate sulcus. The center sulcus is just anterior to it




Cingulate sulcus and Marginal sulcus


Center sulcus

الطب والجراحة لجنة

Parieto-occipital & calcarine sulci



Parieto-occipital & calcarine sulci



Red dotted line: Parieto-occipital sulcus
Yellow dotted line: calcarine sulcus

الطب والجراحة لجنة

Corpus callosum & Internal capsule

Section2: Neuroanatomy


Corpus callosum

Summary

- located inferior to the cerebral cortices, and superior to the thalamus
- connects left and right cerebral hemispheres
- It is divided into four parts (from anterior to posterior):
  - rostrum
  - genu
  - trunk/body
  - splenium




Corpus callosum – Rostrum



Corpus callosum – Genu



Corpus callosum – Body




Corpus callosum – Splenium




Internal capsule

The internal capsule is a deep subcortical structure that contains a concentration of afferent and efferent white matter projection fibers. Anatomically, this is an important area because of the high concentration of both motor and sensory projection fibers. Afferent fibers pass from cell bodies of the thalamus to the cortex, and efferent fibers pass from cell bodies of the cortex to the cerebral peduncle of the midbrain. Fibers from the internal capsule contribute to the corona radiata.




Internal capsule – Anterior limb



anterior limb internal capsule

OASIS

Internal capsule – Genu




genu internal capsule

OASIS

bmv© headneckbrainspine.com

Internal capsule – Posterior limb



Basal Ganglia


Section2: Neuroanatomy

Basal ganglia


The basal ganglia are a group of grey matter nuclei in the deep aspects of the brain that is interconnected with the cerebral cortex, thalami and brainstem.

In a strict anatomical sense, it contains three paired nuclei that together comprise the corpus striatum:


- caudate nucleus
- lentiform nucleus
- putamen
- globus pallidus




Head of caudate nucleus



Putamen



Globus pallidus externa




globus pallidus externa

OASIS

bmv© headneckbrainspine.com

Thalamus



thalamus


OASIS

Brainstem & Cerebellum

Section2: Neuroanatomy


Brainstem


- The brainstem is the most caudal part of the brain. It adjoins, is structurally continuous with the spinal cord and consists of the:
  - Midbrain
  - Pons
  - Medulla oblongata



الطب والجراحة
الجنة


Midbrain – the most rostral part of the brainstem





أهم شيء بالنسبة لنا في الأشعة الأن تعرف الشكل حتى تقدر تفرقه بال axial view وتعرف ال blood supply

Midbrain





At higher level


At lower level

الطب والجراحة لجنة

Pons - the middle portion of the brainstem









أهم شيء بالنسبة لنا في الأشعة الأن تعرف الشكل حتى تقدر تفرقه بال axial view وتعرف ال blood supply


الطب والجراحة الجنة


Pons (Pontine)



Medulla oblongata – the most caudal part of the brainstem








أهم شيء بالنسبة لنا في الأشعة الأن تعرف الشكل حتى تقدر تفرقه بال axial view وتعرف ال blood supply

الطب والجراحة الجنة


Medulla oblongata



medulla oblongata

OASIS

Brainstem Blood Supply




Cerebellum


- The cerebellum, sits at the base of the brain in the posterior cranial fossa below the tentorium and behind the brainstem.

Gross anatomy


- The cerebellum has the following features:
  - three surfaces: anterior (petrosal), superior (tentorial), inferior (suboccipital)
  - three fissures: primary (tentorial), horizontal (petrosal), prebiventral/prepyramidal (suboccipital)
  - two hemispheres
  - single median vermis




Cerebellum


Midsaggital section of cerebellum
Superior view of an "unrolled" cerebellum
Anterior lobe
Primary fissure
Posterior lobe
Posterolateral fissure
Flocculonodular lobe
Vermis
Hemisphere
Nodulus
Flocculus

Cerebellum





Ventricular System


Section2: Neuroanatomy

Ventricular System


- The ventricular system consists of:
  - two lateral ventricles
  - a midline third ventricle
  - a fourth ventricle
- The interconnections between the ventricles occur through the following:
  - interventricular foramen (of Monro): lateral to third
  - cerebral aqueduct (of Sylvius): third to fourth
  - median aperture (of Magendie): fourth to cisterna magna
  - two lateral apertures (of Luschka): fourth to cerebellopontine cistern

الطب والجراحة لجنة

Ventricular System




Ventricular System




Lateral Ventricles


- The lateral ventricles are paired CSF-filled spaces in the cerebrum and part of the ventricular system of the brain.
- They can be asymmetrical.
- Each has five divisions, including:
  - anterior/frontal horn
  - inferior/temporal horn
  - body
  - trigone/atrium
  - posterior/occipital horn
    - bilaterally well developed in only 12% of subjects
    - may be absent, poorly developed, asymmetrical




Lateral Ventricles



Lateral Ventricles



Lateral Ventricles




Third Ventricle

Gross anatomy

- The third ventricle is a median cleft between the two thalami, which make up the superior aspect of the lateral walls. The thalami are separated from the hypothalamus (anteriorly) and subthalamus (posteriorly) by the hypothalamic sulcus.
- The anterior wall of the third ventricle is formed from superior to inferior by the columns of the fornix, the anterior commissure and the lamina terminalis.
- Posteriorly, it is bounded from superior to inferior by the habenular commissure, the pineal gland and the posterior commissure.
- The roof is formed by the choroidal fissure and the bodies of the fornix.
- The floor is bounded from anterior to posterior by the optic chiasm (posterior aspect), the infundibulum of the pituitary gland, the tuber cinereum, the mammillary bodies.

الطب والجراحة لجنة

Third Ventricle




Fourth Ventricle


- The fourth ventricle is one of the components of the ventricular system in the brain, along with the lateral and third ventricles. It extends from the cerebral aqueduct (of Sylvius) rostrally to the obex caudally and is filled with CSF.
- CSF enters the ventricle via the cerebral aqueduct and leaves via one of four routes:
  - via the obex and into the central canal
  - via the median aperture (of Magendie) into the cisterna magna
  - via one of the two lateral apertures (of Luschka) into the cerebellopontine cistern
- It is characteristically diamond-shaped in sagittal cross-section and located dorsal to the pons or upper part of the medulla oblongata. The roof, located dorsally is formed by the superior medullary velum and inferior medullary velum overlying the cerebellum. The floor, located ventrally, is formed by the rhomboid fossa, a diamond-shaped depression on the dorsal surface of the pons and upper half of the medulla. The sidewalls are formed by the vela and cerebellar peduncles. The peak of the fourth ventricle within the cerebellum is the fastigium and it underlies the fastigial nucleus. It is widest at the level of the pontomedullary junction. The obex is the most caudal tip of the fourth ventricle.

الطب والجراحة لجنة

Fourth Ventricle



Foramen of Magendi




Cisterns


Section2: Neuroanatomy

Cisterns

1. Interpedicular cistern
2. Ambient cistern
3. Quadrigeminal cistern
4. Suprasellar cistern
5. Cisterna magna
6. Pre pontine-cistern

Interpedicular cistern






Located between the cerebral peduncles.

Ambient cistern


Gross anatomy

- The ambient cistern is a thin, sheet-like extension of the quadrigeminal cistern that extends laterally around the midbrain and posterior to the thalami. It acts as the connection between the quadrigeminal cistern and the interpeduncular cistern. The term "ambient cistern" may be used to refer to these connections and the quadrigeminal cistern as a whole.



Quadrigeminal cistern


- It is located between the colliculi, the splenium of the corpus callosum and the superior surface of the cerebellum and extends from the third ventricle to the great cerebral vein.



Suprasellar cistern

Location

- The suprasellar cistern is located above the Sella turcica, under the hypothalamus and between the uncus of the temporal lobes. It has roughly the shape of a pentagon at the level of the optic chiasma, and of a six-point star at the level of cerebral peduncles.



Cisterna magna


Gross anatomy


- The cisterna magna is located between the cerebellum and the dorsal surface of the medulla oblongata at and above the level of the foramen magnum.



Prepontine cistern

- located ventral to the pons and dorsal to the clivus.





Blood Supply

Section2: Neuroanatomy


Circle of Willis


Gross anatomy

- Vessels comprising the circle of Willis include:
  - ✓Anterior circulation
    - left and right internal carotid arteries (ICA)
    - horizontal (A1) segments of the left and right anterior cerebral arteries (ACA)
    - single anterior communicating artery (ACOM)
  - ✓posterior circulation
    - left and right posterior communicating arteries (PCOM) (although some consider the PCOM to be anterior circulation)
    - horizontal (P1) segments of left and right posterior cerebral arteries (PCA)
    - single basilar artery (tip)


الطب والجراحة لجنة

Circle of Willis





Circle of Willis (MRA)




MRA


Check out the MRA sequence of this normal case

3minutes video explaining normal MRA of the brain


الطب والجراحة
لجنة

Cerebral Blood Supply







Cerebral Blood Supply




Brainstem Blood Supply



Dural venous sinuses






Dural Venous Sinuses | 3D Anatomy Tutorial

Quiz

Section2: Neuroanatomy

Exam 1






Answer Exam 1

A. Orbit
B. Sphenoid Sinus
C. Temporal Lobe
D. External Auditory Canal
E. Mastoid Air Cells
F. Cerebellar Hemisphere

الطب والجراحة
لجنة

Exam 2







Answer Exam 2

A. Frontal Lobe
B. Frontal Bone (Superior Surface of Orbital Part)
C. Dorsum Sellae
D. Basilar Artery
E. Temporal Lobe
F. Mastoid Air Cells
G. Cerebellar Hemisphere

Exam 3






Answer Exam 3

A. Frontal Lobe
B. Sylvian Fissure
C. Temporal Lobe
D. Suprasellar Cistern
E. Midbrain
F. Fourth Ventricle
G. Cerebellar Hemisphere

Exam 4







Answer Exam 4

A. Falx Cerebri
B. Frontal Lobe
C. Anterior Horn of Lateral Ventricle
D. Third Ventricle
E. Quadrigeminal Plate Cistern
F. Cerebellum

Exam 5






Answer Exam 5

A. Anterior Horn of the Lateral Ventricle
B. Caudate Nucleus
C. Anterior Limb of the Internal Capsule
D. Putamen and Globus Pallidus
E. Posterior Limb of the Internal Capsule
F. Third Ventricle
G. Quadrigeminal Plate Cistern
H. Cerebellar Vermis

Exam 6






Answer Exam 6

A. Falx Cerebri
B. Frontal Lobe
C. Body of the Lateral Ventricle
D. Splenium of the Corpus Callosum
E. Parietal Lobe
F. Occipital Lobe
G. Superior Sagittal Sinus

Exam 7






Answer Exam 7

A. Falx Cerebri
B. Frontal Lobe
C. Body of the Lateral Ventricle
D. Splenium of the Corpus Callosum
E. Parietal Lobe
F. Occipital Lobe
G. Superior Sagittal Sinus

Exam 8






Answer Exam 8

A. Falx Cerebri
B. Sulcus
C. Gyrus
D. Superior Sagittal Sinus

Exam 9




Answer Exam 9

a. genu of corpus callosum
b. anterior limb internal capsule
c. lentiform nucleus
d. posterior limb internal capsule
e. Sylvian fissure
f. thalamus
g. caudate head
h. genu of internal capsule
i. third ventricle
j. trigone (atrium) of lateral ventricle
k. splenium of corpus callosum
l. superior sagittal sinus

الطب والجراحة لجنة


Exam 10



Answer Exam 10

a. fornix
b. interventricular foramen (of Munro)
c. genu of corpus callosum
d. optic recess of third ventricle
e. suprasellar cistern
f. basilar artery
g. third ventricle
h. quadrigeminal cistern
i. cerebral aqueduct
j. straight sinus
k. 4th ventricle
l. cisterna magna


Exam 11


Cor neck CTA thick MIP
a
b
c
d
e
f
g
h
i
j
A.Prof Frank Gaillard, rID: 38570

Answer Exam 11

a. MCA (M2 segment)
b. superficial temporal artery
c. internal carotid artery ascending carotid canal portion
d. common carotid artery
e. vertebral artery
f. ACA
g. basilar artery
h. ICA cavernous portion
i. left vertebral artery
j. internal jugular vein

Exam 12



Answer Exam 12

a. right ophthalmic artery
b. left A2 segment of ACA
c. left M1 segment of MCA
d. left posterior communicating artery
e. left M3 branch of MCA
f. left P2 segment of PCA
g. left V4 segment of vertebral artery
h. right P3 segment of PCA
i. right superior cerebellar artery
j. right M2 branch of MCA
k. C2 (petrous) segment of the right ICA

الطب والجراحة لجنة

Scrollable labeling questions

CT head axial - labelling questions

CT head sagittal - labelling questions

CT head coronal - labelling questions

Section3: Pathological Conditions

Traumatic brain injuries

Section3: Pathological Conditions

Traumatic brain injuries

In the acute setting patients can present with primary brain damage

- Hematomas
- Cerebral contusions
- Diffuse Axonal injury

Secondary brain damage can also occur and manifests as

- Generalized cerebral oedema
- Hypoxic-ischemic brain injury
- Ischemic stroke from traumatic arterial dissection
- "big black brain"

Long-term sequelae of head trauma include:

- Encephalomalacia/gliosis
- Chronic subdural hematomas / CSF hygromas
- Chronic traumatic encephalopathy
- Depression, anxiety and alcohol excess
- Increased risk of schizophrenia, bipolar disorder and organic mental disorders

الطب والجراحة لجنة

Traumatic brain injuries

Complications

- Severe mass effect can result in:
  - Midline shift: associated with worse prognosis
  - Cerebral herniation: often requires urgent treatment
  - Hydrocephalus: can also be a chronic non-mass effect related complication

Associations

- Other traumatic injuries are common:
  - Cervical spine injury
  - Skull fracture
  - Facial fracture
  - Pneumocephalus
  - Extracranial injuries

الطب والجراحة لجنة

Primary brain damage

- Hematomas; we will discuss all of them not only the traumatic ones
- Cerebral hemorrhagic contusion
  - intermediary injury
- Diffuse axonal injury (DAI)

الطب والجراحة لجنة


Hematomas

- **Scalp Hematomas**
  - Caput succedaneum
  - Subgaleal hematoma
  - Cephalohematoma
- **Intracranial hemorrhage**
  - **Extra Axial**
    - Epidural hematoma
    - Subdural hematoma
    - Subarachnoid hematoma
    - Intraventricular hematoma
  - **Intra Axial**
    - Intracerebral hemorrhage
      - Lobar hemorrhage
      - Basal ganglia hemorrhage
      - Pontine hemorrhage
      - Cerebellar hemorrhage

الطب والجراحة لجنة

Hematomas

- **Appearance on CT:**
  - Acute (hours to days) - New blood is White
  - Subacute (days to weeks) - Grey
  - Chronic (weeks to months) - Old blood is Dark
  - Acute on Chronic - Layering effect



Scalp Hematomas


- Commonly occurs either following a perinatal injury at delivery or as part of head trauma.
- There are three types of hematoma, which are defined by their location within the scalp, in particular their location as related to the galea aponeurosis and skull periosteum
  - Caput succedaneum: subcutaneous hematoma superficial to the galea aponeurosis, most commonly secondary to vacuum-assisted delivery
  - Subgaleal hematoma: hematoma within the potential space between the galea aponeurosis and the skull periosteum
  - Cephalohematoma: subperiosteal and therefore bound by the suture line

الطب والجراحة لجنة

Layers of the scalp (mnemonic)


Mnemonic

- **S:** skin
- **C:** connective tissue
- **A:** (galea) aponeurosis
- **L:** loose connective tissue
- **P:** periosteum



Caput succedaneum

- Caput succedaneum is a manifestation of birth trauma, and it consists of a subcutaneous serosanguineous fluid collection external to the galea aponeurosis in the newborn's scalp. The fluid collection is extra-periosteal.
- Caput succedaneum results from pressure on the presenting part of the skull against the dilating cervix. It extends across the midline and over suture lines and is associated with head molding.
- Caput succedaneum does not usually cause any complication and usually resolves over the first few days. Management is expectant.





Subgaleal hematoma

Subgaleal hematoma describes scalp bleeding in the potential space between the periosteum and the galea aponeurosis. It is a rare but possibly lethal emergency.

It most commonly occurs after vacuum-assisted and forceps delivery but may also be seen following head trauma or occur spontaneously.

مع انها مش مشروحة بس جاءت ضمن خيارات واحد من الامتحانات فاعرفها ويفضل تعرف أي شيء traumatic كثير بيجي عليهم






View case on radiopaedia

Subgaleal hematoma


Brain Window

Bone Window



Cephalohematoma

Cephalohematomas are traumatic subperiosteal hematomas of the skull that are usually caused by birth injury. They are bound between the periosteum and cranium, and therefore cannot cross sutures. Being bound by a suture line distinguishes them from Subgaleal hematoma, which can cross sutures.



Extra Axial Hematomas Summary

❖Epidural hematoma

- Lens (lentiform) in shape
- Respect the sutures
- Can cause mass effect (discussed later)
- Most common from the middle meningeal artery

❖Subdural hematoma

- Lacunar in shape
- Doesn't respect suture
- Less likely to cause mass effect
- Most common from the bridging veins

السلايد هذا تلخيص للأكم من سلايد الي بعده مهم جدا

❖Subarachnoid hematoma


- Bleeding in sulci & citterns
- Usually found centrally (around the circle of Willis) but can occur in other parts of the brain
- Most commonly due to rupture of an intracranial aneurysm (berry aneurysm)
- Commonly present with "The worst headache in my life" complain

❖Intraventricular hematoma


- Inside cerebral ventricles
- Can be primary or secondary to a large extraventricular component with secondary extension into the ventricles

الطب والجراحة لجنة

Extra Axial Hematoma



Epidural Hematoma





Epidural hematoma (Extradural hematoma)

- Extradural (or epidural) hematoma is a post-traumatic event resulting from injury to an intracranial artery, most commonly the middle meningeal artery. Leakage from an injured artery results in collection of blood which strips the dura mater away from the inner table of the skull.
- Extradural hematoma results in formation of a lens-shaped collection. This is because the dura mater is strongly adhered to the skull in the region of the sutures. Stripping of the dura from the skull is limited at these points resulting in limitation of the extent of a collection (no cross of the sutures)

الطب والجراحة لجنة

Subdural Hematoma





Subdural Hematoma

Unlike other hematomas that always present acutely, subdural hematomas can present acute or chronic




Acute SDH
(Hyperdense)



Subacute SDH
(Iso dense)



Chronic SDH
(Hypodense)



Acute on Top
of Chronic

Subdural Hematoma

- The cerebral veins are fragile. Risk of injury to these veins is increased in elderly and anti-coagulated patients. Subdural hematoma may result from minor trauma. There is often no clear history of trauma at all.
- A subdural collection is not limited by the attachment points of the dura to bone, as is seen in an extradural hematoma. The result is the formation of a crescent-shaped collection. The arachnoid remains intact and so blood does not pass into the sulci.

الطب والجراحة لجنة

Subdural Window

- The subdural (blood) window can be used when reviewing a CT brain as it makes intracranial haemorrhage more conspicuous and may help in the detection of thin acute subdural hematomas that are against the calvaria that are similar density to adjacent bone. It is a wider setting than the standard non-contrast window.



Axial non-contrast
Subdural Window

الطب والجراحة لجنة

Subdural Window




Axial non-contrast
Brain Window



Axial non-contrast
Subdural Window

الطب والجراحة لجنة

Subarachnoid Hemorrhage




Hyperdensity in both sylvian and basal cistern (subarachnoid hemorrhage) with small intraventricular hemorrhage in the fourth ventricle

الطب والجراحة لجنة


Subarachnoid hemorrhage

Subarachnoid hemorrhage may result in blood collecting in the ventricles
Occasionally a layer of blood in the ventricles is the only sign



Subarachnoid hemorrhage

Blood within the subarachnoid space is seen in a sulcus of the left cerebral hemisphere. This may be a subtle finding only seen on a few slices



Subarachnoid hemorrhage


- Can occur after trauma, or in individuals with arteriovenous malformations (AVMs), or berry aneurysms (particularly of the Circle of Willis). Patients describe a sudden onset of extremely severe pain ("thunderclap headache"). Blood in the subarachnoid space can cause meningeal irritation and result in symptoms similar to meningitis.

Appearance: High density blood in the sulci, basal cisterns and fissures:

  - Suprasellar cistern (large pentagon/star shape),
  - Quadrigeminal cistern (smaller W/smile shape below suprasellar)
  - Blood may also extend into the ventricles.

الطب والجراحة لجنة

Intraventricular hematoma



Subacute intraventricular hemorrhage



Hypertensive basal ganglial bleed



Basal ganglia hemorrhage with intraventricular extension

الطب والجراحة لجنة

Intraventricular hematoma

- Intraventricular hemorrhage denotes the presence of blood within the cerebral ventricular system.
- It is associated with significant morbidity due to the risk of obstructive hydrocephalus.
- It can be divided into primary or secondary, primary hemorrhage being far less common than secondary:
  - Primary: blood in the ventricles with little (if any) parenchymal blood
  - Secondary: a large extraventricular component is present (e.g., parenchymal or subarachnoid) with secondary extension into the ventricles
- In adults, secondary intraventricular hemorrhage is usually the result of an intracerebral hemorrhage (typically basal ganglia hypertensive hemorrhage) or subarachnoid hemorrhage with ventricular reflux.

الطب والجراحة لجنة

Other Extra Axial Hemorrhages

- **Intralaminar dural haematoma** in the intracranial space, are exceedingly rare, and represent hemorrhage between the two layers of the dura mater. As a result, it has distinctive features allowing it to be distinguished from extradural haematomas but mimics subdural haematoma. They are equivalent to epidural haematoma of the spine.
- **Subpial hemorrhage** is a rare form of extra-axial intracranial hemorrhage defined as hemorrhage between the cortical surface and the pia mater. It is an entity that is generally difficult to distinguish from subarachnoid hemorrhage.

الطب والجراحة لجنة

Intra Axial Hemorrhage

Intra Axial Hemorrhage = Intracerebral hemorrhage (roughly speaking)

- Intracerebral hemorrhage, or intraparenchymal hemorrhage, is a subset of an intracranial hemorrhage and includes a number of entities that have in common the acute accumulation of blood within the parenchyma of the brain

- Intracerebral hemorrhage are usually (but not always - it can be traumatic -) hemorrhagic strokes thus will be discussed in our next part of this section (Strokes)


Cerebral hemorrhagic contusion

- Cerebral hemorrhagic contusions are a type of intracerebral haemorrhage and are common in the setting of significant head injury. They are usually characterized on CT as hyperattenuating foci in the frontal lobes adjacent to the floor of the anterior cranial fossa and in the temporal poles.
- Radiographic features
  - Cerebral contusions can occur anywhere, but have a predilection for certain locations, as a result of the direction of the head strike and the intrinsic shape of the skull cavity.
    - anterior cranial fossa floor
    - temporal pole
    - coup and contrecoup pattern (see next slide)


الطب والجراحة لجنة

Coup and Contrecoup pattern


- A coup-contrecoup injury is a term applied to head injuries and most often cerebral contusions and traumatic subarachnoid haemorrhage. It refers to the common pattern of injury whereby damage is located both at the site of impact (often less marked) and on the opposite side of the head to the point of maximum external trauma.
- The impact accelerates first the skull, at which point the brain immediately subjacent to the point of impact may be damaged as it is forcibly impinged up on by the moving skull – this is sometimes referred to as the coup injury. Shortly thereafter the intracranial content has accelerated to be travelling with the skull, away from the initial impact. As the skull then stops, the brain once again impacts on the internal surface of the skull resulting in damage.
- Areas most frequently affected are the inferior surface of the frontal lobes and temporal poles




Cerebral hemorrhagic contusion



From the case: Cerebral contusions - bifrontal




From the case: Cerebral haemorrhagic contusion



From the case: Cerebral haemorrhagic contusion

Cerebral hemorrhagic contusion



From the case: Cerebral haemorrhagic contusion with subdural and subarachnoid haemorrhage




From the case: Cerebral contusion with acute extravasation of blood



From the case: Coup-contrecoup injury

Diffuse axonal injury (DAI)

- Diffuse axonal injury (DAI), also known as traumatic axonal injury (TAI), is a severe form of traumatic brain injury due to shearing forces. It is a potentially difficult diagnosis to make on imaging alone, especially on CT as the finding can be subtle, however, it has the potential to result in severe neurological impairment.
- The diagnosis is best made on MRI where it is characterized by several small regions of susceptibility artifact at the grey-white matter junction, in the corpus callosum, and in more severe cases in the brainstem (see next slide), surrounded by FLAIR hyperintensity.
- Acceleration, deceleration and rotational forces cause portions of the brain with different densities to move relative to each other resulting in the deformation and tearing of axons



DAI – Intermediary injury

- Intermediary injuries or intermediary contusions affect the basal ganglia and/or thalami and are uncommon manifestations of diffuse axonal injury associated with a poor prognosis.

Radiographic features

- CT
  - hyperdense hemorrhagic contusions of the lentiform nucleus and/or thalami
  - often bilateral

Intermediary injury: Late



From the case: Intermediary injury

Diffuse axonal injury (DAI)

Radiographic features

❖CT

- Unfortunately, it is not sensitive to subtle diffuse axonal injury and as such, some patients with relatively normal CT scans may have significant unexplained neurological deficit
- The appearance depends on whether the lesions are overtly hemorrhagic. Hemorrhagic lesions will be hyperdense and range in size from a few millimeters to a few centimeters in diameter. Non-hemorrhagic lesions are hypodense. They typically become more evident over the first few days as oedema develops around them. They may be associated with significant and disproportionate cerebral swelling.

الطب والجراحة لجنة

Diffuse axonal injury (DAI)

Radiographic features

❖ MRI


- MRI is the modality of choice for assessing suspected diffuse axonal injury even in patients with entirely normal CT of the brain. MRI, especially SWI or GRE sequences, exquisitely sensitive to paramagnetic blood products may demonstrate small regions of susceptibility artefact at the grey-white matter junction, in the corpus callosum or the brain stem. Some lesions may be entirely non-hemorrhagic (even using high field strength SWI sequences). These will, however, be visible as regions of high FLAIR signal.

الطب والجراحة لجنة

Diffuse axonal injury (DAI)



From the case: Diffuse axonal injury



From the case: Diffuse axonal injury

Diffuse axonal injury (DAI)

- With DAI we can see both hemorrhagic and non-hemorrhagic lesions (contusion)
- **Hemorrhagic lesions** will be hyperdense and range in size from a few millimeters to a few centimeters in diameter.
- **Non-hemorrhagic lesions** are hypodense. They typically become more evident over the first few days as oedema develops around them. They may be associated with significant and disproportionate cerebral swelling.

الطب والجراحة لجنة

Secondary brain damage

- Generalized (Diffuse) cerebral oedema
- Hypoxic-ischemic brain injury
- Ischemic stroke from traumatic arterial dissection
- “Big black brain”

الطب والجراحة لجنة

Generalized (Diffuse) cerebral oedema

Radiographic features

- **CT**
  - Diffuse hypodensity
  - White cerebellar sign
  - Effaced sulci and ventricle
  - Loss of grey white matter differentiation



From the case: Diffuse cerebral oedema

White cerebellum sign

- White cerebellum sign, also called reversal sign or dense cerebellum sign, is encountered when there is a diffuse decrease in density of the supratentorial brain parenchyma, with relatively increased attenuation of the thalami, brainstem and cerebellum.
- This sign indicates irreversible brain damage and has a very poor prognosis, which is why radiologists must be aware of this sign.



From the case: Dense cerebellum sign

الطب والجراحة لجنة

Hypoxic-ischemic encephalopathy

- Severe global hypoxic-ischemic injury in this population primarily affects the grey matter structures

Radiographic features

- **CT**
  - diffuse oedema with effacement of the CSF-containing spaces
  - decreased cortical grey matter attenuation with a loss of normal grey-white differentiation
  - decreased bilateral basal ganglia attenuation
  - reversal sign: reversal of the normal CT attenuation of grey and white matter, demonstrated within the first 24 hours in a small number of these patients
  - white cerebellum sign
  - linear hyperdensity outlining the cortex
  - pseudosubarachnoid haemorrhage

الطب والجراحة لجنة

“Big black brain”

- is a radiologic pattern unique to infants and toddlers in the context of traumatic brain injury, most frequently due to abusive head trauma.
- The CT presentation is a parenchymal hypodensity and uniform loss of gray-white differentiation in the entire hemisphere, associated with ipsilateral subdural haematoma.
- The pattern can appear in the vascular territories of the ACA, MCA and PCA, without evidence of occluded blood vessels.
- Although "black" may refer to hypodensity on CT, the pathology is apparent on MRI sequences, which will show signal changes, as well.
- Frequently, there is subsequent hemispheric atrophy, resulting in severe motor impairment, cognitive impairment, epilepsy, and blindness.

الطب والجراحة لجنة


Long-term sequelae of head trauma

- Encephalomalacia/Gliosis (discussed later in strokes)
- Chronic subdural haematomas / CSF Hygromas
- Chronic traumatic encephalopathy (a neurodegenerative tauopathy that is thought to result from mild repetitive head trauma).
- Depression, anxiety and alcohol excess
- Increased risk of schizophrenia, bipolar disorder and organic mental disorders

الطب والجراحة لجنة

Subdural hygroma


- Subdural hygromas refer to the accumulation of fluid in the subdural space. In many cases, it is considered an epiphenomenon of head injury when it is called a traumatic subdural hygroma.
- Subdural hygromas generally occur along the supratentorial cerebral convexities
- Most subdural hygromas are believed to be derived from chronic subdural hematomas.



Subdural hygroma

CT/MRI

- A subdural hygroma radiographically appears as a crescentic near-CSF density/signal accumulation in the subdural space that does not extend into the sulci and rarely exerts significant mass-effect.
- Vessels rarely cross through the lesion in contrast-enhanced studies (cortical vein sign). Importantly these collections do not entirely follow CSF on FLAIR, often appearing hyperintense.



الطب والجراحة لجنة

Subdural hygroma



From the case: Subdural hygroma



From the case: Bilateral subdural hygroma

الطب والجراحة لجنة

Complications of Traumatic Brain Injuries

Severe mass effect can result in:

- Midline shift: associated with worse prognosis
- Cerebral herniation: often requires urgent treatment
- Hydrocephalus: can also be a chronic non-mass effect related complication

- Midline shift and hydrocephalus are discussed later

الطب والجراحة لجنة


Associations


Other traumatic injuries are common:

- Cervical spine injury
- Skull fracture
- Facial fracture
- Pneumocephalus
- Extracranial injuries in 35%

الطب والجراحة لجنة

Compressed Fracture





From the case: Vertebral body compression fracture

Compressed Fracture



X-Ray



CT Sagittal reformat

الطب والجراحة لجنة

Skull Fracture – Bone window








Depressed / Displace Fracture

الطب والجراحة لجنة

Skull Fracture – Bone window



Nondisplaced



Linear

الطب والجراحة لجنة

Pneumocephalus

- Refers to the presence of intracranial gas, and in most cases the gas is air. The term encompasses gas in any of the intracranial compartments and is most encountered following trauma or surgery.

Radiographic features

CT

- Gas on CT will have a very low density (~ -1000HU) but care needs to be taken in ensuring that it is not fat which although of much higher density (-90HU) also appear completely black on routine brain windows.

MRI


- On MRI the diagnosis may be more difficult as there is no 'objective' density measurement. Gas will appear completely black on all sequences, but depending on the location and morphology can be mistaken for blood product or flow voids

الطب والجراحة لجنة

Pneumocephalus



From the case: Pneumocephalus following sinus surgery



From the case: Pneumocephalus



From the case: Traumatic pneumocranium

Strokes

Section3: Pathological Conditions

Stroke

- A stroke is a clinical diagnosis that refers to a sudden onset focal neurological deficit of presumed vascular origin.
- It is divided into two broad categories:
  1. hemorrhagic stroke (15%)
  2. ischemic stroke (80%)
- Hemorrhage may be primary (usually as a result of hypertension or cerebral amyloid angiopathy) or secondary (the result of an underlying lesion such as a vascular malformation or tumor).

الطب والجراحة لجنة

Hemorrhagic Stroke (Intracerebral Hemorrhage)

They are most often broadly divided according to whether they are spontaneous (primary) or due to an underlying lesion (secondary), and then further divided according to aetiology and/or location.

1. **primary haemorrhages** (no underlying lesion)
    1. Lobar haemorrhages secondary to cerebral amyloid angiopathy
    2. Hypertensive haemorrhages
2. **secondary haemorrhages** (some other lesion complicated by hemorrhage)
    1. Vascular malformation
    2. Cerebral venous thrombosis
    3. Tumor (primary or secondary)

الطب والجراحة لجنة

Primary Hemorrhages

- Primary hemorrhages are subclassified into 2 subtypes and tend to have favorable location, unlike secondary hemorrhages that can acquire anywhere.
  - Lobar hemorrhage secondary to cerebral amyloid angiopathy
  - Hypertensive hemorrhage secondary to uncontrolled hypertension
- According to the location of the hemorrhage
  - Cerebral amyloid angiopathy → Lobes
  - Hypertensive hemorrhage → Basal ganglia, Pontine, and Cerebellum

الطب والجراحة لجنة


Lobar haemorrhage

- **Aetiology**: most commonly due to cerebral amyloid angiopathy especially in the elderly; note that the confirm of this diagnosis needs a brain autopsy thus clinically we search and exclude secondary causes especially in younger individuals
- **Key point**: cerebral amyloid angiopathy microhemorrhages occur in the same distribution as lobar haemorrhages, thus in older adults with a known history of past cerebral amyloid angiopathy microhemorrhages we don't need to exclude 2ry causes
- **Location**: its usually peripherally located
- **Can extended** commonly to the subdural space and less commonly if the hemorrhage is massive, it can extend to the intraventricular space

الطب والجراحة لجنة

Lobar Haemorrhage Case

Presentation: 70-year-old woman, 3 days worsening headache. Sudden onset, associated vomiting and confusion.



This is a calcified cordial plexus NOT a hemorrhage
A normal finding especially in older individuals

الطب والجراحة لجنة


Lobar Haemorrhage Case

Presentation: 70-year-old woman, 3 days worsening headache. Sudden onset, associated vomiting and confusion.

- MRI (days later)

Presented with a new lobar hemorrhage.

Also notice the microaneurysms through out the peripheral field




Hypertensive hemorrhage

- Hypertension is the most common cause of intracerebral haemorrhages.
- **Aetiology:** Most commonly due to rupture of Charcot-Bouchard aneurysms in lenticulostriate arteries 80%, pons 10%, cerebellum 10%
- Divided according to their typical **locations** (centrally located) which include, in order of frequency:
  1. Basal ganglia haemorrhage (especially the putamen)
  2. Thalamic haemorrhage
  3. Pontine haemorrhage
  4. Cerebellar haemorrhage
- **Can extended** to the ventricles causing secondary intraventricular hemorrhages resulting in hydrocephalus

الطب والجراحة لجنة

Hypertensive hemorrhage

- This diagram depicts the Charcot Bouchard microaneurysms arising from perforating lenticulostriate arteries.
- These aneurysms typically arise from arteries no larger than 300 micrometers in diameter. Thus, can’t be seen on imaging




Hypertensive hemorrhage

CT Axial



Hypertensive basal ganglial bleed

Axial FLAIR



Thalamic haematoma and intraventricular haemorrhage

CT Axial



Cerebellar haemorrhage

الطب والجراحة لجنة

Lobar Vs Hypertensive hemorrhages

	Lobar	Hypertensive
Aetiology	Cerebral amyloid angiopathy	Charcot-Bouchard aneurysm rupture
Location	Lobes Aka peripherally located	Deep gray matter structures Aka centrally located
Subdural extension	Can extend to the subdural space	-
2ry intraventricular hemorrhage	Massive lobar hemorrhage can extend to the intraventricular space causing secondary intraventricular hemorrhage	can extend to the intraventricular space causing secondary intraventricular hemorrhage resulting in hydrocephalus

الطب والجراحة لجنة

Ischemic strokes

- **Ischemic stroke** results from a sudden cessation of adequate amounts of blood reaching parts of the brain and typically presents with rapid onset neurological deficit, which is determined by the area of the brain that is involved.. Ischemic strokes can be **divided according to** territory affected or mechanism.
- Ageing ischemic strokes can be important in a number of clinical and medicolegal settings. Both CT and MRI can help in determining when a stroke occurred as imaging features evolve in a reasonably predictable fashion.
- CT scan is the most modality we use to diagnose acute strokes; more widely available and less time consuming than MRI

في اختلاف بين المراجع على الageing تاع الstrokes بالسلايد التالي حنكتب الي رح نعتمده في هذه الدوسية مع شوية تطرق لرأي ثاني في التصنيف معلش عيني تحملنا شوي بالفلسفة

الطب والجراحة الجنة

Ischemic strokes Ageing

- Early hyperacute: 0 to 6 hours
- Late hyperacute: 6 to 24 hours
- Acute: 24 hours to 1 week
- Subacute: 1 to 3 weeks
- Chronic: more than 3 weeks

The above definition of hyperacute as 0-24 hours and acute as 1-7 days was affirmed by the international Stroke Recovery and Rehabilitation Roundtable. However, this group defined subacute as 1 week to 6 months (with 3 months dividing early and late subacute phases) and chronic as older than 6 months.


الطب والجراحة لجنة

Ischemic strokes Evolution

- **During few hours (Hyperacute)**: Subtle loss of gray-white matter differentiation (see next slides for more info about hyperacute infarct findings)
- **First week (Acute)**: Region become more defined and lower attenuation & swells. In the middle of the first week there is maximal swelling.
- **2-3 weeks (Subacute)**: Swelling gradually subsides and attenuation increases, in some patient returning to near normal Aka fogging phenomenon (discussed latter)
- **More than 3 weeks (Chronic)**: The infarcted tissue is removed leaving behind cystic encephalomalacia

الطب والجراحة لجنة

Ischemic strokes Evolution



Click here to view the video
Stroke: Evolution from acute to chronic infarction - radiology video tutorial (CT, MRI)

Ischemic strokes Evolution

Days – Weeks	Months – Years
Less well demarcated	Very well demarcated
Positive mass effect	Negative mass effect

الطب والجراحة لجنة

Ischemic strokes Evolution

In the next slides we will discuss

- The hyperacute findings of a stroke
- Fogging phenomenon
- Hemorrhagic transformation of ischemic
- Encephalomalacia

الطب والجراحة
لجنة

Hyperacute findings

- Visualization of clot
  - Hyperdense artery sign
    - refers to focal hyperdensity of arteries on non-contrast brain
    - Can be seen in thromboembolisms but not acute thrombus that has been formed on top of atherosclerotic plaque this is due to the fact the clot increase in density in time (emboli are developed clots so appear hyperdense while an acute thrombus appear Isodense or mildly hyperdense)
- Early parenchymal signs
  - When ischemic the gray matter become hypodense, and the gray matter differentiation become lost
    - Loss of the insular ribbon sign

الطب والجراحة لجنة

Hyperdense vessel sign

- Hyperdense MCA sign
  - refers to focal hyperdensity of the middle cerebral artery (MCA) on non-contrast brain CT and is the direct visualization of thromboembolic material within the lumen.
  - It is thus the earliest visible sign of MCA infarction as it is seen within 90 minutes after the event.
  - It is the longitudinal equivalent of the MCA dot sign and hyperdense basilar tip sign.




Hyperdense vessel sign

- MCA dot sign
  - The middle cerebral artery (MCA) dot sign, also known as the Sylvian fissure sign, is seen on non-contrast brain CT and represents the cross-sectional M2 equivalent of the hyperdense MCA sign.
  - Rather than imaging a length of middle cerebral artery (typically the M1 segment), the dot sign represents a thromboembolism within a segmental branch of the MCA located within the Sylvian fissure (M2 segment).
  - The sign appears when the high-attenuation MCA is viewed in axial section, since the occluded vessel courses in a plane perpendicular to the transverse plane of imaging. (consider slice thickness and the orientation of the vessel when determining artery hyperdensity; use sagittal and coronal images to problem solve)

MCA dot sign


Case link




In cases like this where the hyperdensity is not seen will or you have some doubts about the diagnosis check the sagittal and coronal views for confirmation

Hyperdense vessel sign

- Hyperdense basilar artery
  - It is a sign that can indicates an acute occlusion of the basilar artery which may result in brainstem or thalamic ischemia or infarction






Posterior circulation infarction due to basilar artery occlusion


Loss of the insular ribbon sign

- The loss of the insular ribbon sign refers to a loss of definition of the gray-white interface in the lateral margin of the insular cortex ("insular ribbon") and is considered an early CT sign of MCA infarction.
- The insular cortex is more susceptible to ischemia following MCA occlusion than other portions of the MCA territory because it has the least potential for collateral supply from the anterior cerebral and posterior cerebral arteries.
- Best appreciated on stroke window




Stroke Window

Loss of the insular ribbon sign



Brain Window



Stroke Window

Fogging phenomenon

- The fogging phenomenon is seen on non-contrast CT of the brain and represents a transient phase of the evolution of cerebral infarct where the region of cortical infarction regains a near-normal appearance.
- During the first week following a cortical infarct, hypoattenuation and swelling become more marked, resulting in significant mass effect and clear demarcation of the infarct often with vivid gyral enhancement. As time goes on, the swelling starts to subside, and the cortex begins to increase in attenuation.
- At 2 to 3 weeks following an infarct, the cortex regains near-normal density and imaging at this time can lead to confusion or missed diagnosis. Fogging has been demonstrated in around 50% of cases.
- If in doubt, the administration of IV contrast will demarcate the region of infarction.
- A similar phenomenon is also seen on T2 MRI sequences of the brain and is believed to be due to similar cellular processes, as the timing is similar. It has been found to occur in approximately 50% of patients between 6 and 36 days (median 10 days) after the onset of infarction.

الطب والجراحة لجنة

Hemorrhagic Transformation (Hemorrhagic Infarction)

- It occur during the first week in more than 50% of infarcts but rarely seen during the first 6 hours
- The term hemorrhagic transformation is somewhat variably used and collectively refers to two different processes, which have different incidence, appearance and prognostic implications. These are:
  1. hemorrhagic infarction (petechial haemorrhages)
  2. parenchymal haematoma

الطب والجراحة لجنة

Petechial Haemorrhages Vs Parenchymal Haematoma

- Petechial haemorrhages, as the name suggests, usually appear as tiny punctate regions of haemorrhage, often not able to be individually resolved, but rather resulting in increased attenuation of the region on CT or signal loss on MRI. Although this petechial change can result in cortex appearing near-normal it should not be confused with the phenomenon of fogging seen on CT which occurs 2 to 3 weeks after infarction. It is associated with no mass effect and rarely affect prognosis.
- In the case of secondary haematoma, the radiographic features on both CT and MRI are merely a summation of the features of an ischemic infarct, with superimposed cerebral haemorrhage. The amount of haemorrhage relative to the size of the infarct can vary widely, but usually, it is possible to identify significant areas of the brain which are infarcted but not hemorrhagic. This may not be the case if the haemorrhage is large and the underlying infarct small. It is associated with mass effect that varies according to its size relatively to infarct and have poor prognosis
- Reperfusion therapy increase the risk of secondary haematoma

الطب والجراحة لجنة

Encephalomalacia

- Encephalomalacia is frequently used by radiologists to describe any area of cerebral parenchymal loss with or without surrounding gliosis.
- Encephalomalacia is the end result of liquefactive necrosis of brain parenchyma following insult, usually occurring after cerebral ischemia, cerebral infection, haemorrhage, traumatic brain injury, surgery or other insults.
- **Radiographic features**

CT

- hypoattenuation, somewhat, higher than CSF
- volume loss
- often associated with gliosis and Wallerian degeneration

MRI

Follows CSF signal on all sequences including FLAIR.

- T1: low signal
- T2: high signal, attenuating fully on FLAIR
- ADC: facilitated diffusion

الطب والجراحة لجنة

Ischemic strokes according to affected territory

- The vascular territory affected will determine the exact symptoms and clinical behavior of the lesion:
  - Anterior circulation infarct
    - Anterior cerebral artery infarct
    - Middle cerebral artery infarct
    - Lacunar infarct
    - Sriatocapsular infarct
  - Posterior circulation infarct
    - Posterior cerebral artery infarct
    - Cerebellar infarct
    - Brainstem infarct (midbrain, pons, medulla)


الطب والجراحة لجنة

Ischemic strokes according to affected territory


- **Anterior cerebral artery (ACA) territory infarcts** are much less common than either middle or posterior cerebral artery territory infarcts.
- **The middle cerebral artery territory** is the most commonly affected territory in a cerebral infarction, due to the size of the territory and the direct flow from internal carotid artery into the middle cerebral artery, providing the easiest path for thromboembolism.
- **Posterior circulation infarction (POCI),** also referred as posterior circulation stroke, corresponds to any infarction occurring within the vertebrobasilar vascular territory, which includes the brainstem, cerebellum, midbrain, thalami, and areas of temporal and occipital lobes

الطب والجراحة لجنة


Remember: Cerebral Blood Supply




Ischemic strokes according to affected territory



ACA Infarction



MCA Infarction



PCA Infarction

الطب والجراحة
الجنة


Ischemic strokes according to affected territory

Thalamic lacunar infarct



Lacunar infarction: small infarction less than 1.5 cm usually around basothalamus occure mainly in DM and hypertensive patients

Cerebellar infarction



Pay attention while differentiating between cerebellar infarcts and PCA infarcts

الطب والجراحة لجنة

Intracranial Masses

Section3: Pathological Conditions

Intracranial Masses

In this part of our dossier, we will talk only about selected cases as if this topic is very wide

- Extra-axial masses
  - Meningiomas
- Intra-axial masses
  - Low grade gliomas
  - High grade gliomas
  - Abscess

الطب والجراحة لجنة


Extra-axial Vs Intra-axial

Often it is trivially easy to distinguish an intra-axial from an extra-axial mass. In many cases, especially when the mass is large and associated with parenchymal changes, such as oedema, localization can be more difficult. Several features are helpful in suggesting that a mass or lesion is extra-axial, including:

- subarachnoid space
  - CSF cleft sign
  - widening of adjacent subarachnoid space/cistern\
  - intervening pial arteries or veins
- brain parenchyma
  - absence of a claw sign
  - intervening cortex between mass and white matter
  - white matter buckling sign
- bone and meninges
  - dural tail sign
  - erosion, invasion or destruction of adjacent bone
  - hyperostosis


الطب والجراحة لجنة

Extra-axial Vs Intra-axial




Classically, the cleft was regarded as representing a thin rim of CSF between a tumor and brain parenchyma.


Claw sign refers to the sharp angles on either side of the mass, which the surrounding normal parenchyma forms when the mass has arisen from the parenchyma. As such, normal parenchyma extends some way around the mass.



Extra-axial Vs Intra-axial



white matter buckling sign represents the white matter projecting into gyri being compressed and displaced by the mass, even in the presence of oedema (which would usually expand gyri, if the mass were intra-axial)



The **dural tail sign** occurs as a result of thickening and enhancement of the dura and is most often seen adjacent to a meningioma.

Meningiomas

- Meningiomas are extra-axial tumors and represent the most common tumor of the meninges.

Radiographic features

- CT is often the first modality employed to investigate neurological signs or symptoms, and often is the modality which detects an incidental lesion:
  - Non-contrast CT
    - 60% slightly hyperdense to normal brain, the rest are more isodense
    - 20-30% have some calcification
  - Post-contrast CT
    - 72% brightly and homogeneously contrast enhance
    - malignant or cystic variants demonstrate more heterogeneity/less intense enhancement


الطب والجراحة لجنة

Meningioma Vs Intra-axial tumors


How to differentiate between the meningioma and Intra-axial tumors ?

1. Meningioma present with extra-axial features:
   - CSF cleft sign
   - intervening pial arteries or veins
   - Absence of a claw sign
   - white matter buckling sign
   - dural tail sign
2. They grow at one spot and bring their blood supply with them giving them sunburst or spoke-wheel appearance of the vessels

Meningioma



Spoke wheel sign



Sunburst sign

الطب والجراحة لجنة

Astrocytoma (Low grade glioma)

CT


- Typically, low-grade infiltrating astrocytoma's appear as isodense or hypodense regions of positive mass effect, often without any enhancement (in fact, the presence of enhancement would suggest higher grade tumors)

MRI

- T1: isointense to hypointense compared to white matter
- T2: mass-like hyperintense signal
- T1 C+ (Gd):
  - No enhancement in grade 2 tumors
  - Solid areas of enhancement +/- necrosis suggest higher grade

الطب والجراحة لجنة

Astrocytoma (Low grade glioma) MRI



T1: hypointense


T2: hyperintense

T1 C+ (Gd): No enhancement

الطب والجراحة لجنة


Glioblastomas (High grade glioma)

- Glioblastomas are typically large tumors at diagnosis. They often have thick, irregular-enhancing margins and a central necrotic core, which may also have a hemorrhagic component.
- They are surrounded by vasogenic-type oedema, which in fact usually contains infiltration by neoplastic cells.
- They can cross the midline giving them butterfly appearance




From the case: Glioblastoma NOS (butterfly morphology)

Glioblastomas



From the case: Glioblastoma NOS



From the case: Glioblastoma NOS

Cerebral Abscess (Brain Abscess)

- **Cerebral abscesses** represent focal areas of infection within brain parenchyma, usually containing pus and having a thick capsule.
- It is a potentially life-threatening condition requiring prompt radiological identification and rapid treatment.
- They typically have enhancing walls and can mimic several other significant pathologies (e.g., GBM and metastasis)
- Fortunately, MRI is usually able to convincingly make the diagnosis, distinguishing abscesses from other ring-enhancing lesions.

الطب والجراحة لجنة


Cerebral Abscess (Brain Abscess)

How to differentiate between cerebral abscess and (GBM and metastasis) ?

- Pattern of enhancement in abscess are usually thin & regular rim (not very helpful as abscess rim can be irregular or tumor rim to be regular)
- The dual or double rim sign is seen on MRI in approximately 75% of cerebral abscesses and is helpful in distinguishing an abscess from a glioblastoma. On both SWI and T2WI
- Diffuse restriction (DWI) is your best clue where abscess present with high DWI signal centrally

الطب والجراحة لجنة

Cerebral Abscess (Brain Abscess)



Miscellaneous Topics

Section3: Pathological Conditions

Miscellaneous Topics

- Multiple Sclerosis
- Brain Atrophic Changes
- Venous Sinus Thrombosis
- Small Vessel Disease

الطب والجراحة
الجنة

Multiple Sclerosis

- Multiple sclerosis (MS) is a chronic demyelinating condition affecting the central nervous system. Diagnosis requires good history, clinical examination, appropriate imaging, and laboratory tests (cerebrospinal fluid for IgG index and oligoclonal bands).
- It is typified by "plaques" of disease which are spread by location and time. MRI is frequently the first-line imaging modality for diagnosis and follow-up.
- Associated with white matter demyelination which is visible on MRI as plaques of disease

Investigation

- CT It is insensitive, although some plaques may show contrast enhancement in the active phase
- **MRI** is the investigation of choice, should be performed with contrast and include the spine

الطب والجراحة لجنة


Multiple Sclerosis – MRI

- Preferred imaging modality for diagnosis and follow up
- Best visualized on a fluid-sensitive sequence, e.g. T2, FLAIR
- White matter lesions; characteristic periventricular, they tend to extend perpendicularly and on sagittal views are called "Dawson's fingers
- Lesions may occur throughout the CNS and be found in a variety of locations, e.g., juxtacortical, brainstem, cerebellum, or spinal cord
- Enhancement is a marker of activity and can be used to determine lesion load and disease progression

الطب والجراحة لجنة

Multiple Sclerosis – MRI – Case 1

- Multiple white matter plaques, some of them are perpendicular to the corpus callosum - so-called Dawson fingers - are typical of multiple sclerosis. On some slices (see next slide), subcallosal dot-dash sign is also evident.




From the case: Multiple sclerosis

الطب والجراحة لجنة

Multiple Sclerosis – MRI – Case 1

- The ependymal dot-dash sign has been described as an early MRI imaging feature of multiple sclerosis before other more florid white matter changes (e.g., Dawson's fingers) become evident
- The sign is best seen on sagittal FLAIR along the inferior surface of the corpus callosum and roof of the lateral ventricle bodies. It is seen as tiny (~1 mm) dots of high signal along the ependymal surface that may coalesce into short dashes.




From the case: Multiple sclerosis


الطب والجراحة لجنة

Multiple Sclerosis – MRI – Case 2


T1 weighted



FLAIR



T2 weighted




FLAIR Sagittal




Multiple scattered supra and infratentorial white matter plaques, most of them are seen at the periventricular region perpendicular to the lateral ventricles giving the appearance of Dawson fingers, other plaques involve the subcortical U fibers, juxta cortical and brainstem. Most of the lesions show T2 shine through effect on DWI/ADC. Characteristically they display low signal on T1 WIs.

الطب والجراحة لجنة


Multiple Sclerosis – MRI – Case 3



T2 weighted



T1 with contrast



T2 weighted

From the case: Spinal cord expansion in multiple sclerosis

Multiple Sclerosis – MRI – Case 4


Enhancement of plague post contrast T1



T1 with Gad
subtle enhancement



DWI acute restricted
diffusion




Matching FLAIR
abnormality

From the case: Multiple sclerosis

Brain atrophic changes

- CT and MRI are equally able to demonstrate cortical atrophy, but MRI is more sensitive in detecting focal atrophic changes in the nuclei.
- Characteristic features include prominent cerebral sulci (i.e., cortical atrophy) and ventriculomegaly (i.e., central atrophy) without bulging of the third ventricular recesses.



Alzheimer disease

❖ ما اتوقعش حد بعرفش شو الزهايمر فرح ندخل بالCT والMRI مباشرة

Radiographic features

❖Although CT is able to demonstrate the characteristic patterns of cortical atrophy, MRI is more sensitive to these changes and better able to exclude other causes of dementia (e.g., multi-infarct dementia) and as such is the favored modality.

❖**Braak staging** on imaging shows the sequential involvement of various structures as follows:


- Stage I and II : earliest involvement of entorhinal cortex.
- Stage III and IV: involves limbic system and hippocampus.
- Stage V and VI: involves cortex (precuneus in particular) with temporal lobes

الطب والجراحة لجنة

Alzheimer disease – MRI


- The primary role of MRI (and CT for that matter) in the diagnosis of Alzheimer disease is the assessment of volume change in characteristic locations, such volume loss is not apparent early in the course of the disease.
- The diagnosis should be made on the basis of two features:

1. Mesial temporal lobe atrophy (particularly the hippocampus, entorhinal cortex and perirhinal cortex)
2. Temporoparietal cortical atrophy



Alzheimer disease – MRI

The sulci and ventricles are mildly more prominent than expected for age. The mesiotemporal structures, including the temporal horns of the lateral ventricles and the collateral sulci, are slightly more prominent, with bilateral hippocampus, amygdala and entorhinal cortex reduced in volume.



In Alzheimer disease we look for atrophic changes in these particular areas, in this case no prominent atrophy can be seen Aka it's not a late stage (stage 3 or 4)

Dural Venous Sinus Thrombosis

- Dural venous sinus thrombosis is a subset of cerebral venous thrombosis, often coexisting with cortical or deep vein thrombosis, and presenting in similar fashions, depending mainly on which sinus is involved.
- Presentation is variable and can range from asymptomatic to coma and death. Typically, patients complain of headache, nausea, and vomiting. Neurological deficits are variable.

Radiographic features

- Unenhanced CT is usually the first imaging investigation performed given the nonspecific clinical presentation in these cases. Potential findings include cord sign, dense vein sign, vasogenic oedema - discused later - secondary to venous hypertension and unilateral or bilateral cortical or peripheral venous haemorrhage
- With contrast administration, especially with a CT venogram, a sinus filling defect is sought. Signs on contrast CT include empty delta sign, gyral enhancement, and prominent intramedullary vein

الطب والجراحة لجنة

Dural Venous Sinus Thrombosis

Unenhanced CT


- The **cord sign** refers to cordlike hyperattenuation within a dural venous sinus on non-contrast enhanced CT of the brain due to dural venous sinus thrombosis. The sign is most commonly seen in the transverse sinus because along the origin of the tentorium it runs approximately in the axial plane such that it is visible on one image.



Dural Venous Sinus Thrombosis

Unenhanced CT

- The **dense vein sign** refers to hyperattenuating thrombus within a cortical vein or dural venous sinus due to acute venous thrombosis.
- When located in the superior sagittal sinus, particularly posteriorly, it is sometimes referred to as the delta, triangle or pseudodelta sign. It is really the same as the cord sign except the vessel is imaged in cross-section.




الطب والجراحة لجنة

Dural Venous Sinus Thrombosis

Enhanced CT

- The **empty delta sign** is a CT sign of dural venous sinus thrombosis of the superior sagittal sinus, where contrast outlines a triangular filling defect, which represents thrombus. It is only described in contrast-enhanced CT, not in unenhanced scans.
- An equivalent appearance can be noted in the transverse sinus on sagittal or coronal planes.



الطب والجراحة لجنة

Cerebral Small Vessel Disease

- It is an umbrella term for lesions in the subcortical brain attributed to pathologic changes in the small vessels. It is the most common cause of vascular dementia/cognitive impairment and is a major cause of ischemic and hemorrhagic strokes.
- **Aetiology**: (most common causes)
  - Age-related
  - Hypertension-related small vessel diseases
  - Cerebral amyloid angiopathy
- **Radiographic features**
  - CT
    - non-enhancing white matter hypodensities
  - MRI
    - T1: hypointense or isointense, less conspicuous than on T2/FLAIR
    - T2/FLAIR: hyperintense
    - DWI: no diffusion restriction
    - T1 C+ (Gd): non-enhancing


الطب والجراحة لجنة

Cerebral Small Vessel Disease

The Fazekas scale is used to simply quantify the amount of white matter T2 hyperintense lesions usually attributed to chronic small vessel ischemia

The scale divides the white matter in periventricular and deep white matter, and each region is given a grade depending on the size and confluence of lesions

- periventricular white matter (PVWM)
  - 0 = absent
  - 1 = “caps” or pencil-thin lining
  - 2 = smooth “halo”
  - 3 = irregular periventricular signal extending into the deep white matter
- deep white matter (DWM)
  - 0 = absent
  - 1 = punctate foci
  - 2 = beginning confluence
  - 3 = large confluent areas



Section4: Important key findings

Important key findings

- Midline shift
- Mass effect
- Hydrocephalus
- Brain mass lesion (Focal brain lesion)
- Cerebral oedema
- Intracranial haemorrhage
- Extra-axial collection

الطب والجراحة لجنة

Midline shift

- **Midline shift** is a finding described on transverse (axial) slices from CT and MRI studies. It describes the situation where the midline of the intracranial anatomy is no longer in the midline and is the result of pushing or pulling forces within either side of the intracranial compartment.

- midline structures
  - Falx cerebri
  - Septum pellucidum
  - 3rd ventricle
  - Interpeduncular cistern
  - 4th ventricle



Midline shift

Role of imaging

- Is there midline shift?
- If it was present previously, is it worse?
- What is the likely cause?
- Is there an acute cause that needs an immediate intervention?
- Who needs to know now?

الطب والجراحة لجنة

Mass effect

- Intracranial mass effect describes what happens around a focal lesion like tumor in the brain. It is important to make the distinction between an abnormality that causes mass effect and compresses adjacent structures, and one that does not. Most tumors will cause mass effect on surrounding structures and in turn cause midline shift or hydrocephalus



Mass effect

Role of imaging


- Quantify the degree of mass effect
- What is the likely cause?
- Are any complications of mass effect, e.G hydrocephalus?

الطب والجراحة لجنة

Hydrocephalus

- **Hydrocephalus** describes the situation where the intracranial ventricular system is enlarged because of increased pressure.
- Radiographic features
  - Dilatation of the ventricular system
    - Lateral ventricles: temporal horns dilate first
  - Transependymal CSF shift
    - Periventricular oedema
    - Reduced density on CT
    - High signal on fluid-sensitive MRI sequences (T2, FLAIR)

الطب والجراحة لجنة


Enlarged ventrilces
Communicating
Non-communicating
No obstruction to absorption of CSF
Absorption of CSF obstructed
Obstruction to normal absorption of CSF thus always present
Special types
Communicating hydrocephalus
Obstructive hydrocephalus

Hydrocephalus

Role of imaging

- Is there hydrocephalus?
- Is it stable or progressive? (Pre-existing hydrocephalus)
- Obstructive or non-obstructive?
  - Level of obstruction?
  - Likely cause? E.G. Tumour, stroke
- Transependymal CSF spread?
- Complications?

الطب والجراحة لجنة

Hydrocephalus





Brain mass lesion (Focal brain lesion)

They are a very disparate group of conditions ranging from infection (abscess) to brain tumors

Role of imaging

- ❖Is there a mass lesion?
- ❖What's the likely cause?
- ❖Are there complications (e.g., Mass effect)?
- ❖Will further imaging (or other tests) be helpful?

When assessing a brain lesion, use these questions:


- ❖Is the lesion solitary?
- ❖Is the lesion definitely in the brain (intra-axial)?
- ❖Is there oedema around the lesion?
- ❖Is there associated mass effect?
- ❖Is there clinical history or previous imaging that helps in the diagnosis, e.g., History of cancer

الطب والجراحة لجنة


Brain mass lesion examples



From the case: Haemorrhagic brain metastases



From the case: Cerebral abscess



From the case: Glioblastoma NOS

الطب والجراحة لجنة

Cerebral oedema

❖Cerebral oedema describes the collection of additional fluid within the white matter of the brain. It is the brain's response to an insult and may take one of two broad forms: vasogenic (white matter) and cytotoxic (grey and white matter) oedema.

❖ **Role of imaging**

- Confirm the presence of oedema
- Determine the likely underlying cause
- Determine whether there is sequela from any mass effect

❖ **Radiographic features**

- Compression and displacement of adjacent structures
- Change in parencymal density (CT) and signal (MRI)
- Reduced density (attenuation) on CT
- High-signal on t2-weighted sequences (MRI)
- Low-signal on t1-weighted sequences (MRI)

الطب والجراحة لجنة

Cerebral oedema

- It has traditionally been broadly divided into vasogenic cerebral oedema and cytotoxic cerebral oedema
- As such a more precise classification is:
  - Cytotoxic cerebral oedema
    - Ionic cerebral oedema
  - Vasogenic cerebral oedema
  - Hemorrhagic transformation
- Special types of oedema to be considered:
  - Transependymal oedema (also known as interstitial cerebral oedema) (discussed in hydrocephalus)
  - Combined cerebral oedema (a type of cerebral oedema in which there is a mixed pattern of both cytotoxic and vasogenic cerebral oedema.)

الطب والجراحة لجنة

Vasogenic cerebral oedema

- Vasogenic cerebral oedema refers to a type of cerebral oedema in which the blood brain barrier (BBB) is disrupted (compared to cytotoxic cerebral oedema, where the blood-brain barrier remains intact). It is an extracellular oedema which mainly affects the white matter via leakage of fluid from capillaries.
- It is most frequently seen around brain tumors (both primary and secondary) and cerebral abscesses, though some vasogenic oedema may be seen around maturing cerebral contusion and cerebral haemorrhage.


الطب والجراحة لجنة

Vasogenic cerebral oedema

Radiographic features

❖CT

- grey-white matter differentiation is maintained, and the oedema involves mainly white matter, extending in a finger-like fashion
- secondary effects of vasogenic oedema are similar to cytotoxic oedema, with effacement of cerebral sulci, with or without midline shift



From the case: Brain metastases from breast cancer


ملاحظة حلو زي عيونك عيني انه مرات الlesion ما ببين نفسه على الnon contrast والشيء الوحيد الي ببين هو أثره زي مثلا في هذه الحالة الedema

Vasogenic cerebral oedema

Radiographic features

❖MRI

- hyperintense T2 and FLAIR signals, which do not show restricted diffusion (compared to cytotoxic cerebral oedema, which shows diffusion restriction)



From the case: Brain metastasis (large cystic mass)

الطب والجراحة لجنة

Cytotoxic cerebral oedema

- Cytotoxic cerebral oedema refers to a type of cerebral oedema, most seen in cerebral ischemia, in which extracellular water passes into cells, resulting in their swelling.
- The term is frequently used in clinical practice to denote the combination of true cytotoxic oedema and ionic cerebral oedema. As the pathophysiology of these two types of oedema is different, as is their imaging, they are discussed separately.
- Ionic cerebral oedema is a form of cerebral oedema usually associated with cytotoxic oedema and represents the passage of water and sodium from capillaries into the brain parenchymal extracellular space. It is distinguished from vasogenic oedema as the blood brain barrier (BBB) remains intact and the oedema occurs as a result of cellular damage, often from ischemia.

الطب والجراحة لجنة

True Cytotoxic Cerebral Oedema

Radiographic features

❖CT


- In true isolated cytotoxic oedema little change is evident on CT as a mere redistribution of water from extracellular to intracellular compartments does not result in attenuation changes. The changes colloquially ascribed to 'cytotoxic oedema' are in fact mostly due to ionic oedema. This is why brain CT is often normal in patients with an acute ischemic stroke.

❖MRI


- The one sequence which is able to identify cytotoxic oedema is diffusion weighted imaging (DWI). As cells swell due to inward shift of water, there is a commensurate decrease in diffusion, identified as high signal on DWI and low signal on ADC.
- These changes persist into the subacute phase until about two weeks when the ADC signal begins to rise above the normal parenchyma and eventually becomes hyperintense.

الطب والجراحة لجنة

True Cytotoxic Cerebral Oedema



From the case: MCA stroke - evolution



From the case: Infarction in an eloquent area

Intracranial haemorrhage

Discussed earlier in details here we'll talk just about the interpretation

Role of imaging

- Is there any haemorrhage?
- Where is it?
- Is the cause visible on the scan, e.g., Fracture, aneurysm)?
- Are there any complications, e.g., Mass effect, midline shift?


Radiographic features

- Acute haemorrhage is hyperdense (white) on ct
- As the blood ages, the density decreases
- Chronic blood approaches the density of CSF
- sizeable haemorrhage may cause mass effect and midline shift


الطب والجراحة لجنة

Extra-axial collection

- **Extra-axial collections** are collections of fluid within the skull, but outside the brain parenchyma. They may be comprised of CSF, blood or pus and may exist in the extradural, subdural or subarachnoid space.
- **Pathophysiology**: fluid collection in the extradural, subdural or subarachnoid space
  - Blood, e.g., post-traumatic extradural haemorrhage
  - Fluid, e.g., chronic subdural haemorrhage
  - Pus, e.g., subdural empyema



From the case: Extradural haematoma



From the case: Subdural empyema



From the case: Chronic subdural haematoma

الطب والجراحة لجنة

Extra-axial collection

Role of imaging

- Determine whether there is an extra-axial collection
- Determine where the collection is:
  - Subarachnoid
    - Extends down into the sulci
    - Extends into the basal cisterns
  - Subdural
    - More crescent-like shape
    - May cross sutures
  - Extradural
    - Lentiform (lens-like) shape
    - Do not cross cranial sutures
- Determine whether there is mass effect and:
  - Midline shift
  - Hydrocephalus

الطب والجراحة لجنة

Section5: Systematic Approach to the Interpretation of CT Head

Sources:

CT Interpretation: Head

Emergency CT head

Systematic Approach to the Interpretation of CT Head

1. Take a fast look through out the CT head to spot any obvious findings, then start from the top or the bottom of the CT according to your preferences (many people prefer starting from above because it has a smaller number of structures to look for)
2. Follow the mnemonic "**B**lood **C**an **B**e **V**ery **B**ad" which is used to read an **emergency head CT scan**.
   - Blood: Blood
   - Can: Cisterns
   - Be: Brain
   - Very: Ventricles
   - Bad: Bone
3. Don't forget to assess the important key findings

الطب والجراحة لجنة

Blood: Blood

- **Appearance on CT: (Density)**
  - Acute (hours to days) - New blood is White
  - Subacute (days to weeks) - Grey
  - Chronic (weeks to months) - Old blood is Dark
  - Acute on Chronic - Layering effect

- **Types/Locations**
  - Intraparenchymal Hemorrhage/Contusions
  - Intraventricular Hemorrhage
  - Extra axial Hemorrhage (e.g., subarachnoid, subdural, epidural)
  - Extracranial Hemorrhage (e.g., Scalp hematomas)

الطب والجراحة لجنة

Hemorrhagic Stroke

With any intracerebral hemorrhage the following points should be included in a report as they have prognostic implications:

- Location
- size/volume
- shape (irregular vs regular)
- density (homogeneous vs heterogeneous)
- presence/absence of substantial surrounding oedema that may indicate an underlying tumor
- presence/absence of intraventricular hemorrhage
- presence/absence of hydrocephalus
- when CT angiography is performed, the presence/absence of the CTA spot sign or a vascular malformation

الطب والجراحة
لجنة

Can: Cisterns

look for the presence of blood, effacement and asymmetry in four key cisterns

- **Circummesencephalic cistern**: CSF ring around midbrain and most sensitive marker for elevated ICP
- **Suprasellar cistern**: Star-shaped space above the sella
- **Quadrigeminal cistern**: W-shaped space at the top of the midbrain
- **Sylvian cistern**: Bilateral space between temporal/frontal lobes

الطب والجراحة لجنة

Be: Brain

Evaluate the brain parenchyma, including:

- Look for asymmetry or effacement of the gyri/sulci pattern
- structural shifts
- gray-white matter differentiation (including the insular ribbon sign)
- abnormal hypodensities (e.g., air, oedema, fat) or hyperdensities (e.g., blood, calcification)

الطب والجراحة لجنة

Very: Ventricles

- Evaluate the ventricles for
  - Dilation or compression
  - Look for intraventricular haemorrhage
  - Ventricular effacement or shift
  - Hydrocephalus
- Compare the ventricle size to the size of cisterns, large ventricles with normal/compressed cisterns and sulcal spaces suggests obstruction.

الطب والجراحة لجنة

Bad: Bone

- Soft tissue swelling, mastoid air cells and paranasal sinuses fluid in the setting of trauma should raise the possibility of a skull fracture; intracranial air means that the skull and the dura have been violated somewhere (we still on brain window)
- Switch to bone windows to evaluate for fracture.
- The identification of small, linear, non-depressed skull fractures may be difficult to identify as they are often confused with sutures – surrogates include pneumocephalus, and abnormal aeration of mastoid air cells and sinuses.
- The Presence of fractures increases the suspicion for intracranial injury, search adjacent and opposing parenchyma and extra-axial spaces.


الطب والجراحة لجنة

Section6: Clinical Cases

Case 1

- **Presentation**: Patient with Alzheimer's disease, fever and lowered level of consciousness.
- **Patient Data**: Age: 82, Gender: Male
- What is your spot diagnosis ? What do you think is the cause (pathology)?





الطب والجراحة لجنة

Case 1


- **Findings**: Hemorrhagic focus in the right frontal lobe, flanked by oedema of the adjacent parenchyma. Extension of bleeding for the subdural space along the right cerebral hemisphere, cerebellum tent and space interhemispheric.

- **Discussion**: **Large frontal lobar haemorrhage extending into the subdural space**. In an elderly patient this is most likely due to **cerebral amyloid angiopathy** rather than an underlying lesion, however no further exams were available.

- **From the case:** Lobar and subdural haemorrhage

Case 2

- **Presentation**: The patient was admitted to the emergency department following severe road traffic injury. His Glasgow Coma Scale (GCS) was 5 though he was intubated.
- **Patient Data**: Age: 25, Gender: Male
- What is your spot diagnosis ? What is this sequence?




Case 2

- **Findings**: Right fronto-temporal cortical and subcortical non-hemorrhagic contusion. Similar but smaller areas are noted on left occipital, left temporal and dorsal aspect of brainstem. Moreover, there is edema of the corpus callosum.
- **Discussion**: Grey-white matter junction, corpus callosum and dorsal aspect of the brainstem are common sites for **diffuse axonal injury**. The case illustrates findings consistent with grade III diffuse axonal injury. Unfortunately, the patient died few days after this scan.
- That is DWI sequence

- **From the case:** Diffuse axonal injury (DAI)

Case 3


- **Presentation**: Assault
- **Patient Data**: Age: 55 Gender: Male
- Mention as much findings as you can seen in these pictures of this case, note that the pictures might show 1 or more findings or even show nothing



Case 3

❖**Findings**:

- Non-contrast CT demonstrates a moderate left lateral convexity epidural haematoma, measuring 18 mm in maximal depth, associated with an undisplaced left temporoparietal fracture. Thin smear subdural haematomas are seen overlying the left and right temporal lobes, with further blood seen layering on the tentorium cerebelli on the right. 18 mm hemorrhagic contusion involving the left middle/superior temporal gyrus. Left parieto-occipital subgaleal haematoma.




Case 3

- **Discussion**: This case illustrates typical appearances of a significant closed head injury with a coup-contrecoup injury, the largest of which is a left-sided extradural haematoma (EDH).

- **From the case:** Extradural haematoma, subdural haematomas, cerebral contusions and subgaleal haematoma

Case 4


- **Presentation**: A teenager with multiple sensory and motor disturbances.
- **Patient Data**: Age: 16 Gender: Female
- What is your spot diagnosis? What do you expect to see on Axial T1 C+ and on sagittal FLAIR



Case 4

- Spot diagnosis: Multiple Sclerosis
- Axial T1 C+: Uptake of the contrast in active plaques
- Sagittal FLAIR: Dawson`s fingers
- **From the case:** Multiple sclerosis





Case 4

- **Presentation**: Driver in car accident. Reduced GCS.
- **Patient Data**: Age: 65 Gender: Male
- In this patient which of the seen hemorrhages are due to hypertensive aetiology and which are due to traumatic aetiology ?





Case 4

- **Findings**:
  - There are multiple intraparenchymal haemorrhages, the largest of which is centered within the left basal ganglia consistent with hypertensive aetiology.
  - The remaining smaller haemorrhages are located at the grey-white junction in the right frontal lobe and temporal lobe consistent with traumatic aetiology.
- **Discussion**:
  - This case is radiologically cute in that the intraparenchymal haemorrhages have two separate aetiologias. The patient most likely suffered a left basal ganglia hypertensive bleed whilst driving and then crashed his car receiving traumatic right frontal and temporal haemorrhages. (موتة حلوة ما شاء الله)
- **From the case:** Combined traumatic and hypertensive haemorrhage

الطب والجراحة لجنة

Case 5

- **Presentation:** This case is a bit different to the others in that it is actually comprised of 6 non-contrast CT scans from 6 different patients. One or more of them are normal. One or more of them have a hyperacute infarct. Can you work out which is which? And mention the sign you saw on the pathological photos








Case 5







الطب والجراحة
الجنة


Case 5


- Case A: Normal
- Case B: Hyperdense MCA sign
- Case C: MCA dot sign (Sylvian fissure sign)
- Case D: Hyperdense Basilar artery (Basilar tip thromboembolism)
- Case E: Hyperdense MCA sign & MCA dot sign (Sylvian fissure sign)
- Case F: Ribbon sign


Case credit: Frank Gaillard Case A rID:37008, Case B: Jeremy Jones rID: 13190; Case C: Ian Bickle rID: 44812; Case D rID: 19199, and Case E rID: 22117

Case 6

- **Presentation**: Cognitive impairment.
- **Patient data**: Age: 75 Gender: Female
- What is your spot diagnosis ? As this disease progress what changes do you expect to see







الطب والجراحة لجنة


Case 6

- **Findings**: Moderate chronic small vessel ischemic changes scattered throughout both hemispheres. The hippocampi are bilaterally, symmetrically, disproportionately volume depleted. The remainder of the brain demonstrates only minor generalized volume loss. Multiple peripherally located micro haemorrhages are seen throughout both hemispheres suggestive cerebral amyloid angiopathy. Features support a diagnosis of Alzheimer's disease
- We expect to see atrophic changes as the disease progresses

- **From the case:** Alzheimer's disease

Case 7

- **Presentation**: Fall from roof onto head.
- **Patient Data**: Age: 60 years Gender: Male
- What is your diagnosis ? Describe what can you see in these 2 photos



Case 7

- **Findings**:
  - Extensive calvarial, skull base and facial fractures. Associated with pneumocephalus
  - Traumatic subarachnoid and intraventricular haemorrhage.
  - Scattered petechial intraparenchymal haemorrhages including within the midbrain consistent with shearing injury.
  - Cerebral oedema with loss of grey-white differentiation, on the left in particular, possibly due to dissection of the left ICA at the skull base due to fractures.
  - There is a nice example of the 'white cerebellum sign' of cerebral oedema which is an extremely poor prognostic feature. This patient died shortly after.
- **Discussion**: A case of non-survivable head trauma with established diffuse cerebral oedema and petechial brainstem haemorrhages due to shearing injury (diffuse axonal injury).

الطب والجراحة لجنة

Case 8

- These 2 patients present with headache what is your spot diagnosis in each of the cases?



Case 8

- Case A: Subarachnoid hemorrhage
- Case B: Transverse sinus thrombosis

- **From the case:** Subarachnoid haemorrhage
- **From the case:** Transverse sinus thrombosis

Case 9 A

- **Presentation**: A previous history of intractable hypertension and presented with sudden onset of right sided weakness

- **Patient Data**:
  - Age: 77 years
  - Gender: Female


- What is your spot diagnosis ?



الطب والجراحة لجنة

Case 9 A

- Spontaneous intracerebral haemorrhage (Hypertensive haemorrhage)



Case 9 B

- **Presentation**: Left sided weakness after head injury


- **Patient Data**:
  - Age: 45 years
  - Gender: Male

- What is your spot diagnosis ?



Case 9 B


- **Findings**:
  - Hyperdense intra axial lesion in the Rt parietal (hemorrhage) with extra axial extension and compressing lateral ventricle on the rt side with dilated Lt lateral ventricle, there is fracture in the Rt parietal
- **Discussion**: Traumatic ICH
  - This intracerebral haemorrhage has the same characteristics as the image of spontaneous haemorrhage shown in the previous case
  - The bone window image (inset) shows an adjacent skull fracture in this patient following



Case 10

- **Presentation**: Quarrel
- **Patient Data**:
  - Age: 46 years
  - Gender: Male

- What is your spot diagnosis ?




Case 10

- **Findings**:
  - Multiple hemorrhagic foci in both fronto-parietal region
  - Mild shift of midline structure to the left.

- **Diagnosis**: Hemorrhagic contusion

Case 12

- **Presentation**: RTA
- **Patient Data**: Age: 68 years Gender: Male
- What is your spot diagnosis ?




Case 12

- **Findings**:
  - Hypodensities in both frontal region with mild shift of midline structures to the left

- **Diagnosis**: Non hemorrhagic contusion

Case 13 A

- **Presentation**: Motor vehicle collision, GCS 3
- **Patient Data**: Age: 70 years Gender: Female
- What is your spot diagnosis ?



Case 13 A

- **Findings**:
  - Extensive bilateral subarachnoid and subdural haemorrhage. Loss of grey-white differentiation, swelling indicate diffuse cerebral edema.

- **Diagnosis**: Diffuse cerebral edema

- **From the case**: Diffuse cerebral oedema

Case 13 B


- **Presentation**: Post CPR / severe trauma
- **Findings**: diffuse hypodensity of the bilateral cerebral hemispheres with loss of white-gray matter differentiation.
- **Diagnosis**: Diffuse cerebral edema



الطب والجراحة لجنة

Case 14

- **Presentation**: Headache
- **Patient Data**:
  - Age: 44 years
  - Gender: Male
- What is your spot diagnosis ?



- **Diagnosis**:
  - Aneurysm in the Rt MCA

الطب والجراحة لجنة

Case 14

Cerebral Aneurysm

- It is important to identify and treat these as a subarachnoid haemorrhage and cerebral haematoma from a ruptured aneurysm can cause sudden death. There are most commonly found involving the vessels of the Circle of Willis.
- **Appearance:** Localized dilatation along one of the cerebral vessels.
- Possible associated findings: Aneurysm clips (metallic artefact), Old bleeds, Burr holes (defect in skull)

الطب والجراحة لجنة

Case 15

- **Presentation**: History of meningitis
- **Patient Data**: Age: Child Gender: Female
- What is your spot diagnosis ?



Case 15

- **Findings**:
  - Dilated lateral ventricle.
  - Periventricular interstitial edema.
  - No focal lesion.
  - Midline structures no shift.

- **Diagnosis**: Acute hydrocephalus

Case 16

- **Presentation**: Previous vascular event for follow up

- What is your spot diagnosis ?



Case 16

- **Findings**:
  - Fluid like density (hypo dense) lesion in the Lt parietal
  - Dilated ventricular system mainly the Lt side
  - No mass effect or shift of midline structures.
  - Diffusion dark lesion
- **Discussion**: After vascular insult (infarction). The dead tissue replaced by fluid called encephalomalacia that shows negative mass effect and so dilated nearby ventricle.

الطب والجراحة لجنة

Case 17

- **Presentation**: Diabetic hypertensive with headache
- **Patient Data**:
  - Age: 54 years
  - Gender: Male
- What is your spot diagnosis ?



Case 17


- **Findings**:
  - Small hypodense lesion in the Lt thalamus
  - No mass effect
  - No shift of midline structure
  - No dilation of ventricular system
- **Discussion**: Lacunar infarcts
  - Lacunar infarcts are small (<15 mm)
  - They result from occlusion of one of the small penetrating end arteries at the base of the brain and are due to fibrinoid degeneration.
- **Clinical presentation**
  - Most lacunar infarcts are clinically silent, but repeated episodes are associated with vascular dementia.



الطب والجراحة لجنة


Case 18

- **Presentation**: Trauma
- What is your spot diagnosis ?



Case 18

- **Findings**:
  - Hyperdense lesion in the Lt frontal lobe (hemorrhage) surrounded with edema.
  - Slight effacement of surrounded sulci.
  - No shift of midline structures.
  - Focal subarachnoid hemorrhage in the Lt lobe
- **Diagnosis**: ICH with subarachnoid extension



Case 19

- Define the artery responsible of the infarct shown by each of the following photos







Case 19

- **Case A**: ACA
- **Case B**: MCA
- **Case C**: PCA

Case 20

- **Presentation**:
  - Fell out of bed approximately one hour earlier. No loss of consciousness. Alert, asymptomatic, no focal neurological deficit.
- **Patient Data**:
  - Age: 85 years
  - Gender: Male
- What is your spot diagnosis ?



الطب والجراحة لجنة

Case 20

- **Findings**:
  - Bilateral widened subdural space with low density collections (best appreciated near the vertex) indicative of chronic hygromas.
  - Cortical veins traversing medially to the collection, confirming its subdural location.
- **Discussion**: Typical appearance of incidentally encountered chronic subdural hygroma.



Chapter 3
Chest Imaging

Done By:Yousef Tarawneh

Sources: Slides + slides notes + CXR made easy for medical students + Radiopedia / and other sites

Writer's notes

- I made this chapter using the doctors slides + their notes + other useful books and sites (CXR made easy for med students book + Radiopedia / other sites)
- The notes are colored blue
- The info from books and sites are colored green
- And other colors are from the slides

- Try to get used to the normal view of the CXR and know how to interpret the CXR well before proceeding with the pathologies and their CXR mentioned in this chapter

- Good luck

الطب والجراحة لجنة

Table of Content

1. Intro and Normal lung x-rey and its contents
2. How to read and interpret a CXR
3. Consolidation + silloute sign + air bronchogram
4. Pneuomonia and TB
5. Lobar collapse and emphysema
6. Solitary and cavitating lesions of the lung
7. The Mediastinum
8. Sarcoidosis
9. Pleural effusion and pneumothorax
10. Lung tumors
11. Cardiomegaly and heart failure
12. Chest trauma


الطب والجراحة لجنة

CHEST IMAGING

- Chest imaging remains major component of diagnostic radiology.
- The chest x-ray is the most commonly performed diagnostic x-ray examination.
- A chest x-ray makes images of the lungs, heart, airways, blood vessels and the bones of the spine and chest.

Spinous processes (white rings) overlying trachea (black)

Hilar regions



Anatomy of the lungs

The Right lung is larger than the left, because most of the heart encroaches on the left lung.

- **The right lung has three lobes:**
  - upper lobe
  - Middle lobe
  - Lower lobe
- **The left lung has two lobes:**
  - upper lobe
  - lower lobe

Anatomy of the lungs /2


Adjacent lobes are separated by an interlobar fissure.


In the right lung:

- The minor (horizontal) fissure separates the upper lobe from the middle lobe.
- The major (oblique) fissure separates the upper and middle lobes from the lower lobe.

In the left lung:

- **The major or oblique fissure separates the upper lobe from the lower lobe.**


L


Upper Lobe
Major Fissure
Minor Fissure
Middle Lobe
Lower Lobe
Right Lung

Segmental anatomy

Segments of the right lung

Upper lobe segments:

- Apical segment
- Anterior segment
- Posterior segment

Middle lobe segments

- Lateral segment
- Medial segment

Right lower lobe segments:

- Superior segment
- Medial basal
- Anterior basal
- Lateral basal
- Posterior basal

Left lung segments:

Left upper lobe


- Apical posterior segment
- Anterior segment
- Superior lingular segment
- Inferior lingular segment

The lingular segments in the left lung is similar in position to the right middle lobe.


Left lower lobe segments:

- Superior segment
- Medial basal
- Anterior basal
- Lateral basal
- Posterior basal


Or lingular segment



Normal lung x-rey and its contents




A normal chest radiograph showing the left and right side.




A normal chest radiograph demonstrating lung zone anatomy.


1. Trachea (light blue)
2. Carina – spinal level T5 (black dotted line)
3. Aortic arch/knuckle (green)
4. Descending thoracic aorta (green dashed line)
5. Left heart border (yellow)
6. Left hemidiaphragm (pink)
7. Right hemidiaphragm (purple)
8. Right heart border (red)
9. Superior vena cava (blue)




1. Clavicle (green)
2. 5th rib – posterior aspect (red)
3. 5th rib – anterior aspect (yellow)
4. Right costophrenic angle (purple)
5. Left costophrenic angle (pink)
6. Right hilum (containing the right hilar lymph nodes) (light blue)
7. Left hilum (containing the left hilar lymph nodes) (blue)
8. Lung apex (pl. apices) (orange)





الطب والجراحة لجنة





A normal chest radiograph with the pulmonary vascular markings in red.






Left half of a chest radiograph demonstrating changes in lung markings from medial to lateral

How is a chest X-Ray taken

- Upright Posterior-Anterior view (PA)
- Lateral view if needed (usually from right to left to make heart closer to film).

The term PA refers to the direction of the x-ray beam which traverses the patient from posterior to anterior.


Distance between X-ray source and patient is 180 cm (~6 feet)
Center of X-ray beam
Lead plates for protection

PA chest radiography is much superior to AP, why ??

- Less magnification of the heart because it's closer to heart.
- More lung fields are visualized.
- The PA projects the scapula away from the lung fields.
- The apices of the lungs are closer to the film in the PA and appear more clear.

Postro-anterior




Antero-posterior



Other plain chest radiography:

- **Antero–posterior view (AP)**
  - ★ very ill patients who are unable to stand.
  - ★ infants and small children.
- **Inspiration–expiration films**
  - ★ suspected bronchial foreign body aspiration.
  - ★ suspected small pneumothorax.


120 cm

Before reading the x-ray film, the following should be checked:


1. Request form

- name, age, sex and date.
- clinical information (history and physical examination because it helps us in differential diagnosis)

2. Technical quality on the film:

- **markers:** L for left, R for right.
- **Centering (patient position).**
  Medial end of the clavicles equidistant from T4-T5 spinous process.
  another sign is that trachea will be overlying spinous processes unless the patient came with trauma.
- **Degree of inspiration.**
  at least the anterior ends of 6th rib above mid part of diaphragm
  or posterior end of 9th rib above the diaphragm
- **Exposure (penetration):** vertebral bodies and disc spaces behind the heart must be barely visible and the vascular markings should be visible through the heart.
  note : intervertebral discs are invisible in over and underexposure.
  note : overexposure is also known as high penetration, or dark film.
  low exposure is also known as soft film, or low penetrarion.

Normal lung x-rey with no rotation




CXR ROTATION


- Patient is rotated to the left: the medial ends of the clavicles will be deviated to the left (patient's left) of the spinous processes of the upper thoracic vertebrae.


- Patient is rotated to the right: the medial ends of the clavicles will be deviated to the right (patient's right) of the spinous processes of the upper thoracic vertebrae.

الطب والجراحة لجنة




Non-rotated chest radiograph. The clavicles are yellow and the medial ends of the clavicles are marked with a black dotted line. The position of the spinous processes of the upper thoracic vertebrae are shown with a white dotted line. The spinous processes lie midway between the medial ends of the clavicles, therefore this radiograph is not rotated.







Chest radiograph of a patient rotated to the right. The clavicles are marked in yellow and the medial ends of the clavicles are marked with a black dotted line. The position of the spinous processes of the upper thoracic vertebrae are shown with a white dotted line. The medial ends of the clavicles are deviated to the right of the spinous processes, therefore this


Inspiration vs Expiration








Chest radiograph with adequate inspiration. The right radiograph shows
the anterior parts of the 1st to 6th ribs marked in yellow and the left and right





Chest radiograph with inadequate inspiration. Notice how the lung volumes are smaller when compared to the previous figure (Figure 3.4). There is the false appearance of cardiomegaly. The anterior parts of the 1st to 4th ribs are marked in yellow and the left and right hemidiaphragm are marked with dashed white lines. (In this example, you can also see gas under the right hemidiaphragm in keeping with pneumoperitoneum.)

الطب والجراحة لجنة

Improper Exposure

Underexposed




Overexposed



Systematic interpretation of a chest x-ray:

- **Lung fields** : from top to bottom, both lungs should be of same radiolucency , and we must look at blood vessels of lungs which should be visible, and on the cardiophrenic and costophrenic angles
- **Trachea** : must be centralized but towards its end the trachea is slightly shifted to the right especially in elderly because of presence of aortic arch. also in hypertensive patients the aortic arch can be enlarged leading to shift of the trachea.
  note : left beonchus is longer but higher and more curved than right
- **Hilar shadows** : contains lymph nodes which are typicaly invisible due to their small size (5-6mm) but can become enlarged (more than a centimeter) and visible.
  note : left hilum is usually higher than right 1-3 cm.
- **Heart and Mediastinum** : heart borders must be well defined (sharp). Area of heart left to spine must be the same radiopacity as right. Note that lung vessels can been seen through the heart.
- **Diaphragms** : normally the diaphragm is well defined and convex upwards, left part is slightly lower than right due to presence of heart.
- **Below diaphragms** : gas in fundus of stomach appears in PA position while in supine gas appears in body and antrum.
- **Free air or abnormal calcification**
- **Soft tissues and Bones**
- **Outside the chest for subcutaneous emphysema**

Normal CXR



•What to look in the lungs?


- ❖ Size of the lungs.
- ❖ Density (Radiolucency) .

  Both lungs should be of equal radiolucency.


  Compare the two lungs at the same level, starting at the apices.


- ❖ Pulmonary vascular pattern.
- ❖ Look for abnormal shadowing or lucency:
  - Radio-opacity (whiteness) = increased density.
  - Radiolucency (blackness) = decreased density.
- ❖ If a lesion is identified:
  - localize the lesion
  - Describe the lesion.
  - Give diagnosis or differential diagnosis.

Normal CXR



Lateral CXR


R humerus
L humerus
Trachea
Aortic arch
Posterior surface of arms
Manubrium
L main bronchus
Sternum
R Scapula
L Atrium
L Ventricle
R Ventricle
Inferior vena cava
L Costophrenic angle
R Costophrenic angle
Gastric bubble


L wiki
Lung Apicies
Oesophagus
Trachea
Sternum
Scapula
Ribs
Hilar Region
Heart
Thoracic Vertebrae
Ribs (posterior)
Diaphragm
Costophrenic Angles


Consolidation

- Accumulation of inflammatory fluid in the alveoli of the lung, causing lung opacity.
- The fluid can be inflammatory exudate, transudate, pulmonary edema, pus or blood.


The most important radiological findings associated with Consolodation


- **Heterogenous or patchy opacification: the opacification is nonuniform and the border is not well demarcated (it sometimes looks fluffy).**
- **Lobar or segmental distribution: the opacification usually corresponds anatomically to a lobe or lung segment.**
- **No loss of lung volume: as a general rule, there is no significant loss of lung volume in consolidation.**
- **Air bronchogram : we will talk about that later .**

الطب والجراحة لجنة




Large area of consolidation in upper lobe and part of the middle lobe because heart border isn't clear





Consolidation in the lingula (green). There is heterogenous airspace opacification in the left lower zone and no loss of lung volume. The left hemidiaphragm can be clearly seen, however the left heart border is poorly defined
(silhouette sign); therefore, the pathology is in the lung adjacent to the left heart
border, i.e. the lingula of the left upper lobe.


الطب والجراحة لجنة




Consolidation (green) in both lungs with moderate sparing of the right upper zone (which appears darker than the rest). There is heterogenous airspace
opacification, no loss of lung volume, and air bronchograms seen in both lungs. You
can also see an endotracheal tube (white arrowheads), ECG leads (white

Silhouette Sign

- Is the loss of the normal border or contour between two structures of same radiological density and at the same anatomical level.





Consolidation (green) in the middle lobe and left lower lobe. There is heterogenous airspace opacification in the right and left lower zones and no loss of lung volume. We know the consolidation on the right side is in the middle lobe as there is loss of definition of the right heart border (silhouette sign) and the superior border is the horizontal fissure (marked with a white dashed line). We know the consolidation on the left side is in the left lower lobe as there is loss of definition of the left hemidiaphragm (silhouette sign), yet the left heart border is still clearly visible.


الطب والجراحة لجنة




Consolidation in middle lobe of right lung because heart border is obliterated

Air Bronchogram

- Refers to the phenomenon of air filled bronchi being visible by the opacification of the surrounding alveoli.
- It is almost always caused by a pathologic airspace (alveolar) process.
- An air bronchogram is the radiographic appearance of an air-filled bronchus
- that is surrounded by fluid-filled or solid alveoli.
- It can appear when there is consolidation (e.g. pneumonia) or pulmonary oedema in the surrounding alveoli.



This is the same radiograph as Figure 7.6 and is a good example of an air bronchogram caused by severe bilateral consolidation. The air bronchogram is marked in grey on the inferior radiograph. You can also see an endotracheal tube, ECG leads, nasogastric tube in situ, and other overlying tubing.


Air filled bronchi on a backdrop of fluid filled alveoli
Bronchi aren't filled with fluid yet


INFLAMMATORY DISEASE OF THE LUNG

- Pneumonia.
- Lung abscess.
- Pulmonary tuberculosis.
- Fungous disease.
- Hydatid disease.

PNEUMONIA

- Is a lung infection, which can be caused by a variety of micro-organisms, including bacterias, viruses, and fungi.
- Lobar pneumonia: inflammation confined to a lobe of the lung.
- Bronchopneumonia: refers to bilateral multifocal areas of consolidation.


Chest X-ray showing consolidation in lower lobe of the right lung because diaphragm is obliterated and heart border is well defined.
Note: this is a PA position because heart isn't magnified and we can see gases in the fundus of stomach






Confirmation of lower lobe consolidation from lateral view
Note : this is left lateral view because of marker on picture

consolidation in middle lobe of right lung because heart border is oblitrated while diaphragm is well defined




Consolidation in lingular segment of left lung demonstrated by obliteration of heart border




Consolidation in Left lower lobe except superior segment and part of the lingular segment



Consolidation in the entire upper lobe of right lung indicating lobar pneumonia



Bronchopneumonia is characterized by multiple small nodular or reticulonodular opacities which tend to be patchy and/or confluent. This represents areas of the lung where there are patches of inflammation separated by normal lung



Patchy airspace opacity throughout the right lung in keeping with bronchopneumonia. Left lung is clear.
No pleural effusion.


الطب والجراحة لجنة


Multi-focal areas of consolidation in both lungs indicating bronchopneumonia




Atypical pneumonia - Q fever:

Hazy airspace opacity is seen in the left mid and upper zone which may be inflammatory in nature. No areas of confluent consolidation are identified.






❖**Round pneumonias are round-ish and while they are well-circumscribed parenchymal opacities, they tend to have irregular margins. They most commonly occur in superior segments of lower lobes and in the majority of cases (98%), they are solitary .**



Rounded region of consolidation is seen in the right upper lobe, suggestive of round pneumonia.
Bilateral otherwise faint opacities of both lungs, suggestive of associated bronchopneumonic changes.

❖Multilobar pneumonia, as the name suggests, is a lobar pneumonia affecting multiple lobes.



Complete airspace consolidation with air bronchograms of the right upper lobe, with further consolidation within the medial right middle lobe


الطب والجراحة لجنة

Aspiration Pneumonia /2


- Is inflammation of the lung and bronchial tree that develops due to the entrance of foreign materials (food, liquids or vomit) from the mouth or stomach into the airways and lung tissue.
- The location of aspiration pneumonia is often gravity dependent and depends on patient position:
  - The lower lobes, especially right side.
  - Right middle lobe.
  - Right upper lobe (patients who aspirate in **prone** position)

.

pneumonia in middle aspirated lower lobe



Consolidation in right and left upper lobes diagnosed as aspiration pneumonia (probably in prone position) confirmed by history



Pneumonia /3

- With treatment most types of bacterial pneumonia can be resolved radiologically within 10 days to 2 weeks.
- Viral pneumonia may last longer radiologically.
- Atypical pneumonias may take about 4 weeks to resolve completely radiologically.
- Note : clinical recovery precedes radiological

❑ What examinations should be considered in patients with pneumonia that does not resolve as promptly as it should ?

✓ **CT scan : could be tumor especially in patients older than 70 years. Or it could be a foreign body especially in young patients.**

✓ **Bronchoscopy**

PULMONARY TUBERCULOSIS

Is an infection caused by mycobacterium tuberculosis.

Primary pulmonary tuberculosis:

- Is the initial infection and present as an area of consolidation (Ghon focus) in any part of the lung with enlarged mediastinal and hilar lymph nodes (primary complex (Ranke complex)).
- Lymphadenopathy is a common feature of primary infection.

  note : consolidation + lymphadenopathy especially in children we should consider TB after pneumonia
- For every thousand pneumonia case a TB case presents

Post-primary TB (Chronic pulmonary TB):

This follows the primary infection after a latent interval and is due to either reactivation or reinfection.

Chronic pulmonary TB /2

Radiological features of chronic pulmonary TB:

- The disease may affect both upper lobes, or one upper lobe.
- In some cases, one upper lobe and the opposite mid zone.
- In the upper lobes, the apical and posterior segments are usually affected but almost never in the anterior segment.
- The superior segments of the lower lobes are also frequent site of TB (appears on hilar region in PA view).

RADIOLOGY OF PULMONARY TB/3

- Progressive or active TB infection is indicated by development of cavities, which could be single or multiples.
- The presence of fibrosis and volume loss indicate healing.
- Healed lesions often calcify.
- Chronic cavities are sometimes colonized by aspergillus mycetoma.

  note: cavities can occur in primary or chronic TB.

The most important radiological findings associated with TB

■■ Primary TB:
peripheral lung mass/consolidation (Ghon focus); enlarged hilar lymph nodes.

■■ Secondary TB (also known as post-primary or reactive TB):
consolidation often with cavitation; typically in upper lobes or apical segments of the lower lobes;

■■ may be associated with pleural effusions and/or pleural

* Previous (old) TB (as healing progresses, features you may recognize are):
1-fibrosis and volume loss;
2-calcified foci;
3-pleural calcification;
4-tuberculoma : a tumour-like mass resulting from a localised tuberculosis infection.

* Miliary TB:
discrete 1–2 mm nodules distributed evenly throughout the lung fields due to haematogenous spread.


الطب والجراحة لجنة

Pathogenesis :-


- Infections : Direct person to person transmission of airborne organisms from active case to a susceptible host



- The alveolar macrophage engulf the bacteria




- Formation of phagosome by the macrophage



- Bacterial protein inhibit the fusion between lysosome & phagosome

الطب والجراحة لجنة

❖Formation of granuloma with caseating necrosis in the center ***(Ghon`s Focus)***



❖If the Ghon`s focus move to Hilar lymph node ***( Ghon`s complex)***




❖After fibrosis and calcification  ***(Ranke complex)***

❖ ***CHEST X-RAY FINDINGS IN PRIMARY TB :-***
❖ ***1) cavitation***
❖ ***2) Hilar lymphadenopathy***
❖ ***3) pleural effusion***
❖ ***4) Ghon`s complex***
❖ ***5) consolidation***

Ghon`s complex


Usually subpleurial




Ranke complex




Hilar lymphadenopathy



Ranke complex




Hilar lymphadenopathy




Ghon complex






Primary TB with right apical consolidation (green). There is patchy opacification at the apex of the right lung





Primary TB with patchy right upper lobe consolidation (green) and cavity formation (yellow). The wall of the cavity is shown in light yellow.



Primary TB with bilateral upper lobe consolidation (green) and cavity formation (yellow). There is patchy opacification in both upper lobes. We know this is primarily consolidation rather than fibrosis as there is no significant volume loss. If you look closely, you can see that the horizontal fissure (white dashed line) is not deviated upwards. The wall of the cavity is shown in light yellow.

الطب والجراحة لجنة





Secondary TB (reactivation) with bilateral upper lobe volume loss, left upper zone consolidation (green), left hilar lymph node enlargement (orange), and old TB changes in the right upper lobe (green). There is patchy opacification in both upper lobes. There is elevation of both hila with the left and right main bronchi pulled superiorly causing widening of the carinal angle. The trachea and both main bronchi are outlined with a white dotted line and shown in blue.


الطب والجراحة لجنة






Consolidation of right upper lobe diagnosed and confirmed as TB from clinical history (his brother had TB)
Note : if no clues of TB are found in history we diagnose as neumonia because it's the most common (1k pneumonia : 1 TB patient) if patient doesn't improve with treatement we suspect TB

Wide areas of consolidation in upper lobes of both lungs with multiple Sattelite lesions characteristic of TB



Cavity inside consolidation from necrosis

Satellite lesions






This is a case of healing TB
What are the findings :
1- right lung collapse (atelectasis) compared to left lung


2- pleural thickening seen in apices but prominently on the right lung apex

3- calcifications (not confirmed)

4- linear fibrosis

5- tenting of diaphragm due to pleural fibrosis and adhesion




Pleural thickening on right and left apices and collapse of right upper lobe
Minor fissure
Pleural adhesion

MILIARY TUBERCULOSIS

- This is due to hematogenous spread of infection.
- Appear as small, discrete nodules,1-2 mm in diameter, evenly distributed throughout both lungs and no area of lung is spared.


  note : no other disease has such characteristics so it's straight forward diagnosis.

  note : miliary metastasis is variable in size and usually on the periphery of lung or on basal level segments. Miliary sarcoidosis affects the middle and slightly upper lobe



Three different chest radiographs of patients with miliary TB throughout both lungs. There are discrete 1–2 mm nodules distributed evenly throughout the lung fields.

Miliary TB



DISEASES OF THE AIRWAYS

- Chronic bronchitis.
- Pulmonary collapse
- Emphysema.
- Bronchiectasis.
- Cystic fibrosis.

Pulmonary collapse

- Pulmonary collapse or atelectasis refers to a decrease in volume of a lung, lobe or segment.
- Three mechanism that cause collapse:
  - **Obstruction.**
  - **Compression:** pneumothorax , pleural effusion.
  - **Contraction (scarring):** chronic TB, pulmonary fibrosis, also bronchiectasis.
- Obstruction to flow of air is the most common cause of collapse 80% of cases.

Pulmonary collapse / 2

- Common causes of bronchial obstruction causing collapse:

  - Bronchial carcinoma
  - Mucus plug (pneumonia, postoperative).
  - Foreign body.
  - Malpositioned endotracheal tube : usually takes an entire lung. The tip of the endotracheal tube should be at least 4 cm above the carina
  - Extrinsic compression of airway by tumor or enlarged lymph nodes.
  - Note : most common cause of lung collapse :

  1. For elderly : bronchial carcinoma
  2. For young : foreign body
  3. For pneumonia patients, or post-operative : mucal plug which usually takes a segment or a subsegment


0


Pulmonary collapse / 3

- The signs of lung collapse are:
  - The presence of radio-opacity.
  - Displacement of the interlobar fissure.
    Is the most reliable sign of direct lung collapse.
  - Displacemet of the hilar shadow.
  - Displacement of the trachea.
  - Shift of mediastinal structures.
  - **Elevation of the diaphragm (only in lower lobe collapse)**


right upper lobe collapse

- Diagrammatic representation of right upper lobe collapse. Increased density in the right upper zone (1), normal position of horizontal fissure (2), horizontal fissure displaced upwards (3), tracheal deviation to the right (4).






Right upper lobe collapse. There is increased opacification in the right upper zone (blue) with loss of definition of the upper right mediastinal margin. The
position of the horizontal fissure in a normal lung is shown with a white dotted line (1);
however, in this case the horizontal fissure has been pulled upwards (2).
The volume



Pic for comparison : Pneumonia of upper lobe, no displacement of horizontal fissure



Chest X-ray showing upward displacement of horizontal fissure of right lung due to collapse of right upper lobe

Features of middle lobe collapse on a chest radiograph:


■■ Increased density in right mid-zone(1).
■■ Loss of definition of the right heart border (the collapsed lobe lies against the right heart border making it indistinct)(1).

normal position of horizontal fissure (2)

■■ Horizontal fissure displaced downwards (pulled down by Collapsed lobe)(3).


Tracheal deviation to the right (4).

■■ The rest of the right lung looks blacker than the left lung (the right upper lobe and right lower lobe over-inflate to compensate for the reduced volume in the hemithorax caused by the collapsed lobe).



الطب والجراحة لجنة

Middle lobe collapse




Middle lobe collapse. There is increased density in the right mid-zone (blue) with loss of definition of the right heart border. The right hemidiaphragm is still clearly seen. The position of the horizontal fissure in a normal lung is shown with a white dotted line (1); however, in this case the horizontal fissure has been pulled downwards (2). The volume loss from the collapsed middle lobe has slightly pulled the trachea to the right side (3).

الطب والجراحة لجنة


Rt lower lobe collapse

- Triangular opacification at the right base medially(1).
- ■■ Loss of definition of the right hemidiaphragm(1).
- ■■ Elevation of right hemidiaphragm (pulled up by collapsed lobe)(1).
- ■■ Depression of right hilum and tracheal deviation (pulled down by collapsed lobe)(3).
- ■■ The right heart border is not obscured(2).
- ■■ The remainder of the right lung looks blacker than the left lung (the right upper lobe and middle lobe over-inflate to compensate for the reduced volume in the hemithorax caused by the collapsed right lower lobe).



الطب والجراحة لجنة


Rt lower lobe collapse




- Right lower lobe collapse. There is a triangular opacity at the right base medially (blue) with loss of definition of the right hemidiaphragm. The right heart border (1) is still clearly seen. The volume loss from the collapsed right lower lobe has slightly pulled the trachea to the right side (2).

Left upper lobe collapse

- Increased density in the left upper and mid zones with ‘Veil-like’ opacification in the left hemithorax with no clear lower border(1).
- ■■ Loss of definition of left upper cardiac border and left mediastinal margin(2).
- ■■ The descending thoracic aorta remains visible (outlined by air in the adjacent left lower lobe)(3).
- ■■ Elevation of left hilum (pulled up by collapsed lobe) and left hemidiaphragm.
- ■■ Trachea may be deviated to the left (collapsed lobe pulls trachea).



Left upper lobe collapse




Left upper lobe collapse. There is increased density in the left upper and mid zones with no clear inferior border and a loss of definition of the left heart border.
You can, however, see the outline of the left hemidiaphragm and descending thoracic aorta (black dotted line) as there is still air in the left lower lobe, which lies adjacent to these structures.


الطب والجراحة لجنة

Left lower lobe collapse

- Triangular opacification overlying the left heart medially (giving so called ‘double heart border’ sign)(1).
- ■■ Loss of definition of the medial aspect of the left hemidiaphragm(1).
- ■■ Elevation of the left hemidiaphragm (pulled up by collapsed lobe)(1).
- ■■ Outline of collapsed left lower lobe (2).
- ■■ Depression of left hilum (pulled down by collapsed lobe).
- ■■ The left heart border is still visible (3) .
- ■■ The remainder of the left lung looks blacker than the right lung (the left upper lobe over-inflates to compensate for the reduced volume in the hemithorax caused by the collapsed left lower lobe).




Left lower lobe collapse




- Left lower lobe collapse. There is a triangular opacity projected over the left side of the heart (blue). The lateral margin of the collapsed left lower lobe (1) together with the left heart border (2) can both be seen separately giving the ‘double heart border’ sign. There is elevation of the left hemidiaphragm (normally it should lie lower than the right) and reduced density of the rest of the left lung when compared to the right lung due to left upper lobe hyper-inflation (to compensate for the reduced volume in the left hemithorax caused by the collapsed left lower lobe).

Chest Xray showing different opacity of heart to the right and left of spine caused by collapse of posterior basal segment of lower lobe causing displacement of oblique fissure downwards, with elevation of left hemi-diaphragm. Note also left hilum is depressed.



Complete lung collapse



Complete left lung collapse. There is increased density throughout the entire left hemithorax and signs of decreased lung volume, such as displacement of the heart/mediastinum towards the collapsed lung and left tracheal deviation. The heart has moved from its normal position (1) to lie in the left hemithorax (2). The trachea is shown in blue and outlined with a white dotted line.


الطب والجراحة لجنة

Chest X-ray showing collapse of the right lung with shift of mediastinum, trachea, and heart to the right side indicating the mechanism of collapse is contraction. This patient has severe **bronchiectasis**

Note : this contraction collapse could appear in healed TB , and pulmonary fibrosis too.



Chest X-ray showing severely collapsed left lung by compression of pneumothorax in left pleural space demonstrated by invisibility of left lung vessels



Emphysema

- Is an increase in the size of the air spaces distal to the terminal bronchioles, with dilatation or destruction of their walls.
- Emphysema is included in a group of diseases called chronic obstructive pulmonary diseases (COPD).
- Complication: formation of emphysematous bullae and pneumothorax.

Radiological features of Emphysema

Sometimes deep inspiration can be confused with emphysema, so we need to look for these features in the X-ray to confirm emphysema


-PA view:

1. Diffuse hyperinflation of the lungs.
2. The lung appear more translucent with relative reduction in size and number of the small vascular markings.
3. The diaphragms are low and flat or oblique and flat while in very deep inspiration the diaphragm is convex
4. The heart shadow is long and narrow.

-lateral view :

- The postero-anterior diameter of the chest is increased in the lateral view resulting in **barrel chest.**
- Aorta appears more prominent because of translucency of lungs.
- You can also see flattening of diaphragm in this view.

Emphysema. Hyperexpansion with flattened diaphragms





Chest X-ray PA view showing hyperinflated lungs, low and flat diaphragm, narrowing of heart and mediastinum pulmonary vessels appear relatively small.



Chest X-ray lateral view showing increase in anteroposterior diameter (barrel chest) . More air fills the lungs so the background is more translucent making the aortic arch more prominent. We can also see flattening of diaphragm.

الطب والجراحة لجنة

SOLITARY PULMONARY NODULE
(Spot in the lung or Coin lesion)

- Is a small round growth usually less than 3 centimeters in diameter. If it's larger than that we call it a mass.
- Most nodules are benign but some can be malignant.
- A nodule is assessed for its:

- size

The larger the nodule, the greater the likelihood of malignancy.

- Margins (most important indicator)

Irregular contour or spiculated margins increase the probability of malignancy

Note : enlarged lymph nodes, metastasis (not primary tumor), and benign tumors have well defined margins

- Calcification


The presence of calcification within a nodule are in favor of benign lesion.

CAUSES OF SOLITARY PULMONARY NODULE

- Bronchial carcinoma.
- Metastasis.
- Hamartoma.
- Bronchial adenoma.
- Granuloma.
- Abscess.
- Hydatid cyst.

Very rare cases include :

- Bronchogenic cyst
- Arterio-venous malformation.
- Rheumatoid nodule.




Large mass lesion (red) in the middle lobe. You know it is located within the middle lobe because if you look carefully you can see that the mass is pushing up the horizontal fissure (white dashed line), which separates the right upper lobe and middle lobe.

الطب والجراحة لجنة




Small solitary mass lesion (red) in the left upper zone.

الطب والجراحة
لجنة



Chest X-ray : in right middle lobe there is a nodule of well defined margin and calcifications that are barely visible. This indicates a benign nodule



Left picture shows a chest X-ray of a patient showing a small nodule in right lung
Right picture is a chest X-ray of the same patient after 6 months shows how the nodule increased markedly in size indicating malignant tumor.
Note : benign nodules take at least 1.5 - 2 years to double in size

Cavitating lesion in the lung

- A cavity is a gas-filled space surrounded by a complete wall.
- Cavitation occurs when an area of necrosis communicates with a patent airway.

CAUSES OF CAVITATING PULMONARY LESION (in order of prevelance)

- Infection:
  - -Pneumonia (note : 1% of pneumonia present as rounded mass, simply called rounded pneumonia)
  - -TB
  - - Hydatid cyst
- Abscess.
- Bronchial carcinoma (most common cavitating tumor is spuamous cell carcinoma)
- Pulmonary infarct.
- Rheumatoid nodule.

Possible radiological signs of caviting lesions

- Centre of the lesion is darker than the periphery as no blood passes through it.
- Fluid level present: look for a horizontal line within the lesion.


There will be whiteness (fluid) below the line and blackness (air) above the line.

- Compare with old radiographs (it may be possible to see the cavity developing over time).




Cavitating lesion (yellow) in the right upper zone. It is likely to be a malignant mass as there are no large areas of surrounding consolidation or fibrosis.
The wall of the cavity is shown in light yellow.

الطب والجراحة
لجنة





Cavitating lesion (yellow) in the right upper lobe. The edge of the cavity is shown with a white dashed line. Within this cavity there is gas (dark) and a mass lesion (light yellow). This is an example of an aspergilloma or fungal ball, which appears as an intracavitary mass surrounded by a crescent of air.

الطب والجراحة
لجنة



Chest X-ray : in upper lobe of right lung, specifically anterior segment there is an area of consolidation with cavity inside (pneumatocele). We can know it's in the anterior segment because of prominence of horizontal fissure.





Chest X-ray : in right lower lobe there is an area of consolidation with cavity inside. This is rounded pneumonia (has 1% chance) . Use of CT scan to know if it contains pus or just air shows that it contained pus (grey)

Mediastinum

Anterior – Middle – Posterior

الطب والجراحة لجنة

The Mediastinum

- Situated between the lungs and extends from the thoracic inlet superiorly to the diaphragm inferiorly.

- The mediastinum is divided into:

✓ **Superior mediastinum:** Lies above the heart.

✓ **Anterior mediastinum:** Lies in front of the heart.


✓ **Middle mediastinum:** Contains the heart and great vessels.


✓ **Posterior mediastinum:** Lies behind the heart.

المهم بالموضوع هذا تعرف وين الكتلة موجودة بين هاي المناطق (Anterior, middle or posterior)
وتعرف ال (differential diagnosis) المتعلق بكل منطقه
بس تحديد ايش الكتلة بالزبط مش مهم الك في المادة لانه بيتطلب فحوص ثانيه وربط بالسيرة المرضيه وغيره


الطب والجراحة لجنة

normal chest radiograph with the mediastinum highlighted in orange







Mediastinum lateral view




The Mediastinum





ROLE OF IMAGING

- A mediastinal mass is identified by chest radiography as it typically manifests as an extra soft tissue mass or opacity.
- The use of the **silhouette sign ,** which describes the loss of normal borders of intrathoracic structures, increases the sensitivity of detecting mediastinal abnormalities.
- The density of soft tissue masses is similar to the mediastinal structures However, since the mass displaces the air-containing lung from the normal mediastinal structure, **the border of the normal mediastinal structure is lost**.
- Once an abnormality is identified by chest radiography, cross-sectional imaging is used to characterize the lesion, generate a differential diagnosis, assess for other abnormalities, and guide further management.(CT with intravenous (IV) contrast)


Great vessels /
Right paratracheal stripe
Superior vena cava
Arch of azygous vein
Superior vena cava
Right atrium
Inferior vena cava
Great vessels /
Left paratracheal stripe
Arch of aorta
Main pulmonary artery
Auricle of left atrium
Left ventricle
Fat pad



MEDIASTINAL MASSES

(514-559)

Anterior – Middle – Posterior


Note : Mediastinal masses appear as widening of mediastinum, when describing these masses we should describe their margins

الطب والجراحة لجنة


ANTERIOR MEDIASTINAL MASSES


- Lymphoma (Terrible lymphoma).
- Thyroid (Retrosternal goiter).
- Teratoma.
- Thymic tumor.
- Pericardial cyst.
- Diaphragmatic hernia (morgagni hernia).

الطب والجراحة لجنة



Nodal masses
(Terrible) Lymphoma
Aneurysm (fusiform or saccular)
Acute aortic syndrome
Thoracic aorta (ascending)
Anterior Mediastinal Lesions
Thymic lesions
Thymoma
Other thymic neoplasms e.g. carcinoma, carcinoid
Benign lesions e.g. cysts
Thyroid
Multinodular goitre
Teratoma
Other germ cell tumours including primary mediastinal seminoma
Germ cell tumours


an abnormal lobulated contour which is extending out through the left mediastinal contour. On our lateral radiograph, you'll notice that there is loss of the retrosternal clear space so this anatomically places the lesion within the anterior mediastinum.






CT which demonstrated a low-density homogenous structure in the anterior mediastinum, which was a known to be a thymic cyst, This pleural effusion was incidental finding






an abnormal bulging in the left upper mediastinal contour which is seen separate to the aortic knuckle.
Again on the lateral projection, our retrosternal clear space is lost, which places this abnormality in the anterior mediastinum.






cross-sectional imaging which demonstrated a large mass in the anterior mediastinum.
This was homogeneously enhancing with a bit of a lobulated contour.
This patient was diagnosed with lymphoma.






there is a lobulated lesion On the lateral , although the retrosternal clear space is preserved superiorly, the more inferiorly you move you can see that it is effaced.
That allows us to place it in the anterior mediastinum and also we can see here that there is clearance posteriorly.







اضغط على Esc للخروج من وضع ملء الشاشة

CT which demonstrated a large heterogeneously enhancing mass in the anterior mediastinum. Incidentally, this was biopsy-proven thymic neuroendocrine carcinoma.





Heterogeneous with fat, fluid, soft tissue, & calcification → *Benign teratoma



MIDDLE MEDIASTINAL MASSES

- Aneurysm of aortic arch.
- Bronchogenic cyst.
- Lymph node enlargement

الطب والجراحة لجنة

1.Aortic Arch Aneurysm X-ray






Aortic Arch Aneurysm CT scan







2.Bronchogenic Cyst


- abnormal budding of the bronchial tree during embryogenesis
- They can occur in the mediastinum or be intrapulmonary
- The most common location is the middle mediastinum



الطب والجراحة لجنة


EXPIRATION
Retro
AP


Paratracheal Area



Carinal Area

3.Lymph node enlargement


- ❖Chest lymph nodes: Paratracheal and Hilar lymph nodes
- ❖Causes of lymph node enlargement
- ✓Lymphoma
- ✓primary tuberculosis
- ✓sarcoidosis

الطب والجراحة لجنة

Paratracheal stripe - Normal




Right and left paratracheal stripes



Normal chest X-ray

Right Paratracheal Lymphadenopathy




The right paratracheal stripe is not seen, having been obliterated by a right paratracheal mass (arrowheads).




CT scan demonstrates right paratracheal lymphadenopathy (arrow).

الطب والجراحة لجنة

Left Paratracheal Lymphadenopathy




An abnormal convex border (arrow).




CT scan demonstrates lymphadenopathy (arrow).

الطب والجراحة
لجنة



Widening of the right paratracheal space in keeping with soft tissue in the middle mediastinum



There is widening of the right paratracheal stripe which is concerning for lymphadenopathy. The right hilum is also a little bulky which may also be due to lymphadenopathy.


الطب والجراحة
لجنة

Hilum overlay sign


- The sign refers to preserved visualization of the hilar vessels, excluding abnormalities that localize to the middle mediastinum.
- Useful in differentiating whether an opacity on a frontal chest radiograph in the region of the lung hilum is located within the hilum versus anterior or posterior to it.
- If the cause of the opacity arises from the hilum, the silhouette of the normal pulmonary vessels (interlobar artery, upper lobe arteries, and left lower lobar artery) is obliterated. Causes of these opacities include middle mediastinal tumors, hilar adenopathy, pericardial effusion, vascular enlargement, and cardiac enlargement.
- If the edges of the vessels are appreciated, this implies the cause of the opacity is not in contact with the hilum and is, therefore, either anterior or posterior to it. Most of these opacities are masses in the anterior mediastinum.


الطب والجراحة لجنة

Positive Hilum overlay sign – Ant. Mass




Positive Hilum overlay sign – Post. Mass






Negative **Hilum overlay sign**




Tuberculous hilar lymphadenopathy



20 years old female patient presented with non-productive cough. High community prevalence of TB.

Sarcoidosis hilar lymphadenopathy



Multinodular hilar hyperdensities indicating hilar lymphadenopathy are visible bilaterally (green overlay).
The symmetrical appearance of bilateral hilar lymphadenopathy is a characteristic feature of


الطب والجراحة لجنة

POSTERIOR MEDIASTINAL MASSES


- Aneurysm of descending aorta.
- Hiatus hernia.
- Dilated esophagus (especially achalasia).
- Paravertebral mass or abscess.
- Neurogenic tumors
  - ✓Neurofibroma
  - ✓Ganglioneuroma

الطب والجراحة لجنة

1.Descending aortic aneurysm.




This image shows a grossly enlarged descending aorta due to a thoracic aortic aneurysm



2. Hiatus Hernia

Chest X-ray showing a mass behind the heart with air fluid level, Absence of gas below diaphragm indicates hiatal hernia, confirmed by barium meal





الطب والجراحة لجنة

Hiatus Hernia




A plain chest radiograph showing a well-defined, rounded lucency in the retrocardiac region consistent with a sliding hiatal hernia.

الطب والجراحة لجنة

"Hourglass" Stomach Appearance

Sliding hiatal hernias are characterized by an "hourglass" stomach appearance on imaging






3.Dilated esophagus

Most common cause achalasia

Classic appearance of achalasia on radiographs:


- **convex opacity overlapping the right mediastinum**
- **absent gastric bubble**
- **anterior displacement and bowing of the tracheal shadow on the lateral view**
- **bilateral patchy alveolar opacities**



Achalasia

The esophagus is grossly dilated and measures up to 10 cm in diameter. There are two air-fluid levels projected at the level of the posterior third rib. The gastric bubble is however present. These findings are in keeping with achalasia.

The aortic arch and descending thoracic aorta are unremarkable.
The lungs are clear with no evidence of aspiration.




Bird’s Beak Sign


Achalasia present on barium swallow with the bird’s beak sign




Paravertebral mass



Right and left paraspinal stripes



Paravertebral mass




Neurogenic tumors

- ✓ **Neurofibroma; a benign nerve sheath tumor that can arise anywhere on the body**
- ✓ **Ganglioneuroma; rare, benign tumor, most commonly located in the abdomen**

الطب والجراحة لجنة

Ganglioneuroma




A large paravertebral mass with widening of lower neural exit foramina.

الطب والجراحة لجنة


Ganglioneuroma

CT confirms the presence of a very large posterior mediastinal/paravertebral/retroperitoneal mass extending into the neural exit foramen. The liver is displaced anteriorly. the right kidney displaced inferiorly.










Chest X-ray showing middle and anterior mediastinal masses of well defined margin, so most likely diagnosis is lymphoma




Chest X-ray showing widening of mediastinum more to the left side.
Lobulated well defined masses most likely lymph node enlargment



Chest X-ray showing right side mediastinal mass of well defined margin. Diagnosis was lymphoma.




Chest X-ray showing a well defined mass arising from posterior mediastinum to the right. Found to be ganglioneuroma





Chest X-ray showing a mass behind the heart with air fluid level, Absence of gas below diaphragm indicates hiatal hernia, confirmed by barium meal in next pic.

Barium meal indicating a hiatal hernia




Chest X-ray showing mass behind heart (borders of which marked yellow) with no gas below diaphragm but way up the thorax almost at the neck, confirmed to be achalasia by barium swallow in next pic






Bird beak sign



Aneurysm of arch and descending aorta


Normal aortic diameter less than 3.5 cm
3.5 – 4.5 cm dilated
More than 4.5 cm aneurysm
Normal abdominal aorta less than 2.5 cm



Sarcoidosis

- Is multisystemic disease of unknown etiology.
- It is characterized by the development of non-caseating granulomas,which either resolve or become fibrotic.
- Sarcoidosis commonly causes thoracic lymphadenopathy, predominantly affecting the bronchopulmonary nodes (Hilar nodes).
- The hilar Lymph nodes enlargement is almost always bilateral and symmetrical.

Note : enlarged lymph nodes, metastasis (not primary tumor), and benign tumors have well defined margins



Pleural effusion

- Is fluid collection in the space between the parietal and visceral layers of the pleura, usually contains serous fluid, but may have differing contents.

- **Haemothorax**: blood, usually following trauma.
- **Empyema**: purulent fluid (pus).
- **Hydropneumothorax**: fluid and air.


Pleural effusion / 2

- **The radiological features of pleural effusion on a chest x- ray are:**
  - Homogeneous opacification.
  - Loss of the diaphragm outline.
  - No visible pulmonary or bronchial markings.
  - Concave upper border which appear higher laterally.
  - blunting or obliteration of the costophrenic angle.


Pleural effusion

Diagrammatic representation of a left-sided pleural effusion. There is a homogenous dense opacity (1) with a concave upper margin giving a typical ‘meniscus shape’ (2) and loss of visualisation of the left costophrenic angle (3).






Small left-sided pleural effusion (green). There is loss of visualisation of
the left costophrenic angle. The upper edge is concave in shape giving a ‘meniscus’ appearance laterally.




Bilateral pleural effusions (green). The effusions are homogenous, dense opacities and the left-sided effusion is larger than the right. There is a loss of visualisation of the left and right costophrenic angles and both hemidiaphragms. The upper edges are concave in shape giving a ‘meniscus’ appearance laterally. The rightsided effusion appears to have two upper edges; one is fluid tracking anterior to the lung and the other is fluid tracking posteriorly (white and black dashed lines).

الطب والجراحة لجنة





Massive left-sided pleural effusion (green). There is a homogenous, dense opacity in the left hemithorax with loss of visualisation of the left costophrenic angle and left hemidiaphragm. There is no meniscus as the pleural effusion extends all the way to the apex of the left lung. You can tell that this is a pleural effusion and not lung
collapse or a pneumonectomy because there is mass effect, i.e. the trachea (outlined with a white dotted line and shown in blue) and mediastinum are pushed away from the increased density

Chest X-ray showing large area of radiopacity on the lower right side of the chest with concave upper margin (meniscus sign) higher laterally than medially, diaphragm and costophrenic angle are obliterated, collapse of right lung, and shift of the mediastinum to the left side indicating pleural effusion



Chest X-ray showing an area of radiopacity in the left lower part of the thorax of a concave upper margin (meniscus sign) higher laterally than medially, with obliteration of the diaphragm and the costophrenic angle, and no shift of mediastinal structures indicating mild pleural effusion






Same as previous cases

Slight shift in mediastinum

Generalized opacity of the left part of the thorax still with obliteration of diaphragm because the patient is in supine position
To confirm this we use ultrasound or CT (CT is very sensitive to pleural effusion




Pneumothorax


- Is the presence of free air in the pleural space, by a tear in either the parietal or visceral pleura.
- **The most common cause of pneumothorax is chest injury, but the most common cause of spontaneous pneumothorax is rupture of sub-pleural emphysematous bullae (bleb).**
- Types of pneumothorax :
  - Closed or simple pneumothorax : a tear of pleura allows a limited amount of air to enter then it's sealed allowing no further entry of air
  - Open pneumothorax : a tear of pleura allows air to enter during inspiration and exit during expiration
  - Tension pneumothorax : a tear of pleura allows air to enter during inspiration but can't exit during expiration, so air is trapped in the pleural space with continuous addition, this is why it's progressive and fatal.

Radiological features of pneumothorax

- **Lung edge:** a thin white line at the lung margin, represent the visceral pleura.
- **Absent lung markings** between the lung edge and chest wall.
- **Mediastinal shift:** occur when a tension pneumothorax develops.
  - More prominent vascular markings in the opposite lung.
  - As a pneumothorax causes the affected lung to collapse, most of the right ventricular output is delivered to the opposite lung, leading to increased vascular markings on an erect chest radiograph.


Diagrammatic representation of a left-sided pneumothorax. The left lung is collapsed (1), there is air in the surrounding pleural space (2) which appears blacker, and you can see the edge of the lung (3).





Right-sided pneumothorax (blue). The lung edge (white dashed line) is seen and there are no lung markings beyond this lung edge. The vascular markings are also slightly more prominent in the opposite lung

الطب والجراحة لجنة




Large right-sided pneumothorax (blue). The lung edge (white dashed line) is clearly seen and there are no lung markings beyond this lung edge. Remember that the lung always collapses inwards (medially), while the pneumothorax is seen around the edge of the lung (peripherally) and will always be less dense (blacker) than the lung tissue. Sometimes students mistake the outline of the scapula (black dotted line) for the lung edge, so be careful not to make this basic mistake

الطب والجراحة لجنة


Chest X-ray showing severely collapsed left lung and a pneumothorax indicated by absence of blood vessels . There is no mediastinal shift so there is no tension



Chest X-ray showing collapse of right lung and pneumothorax indicated by lack of blood vessels. There is slight shift of mediastinal structure indicating tension pneumothorax



Edge of lung indicated by arrows
No tension




Tension pneumothorax radiological findings

Darkening of the hemithorax and loss of lung markings due to air in the pleural space.

- Increased volume of the hemithorax.
- Displacement of the mediastinum (and trachea) away from the pneumothorax.
- Depressed hemidiaphragm.

الطب والجراحة لجنة



Left-sided tension pneumothorax (blue). There is mediastinal shift to the
right (normal position of the left heart border (1), new position of left heart border (2)), tracheal deviation to the right (outlined with a white dotted line and shown in blue), depression of the left hemidiaphragm (3) and loss of normal lung markings on the left side as the tension pneumothorax occupies the whole hemithorax

الطب والجراحة لجنة

TUMORS OF THE LUNG

- Lung cancer is the commonest fatal malignancy.
- The strongest risk factor is cigarette smoking. 70%
- More than 95% of malignant tumors arise from the respiratory epithelium and are termed bronchogenic carcinoma.
- Less than 5% of lung cancers are of rare cell types, such as carcinoid tumors, lymphoma, or metastasis.
- **Usually, we can't diagnose tumors of the lung with just a simple CXR so more imaging(CT/PET/NUCLEAR IMAGING) and further evaluation is required**

Types of lung cancers

1- Adenocarcinoma

- Is the most common type of lung cancer, making up 30-40% of all cases.
- Usually arise **peripherally** as solitary pulmonary nodule.


The alveolar cell carcinoma is a subtype of adenocarcinoma and arise within the alveoli producing areas of consolidation and the appearance resemble pneumonia. That's why in elderly who don't improve with pneumonia treatment we must consider alveolar cell carcinoma and do CT scan

2- Large cell carcinoma.

- Usually arise at **the periphery of the lung**.
- The growth is relatively rapid.


•CHEST X-RAY adenocarcinoma

This chest x-ray shows adenocarcinoma of the lung. There is a rounded light spot in the right upper lung at the level of the second rib. The light spot has irregular and poorly defined borders and is not uniform in density.



Radiographic features of large cell lung cancer

- Large-cell lung cancer of the lung typically presents as a large peripheral mass of solid attenuation and irregular margin. Focal necrosis can be present.



Types of lung cancers / 2


3- Squamous cell carcinoma

- They typically occur in **central bronchi**.
- Grow slowly and cavitate more often than any other cell types.


4- Small (oat) cell carcinoma.(WORST)

- Are usually central in location.
- Have the **fastest** rate of growth.
- Typically associated **with mediastinal adenopathy**.


Colored frontal X-ray of the chest of a male patient with squamous cell carcinoma (orange mass) in the upper lobe of the right lung.



Small cell lung cancer



Although not particularly S shaped, the collapse of the right upper lobe and a hilar mass are the components of Golden's S sign.



CXR demonstrates a large opacity in the right upper zone consistent with collapse of the right upper lobe secondary to a hilar mass. Note the hyperlucency of the hyperexpanded right lower and middle lobe, and elevation of the right hemidiaphragm

Golden S sign

- The Golden S-sign is seen on frontal chest radiographs. In the most common situation, the appearance is that of right upper lobar collapse (the right upper lobe appearing dense and shifting medially and upwards) with a central mass expanding the hilum. These two changes together form a reverse S-shape.


- Please check this video https://youtu.be/u7Ge-uNFdrk

الطب والجراحة لجنة

Chest X-ray showing a centrally located mass in the right lung, has an irregular margin indicating a malignant tumor



Chest X-ray showing a centrally located mass in the hilar region of the right lung that has an irregular margin indicating malignancy. There is also right lung collapse indicated by visible lung edge, and surrounded by *pneumothorax* indicated by absence of pulmonary vessels (no tension).
Note : this pneumothorax isn't related to the tumor, however, upon taking biopsy a tear of pleura could persist (3% chance) leading to this pneumothorax, for this reason after every chest biopsy a CXR is taken.



Edge of lung pointed out by red arrows

BENIGN TUMORS OF THE LUNG

❖Hamartoma.


- Is the most common benign tumor of the lung.
- Appear as solitary, well marginated,rounded mass.
- Calcification (popcorn) is present in 40% of cases (diagnostic feature)
- Fat is seen in up to 50% of hamartomas.

❖Adenoma.

- The vast majority of cases occur around the hilum and appear as round, well-defined nodule.


hamartoma x-rey

Small rounded lobulated dense nodular opacity at the upper zone of right lung.




hamartoma→ apical view

Small rounded lobulated dense nodular opacity at the upper zone of right lung.





Hamartoma sagittial CT

❖ *Solitary nodule at the right upper lung lobe apical segment showing lobulated outline, with internal hypodense foci of fat (CT density of –60 HU), as well as coarse 'popcorn' calcifications, with faint insignificant enhancement on post-contrast series.*




hamartoma ct axial popcorn






Chest X-ray showing a well defined mass in the middle lobe of the right lung with calcifications that are barely visible (arrow heads) .




Popcorn calcification(with in the blue circles) appears clear on CT scan and is diagnostic of hamartoma with no need of further evaluation

Lung metastasis

- The commonest primary tumors producing lung metastasis are breast, renal tract, thyroid, bone, and testicular tumors.
- Metastasis to the lung are usually bilateral and tend to be peripheral and more numerous at the lung bases.
- Lung metastasis appear as well-defined spherical nodules or masses (variable in size).





Cannonball lung metastasis(from a Renal Cell Carcinoma)
Numerous bilateral lung nodules and masses are noted, suspicious for metastatic disease.


Chest X-ray showing multiple bilateral nodules of spherical shape and well defined margins and a centrally located mass in the right lung indicating metastasis




Cardiomegaly

- On a standard PA chest, the heart size can be expressed as the cardiothoracic ratio.
- Generally, a ratio of over 50% of the heart size to the maximum internal diameter of the chest indicates cardiac enlargement.
- Additionally if the horizontal diameter of the heart is more than 16 cm we consider it enlarged regardless of internal diameter of chest

C is at the level of costophrenic angles or at level of mid part of diaphragm





PA chest radiograph showing cardiomegaly. The shorter red arrow (1) marks the width of the heart. The longer black arrow (2) marks the width of the thoracic cavity. As you can see, the width of the heart is greater than half the width of the thoracic cavity; therefore, the heart is enlarged (cardiomegaly). You can also see median sternotomy wires from previous surgery.

الطب والجراحة لجنة

Cardiomegaly / continuation

- Specific chamber enlargement may be identify on chest film:

Left atrium

double contour to the right heart border with splaying of the carina.

Right atrium

prominent of the right heart border.

Left ventricle

increased convexity of left heart border.

Right ventricle


upward displacement of the cardiac apex.

Right atrial enlargement : increased convexity of right heart border

Left ventricular enlargement : increased convexity of left heart border

Right ventricularl enlargement : displacement of apex above diaphragm. Normally apex is below diaphragm.


Left atrial enlargement :
1- most important sign is double contour sign (see next slide)
2- straightening of left heart border
3- bifurcation of trachea angle wider than 65 degrees and the left main bronchus horizontal rather than oblique (see next slide)




Left Atrial Dilatation :

•Causes : Mitral stenosis/ regurgitation, LVF, LA myxoma, …etc.

•Appearance: Straight left heart border, doubling of cardiac shadow right to the sternum, widening of the carina (Bifurcation of the Trachea), and the left primary bronchus is displaced upwards.





Left Ventricular Dilatation :

•Causes : Pressure overload: HTN & AS, Volume overload: VSD, AR, Aneurysm, Cardiomyopathy.


•Appearance: Left cardiac border gets enlarged and becomes more convex, Lt cardiac border dips into Lt dome of diaphragm, rounded apical segment, cardiophrenic angle is obtuse ((منفرج الزاوية

•Shmoo Sign.

الطب والجراحة
لجنة

Right Atrial Dilatation :

- Causes : Tricuspid stenosis/ regurgitation, ASD, AF, ...etc.
- Appearance: Rt. Cardiac border becomes more convex > 50% of right border, Rt. Atrial border extends below 3rd ICS.
- Measurement from mid vertical line to maximum convexity in Rt Border more then 5 cm in adult & 4cm in children.




Right Ventricular Dilatation :

•Causes : PV stenosis, Cor pulmonale, VSD, Tricuspid regurgitation & Secondary to LVF.

•Appearance: Elevation of the cardiac apex, Straightening/ convexity of left upper cardiac contour.




Note : this picture is from the web
https://radiopaedia.org/articles/double-density-sign-left-atrium
Left Atrial Enlargement
Left main bronchus
Right main bronchus
Double contour
Double contour

Heart failure

The radiological features are :

- **Cardiac enlargement.**
- **Upper lobe vascular dilatation.**
- (cephalization of blood flow): from raised pulmonary venous pressure.
- **Interstitial pulmonary edema:**
- accumulation of fluid in the interstitial space, in the interlobular septa with peripheral septal lines (kerley B lines)
- **Alveolar pulmonary edema**:
- accumulation of fluid in the alveolar spaces. The bilateral central edema is described as bat's wing.
- Vessels contour is hazy
- sometimes there is pleural effusion

1. Cardiomegaly.
2. Pulmonary oedema (pink):
■■ +/- the atypical ‘bat’s wing’ pattern shadowing.
3. Upper lobe vascular prominence (white):
■■ vessels in upper lobes appear larger than vessels in the lower lobes on an erect chest radiograph due to increased pulmonary vascular resistance in the lower zones due to dependent oedema.
4. Septal lines (light grey):
■■ due to fluid in the pulmonary lymphatics.
5. Pleural effusions (green):
■■ fluid leaks from the vessels into the pleural space






Figure 11.1 Diagrammatic representation of the potential radiological features of heart failure.



A patient with heart failure. There is bilateral diffuse ‘fuzzy’ opacification in the mid and lower zones in keeping with pulmonary oedema (pink). There is evidence of upper lobe vascular prominence (red) as vessels in the upper zone appear slightly larger than usual. There are also septal lines which are seen best in the right lower zone (yellow, surrounded by white dashed lines). This patient does not happen to have pleural effusions and the heart is not enlarged (although it is at the upper limit of normal).

الطب والجراحة
لجنة

Chest X-ray showing enlargement of the heart and dilation of upper lobe vessels (compared to lower lobe vessels) (cephalization of blood flow), presence of kerley B land A lines, hazy (not well defined) vessels contours, alveolar bilateral edema



Note : use "bring to front" and "send to back" feature to compare
Kerley B lines (blue) at the lvl of costophrenic angle 1-3cm
Kerley A lines (red) at the lvl of the hilum usually longer than B 1-5cm
Yellow is vessels and kerlry lines because it's branching

Chest trauma


- Chest trauma can be as blunt or penetrating.
- Penetrating chest trauma can injure vital organs such as the heart and lungs.
- The common clinical problems associated with chest injury include pulmonary contusion(hemorrahge in the lung), pneumothorax and hemothorax.
- Fractures of the lower ribs may be associated with diaphragmatic tears and spleen or liver injuries
- Fractures of middle ribs (4^{th} – 9^{th}) may be associated with pneumothorax, and lung contusion
- Fractures of the upper ribs can be associated with injuries to adjacent great vessels. If there is widening of mediastinum there is a high likely chance of great vessel injury and needs CT scan to confirm it (mediastinal hematoma)


Chest trauma / 2

What is the ABCDE approach to guide the radiographic search for thoracic injury ?

- Air : extra pulmonary
  1. Pneumothorax
  2. subcutaneous emphysema
  3. pneumomediastinum

  } Occur in penetrating injuries
- Bones – rib fracture, thoracic spine, scapula and sternum fractures.
- Contusions and lacerations in the lung.
- Diaphragm - rupture. Or tear
- Effusions – hemothorax.







The right 2nd, 3rd, 4th, 5th, 6th, and 7th ribs are fractured posteriorly (white circles). Remember, rib fractures may be easier to see if you rotate the radiograph through 90°.

Subcutaneous Emphysema


- On a chest radiograph, the gas appears as an area of low attenuation within the soft tissues.
- The skin of a patient with subcutaneous/surgical emphysema has a characteristic crackling feel, a sensation that has been described as similar to touching Rice Krispies. This sensation of air under the skin is known as subcutaneous crepitation.






Subcutaneous emphysema (yellow) projected over the left shoulder. In this case the patient was asthmatic and coughed violently, rupturing a bulla or bleb and allowing air from the lung to escape into the mediastinum. The air then tracked superiorly to the left shoulder.


الطب والجراحة لجنة






Subcutaneous emphysema (yellow) projected over both shoulder, neck, and upper mediastinum.

الطب والجراحة
لجنة



Obvious fracture
Depressed direction of neck of rib




Not fracture

Fractures 4th – 7th rib with contusion


Sometimes contusion could present without fractures in traumatic patients




Subcutaneous emphysema(air outside the lung) Indicating penetrating injury



Subcutaneous emphysema
Tension pneumothorax
Rib fractures
Collapsed lung


fracture
Tear in diaphragm
allowing stomach t
ascend upwards



Dextrocardia : right diaphragm is lower than left because of heart

Note : scoliosis

Chapter 4
Abdomen Imaging

Done By: Monther Qatawneh & Rakan Aljabsheh

Sources:

Doctor slides.

Abdominal X-ray for medical student, first edition.

Table of Content

Section 1: Imaging modalities 639
Section 2: AXR introduction 640
- AXR technique
- Plane Positions
- Normal AXR
- Quadrants and Regions
- Difference between S&L bowels
- AXR Checklist
- Presenting an Abd. Radiograph

Section 3: GIT Fluoroscopy 651
- Contrast medium in GIT studies
- Barium swallow
- Barium Meal
- Barium follow-through
- Barium enema

Section 4: Small Bowel Obstruction 661
- Introduction
- Classification
- Radiological signs of SBO
- Mechanical Obstruction
- Functional Obstruction
- Examples

Section 5: Large Bowel Obstruction 675
- Introduction
- Sigmoid volvulus

Section 6: Pneumoperitoneum 682
- Introduction
- Radiological signs
- Examples
- Apple core sign

Section 1: Imaging modalities

- Abdominal Ultrasound :
  - Is a noninvasive procedure used to assess the organs and structures within abdomen. It allows quick visualization of the abdomen. It's safe in children and pregnancy.
  - It's the first imaging technique uses to Dx & characterize abdominal pathology.
- Abdominal X-ray.
- Abdominal CT scan :
  - It's a diagnostic imaging test in abdomen
- Abdominal MRI.

In this chapter we will focus on Abdominal X-ray.

Section 2: Abdominal X-ray introduction

AXR technique :


- Abdominal X-ray (AXR) should be presenting boundaries from **the basal lung fields** to the **pubic bone**.
  - X-ray is toxic in children and pregnancy.
  - Liver, kidneys, spleen and bladder can’t always be identified due to intestinal loops overlapping that contain air + feces.
  - An AXR can be made in standing, supine and lateral positions (See next slide).

الطب والجراحة لجنة

Plane Positions


1) Standing position :

- *patient have to stand about 10 min before taking image.*
- Useful to see :Air under diaphragm (CXR could be used), air–fluid levels, may demonstrate dilated loops of bowel and Radiopaque foreign bodies.




2) Supine position :

- Useful to see : Dilated loops of bowel suggesting obstruction may be seen.
- air–fluid levels are not seen on the supine view.




3) Lateral position :

- Used when Patient can't stand or sit to do PA view.
- Useful to see :Air–fluid levels, Extraluminal free gas.




الطب والجراحة لجنة

Normal AXR – visceral structures




1. **Liver** (purple)
2. **Spleen** (pink)
3. Location of the **pancreas**



1. **Right kidney** (red)
2. **Left kidney** (red)
3. Location of **right ureter**
4. Location of **left ureter**
5. **Urinary bladder** (orange)
6. **Gas in the rectum** (green)
7. Location of **right adrenal gland**
8. Location of **left adrenal gland**
9. Location of **the gallbladder**

Note : Pancreas, ureters, adrenal gland and gall bladder: Not normally visualized.

Normal AXR - Bowel




- Different normal abdominal radiographs showing examples of the normal variation in bowel gas pattern.
- Note the stomach wall rugae (highlighted between the white arrows).

1. Stomach
2. Caecum
3. Ascending colon
4. Hepatic flexure
5. Transverse colon
6. Splenic flexure
7. Descending colon
8. Sigmoid colon

الطب والجراحة لجنة

Normal AXR – Skeletal Structure






1. Left 12th rib
2. Psoas outline – left and right
3. Vertebral body of L3
4. Pedicles of L1 vertebra
5. Right transverse processes of L1–5
6. Spinous process of L4
7. Sacrum
8. Coccyx
9. Right hemi-pelvis
10. Right sacroiliac joint
11. Right femur
12. Left femur

Quadrants and Regions

❖The abdomen can be divided into four quadrants or nine regions:

(RUQ) : Right upper quadrant
(LUQ) : Left upper quadrant
(RLQ) : Right lower quadrant
(LLQ) : Left lower quadrant


1. Right hypochondriac
2. Epigastric
3. Left hypochondriac
4. Right lumbar
5. Umbilical
6. Left lumbar
7. Right iliac
8. Suprapubic
9. Left iliac




الطب والجراحة
لجنة

Difference between Small & Large bowels

	Small bowel (Figure A)	**Large bowel** (Figure B)
Location	Central	Peripheral
Mucosal fold	Valvulae conlventae (Small Figure)	Haustration (Small Figure)
Diameter	<3 cm	<6 cm (cecum:<9cm)
Fluid level/feces	2 or 3 fluid level normal	Solid feces normal
Cause of obstruction	**Adhesion**, hernia, tumor, Gall stone	**Tumor**, abscess, diverticulitis, volvulus.





AXR Checklist - ABCDE

- The following structures should be checked in 2 views of AXR :

A: Air in the wrong place :

- Look for pneumoperitoneum and pneumoretroperitoneum & Look for gas in the biliary tree and portal vein.
- Normally we can see air fluid level in stomach (always), two to three (not more) in Small bowel is acceptable & none (free) in Large bowel. it's seen in standing and lateral positions.

B: Bowel :

- Look for : Dilated small and large bowel, volvulus , distended stomach, hernia, evidence of bowel wall thickening.

C: Calcification :

- Look for clinically significant calcified structures.

D: Disability (bones and solid organs) :

- Look at the bony skeleton for fractures and sclerotic/lytic bone lesions, look at the spine for vertebral body height, alignment, pedicles and a 'bamboo spine', solid organ enlargement.

E: Everything else :

- Look for : Evidence of previous surgery and other medical devices, foreign bodies, the lung base.

الطب والجراحة لجنة

AXR Checklist - **ABDO X**

- An alternative mnemonic to the **ABCDE** is the **ABDO X**

A: **A**ir - where it should and shouldn't be

B: **B**owel - position, size and wall thickness

D: **D**ense structures, calcification and bones

O: **O**rgans and soft tissues

X: e**X**ternal objects and artifacts

الطب والجراحة لجنة

Presenting an Abdominal Radiograph

You should present an abdominal radiograph in a systematic way to ensure you cover all areas and do not miss anything important. This is how you should present:

1) Give the type of radiograph.
2) Give the patient's name.
3) Give the date the radiograph was taken.
4) Briefly assess the radiograph quality.
5) Run to ABCDE of abdominal radiograph.
6) Give the short summary at the end

e.g. " this is an **AP supine abdominal radiograph** of Monther Qatawneh, taken on the 1st of January, 2022.

Section 3: GIT Fluoroscopy

Contrast medium in GIT studies:

- **Barium sulfate :**
  - is the most commonly used in barium studies of the GIT.
- **Gastrografin :**
  - is used in: Suspected bowel perforation, post-operative for leakage, meconium ileus (in neonates).
  - It is contra-indicated in Barium swallow and meal if the patient is at risk of aspiration such as in infants, because gastrografin can induce pulmonary edema and chemical pneumonitis.
- **Non ionic contrast medium:**
  - It's used in barium swallow studies in infants.


الطب والجراحة لجنة

1) Barium swallow :

- Is the examination of the esophagus with contrast medium.
- **Indication** :
  - Symptoms of gastro-esophageal reflux (most common)
  - Dysphagia which could be related to :
    - Esophageal stricture
    - Esophageal tumor (important)
    - Suspected achalasia
    - Vascular abnormalities
    - Esophageal web

الطب والجراحة لجنة

1) Barium swallow :



Multiple images of Barium swallow study:

A: Normal radiograph.

B: Tubular structure with filling defect and mass lesion.

C: Apple eaten sign; suspected esophageal cancer.


D: Dilated end of the esophegus (bird beak sing); Suspected achalasia.

2) Barium Meal :

- Is the examination of the Stomach & Duodenum with contrast medium.
- **Double contrast** is obtained by introduction of gas into the stomach using effervescent powders and barium.
- **Indication** :
  - Gastro esophageal reflux
  - Gastric or duodenal ulcer
  - Hiatus hernia
  - Suspected gastric tumor

الطب والجراحة لجنة

2) Barium Meal :



Multiple images of Barium meal study:
A: Normal radiograph.
B & C: Suspected Hiatus Hernia

الطب والجراحة لجنة


3) Barium follow-through :

- Is the examination of the Small bowel with contrast medium.
- Abdominal films are taken **every half hour** until barium reaches the large bowel .
- **Indication** :
  - Inflammatory bowel disease, most often Crohn's disease (most common)
  - Small bowel tumors / lymphoma
  - Small bowel obstruction

الطب والجراحة لجنة

3) Barium follow-through :





Multiple images of Barium follow-throw study:
A: Normal radiograph.
B: multiple strictures and stenosis and mostly there is complete cut in Crohn's disease.

4) Barium enema :


- Is the examination of the Large bowel with contrast medium.
- Barium is run into the colon by Foleys catheter placed in the rectum.
- Air insufflation into the large bowel produces a double contrast examination.
- **Indication** :
  - Investigation of abdominal mass.
  - Large bowel obstruction : Suspected colonic tumor, diverticular disease & volvulus.

الطب والجراحة لجنة

4) Barium enema :







Multiple images of Barium enema study:

A: Normal radiograph.

B: We see diverticulosis in descending colon.

C: We see a stricture (apple eaten sign); suspected colon cancer.

Section 4: Small Bowel Obstruction (SBO)


- It’s covered the lecture and the seminar.

Introduction

- Small bowel obstruction (SBO) accounts for 80% of all mechanical intestinal obstruction.
- It has a mortality rate of ~5%
- **Clinical presentation:** Classical presentation is: Cramping abdominal pain, abdominal distension with nausea and vomiting.
- Radiographic findings can be evident 6-12 hours before the onset of clinical symptoms.


الطب والجراحة لجنة

Classification



Radiological signs of SBO

- Dilatation of small bowel loops more than 3 cm.
  - In Mechanical obstruction : Proximal dilatation and distally collapsed.
- Multiple air fluid levels (more than 3).
- Absence of gas in the rectum.
  - If gas is still present, it indicate that the obstruction is recent or that it is incomplete.




This radiograph represent SBO, we see dilation more than 3 cm (Black arrow), multiple air fluid level (red arrows) and absence of gas in rectum (blue arrow).

Mechanical Obstruction

1) Adhesions :


- Are bands of fibrous or scar-like tissue that form between abdominal tissues and organs.
- Its not accurate seen in X-ray so we depend on the history and the CT scan.
- Are the most common cause of mechanical obstruction, about 70% of cases.



Mechanical Obstruction

2) Strangulated Hernia :

- Occurs when a section of the small intestines pushes through a weakened area of the abdominal muscle. The surrounding muscle then clamps down around the tissue, cutting off the blood supply to the small intestine.




Mechanical Obstruction


3) Volvulus: (Figure 1)


- An abnormal twisting of a loop of small bowel around the axis of its own mesentery.

4) Intussusception: (Figures 2,3)

- A serious condition in which part of the intestine slides into an adjacent part of the intestine , one segment of intestine "telescopes" inside of another.







Functional Obstruction (Paralytic ileus)

Definition : Temporarily impaired peristalsis of the gastrointestinal tract in absence of mechanical obstruction.

Causes : (Mnemonic: 5P)

Postoperative, Parturition, Peritonitis, Pelvic and spinal fractures & Potassium level decrease.

Types :

1. **Generalized:** Affect all loop of small & large bowel.
2. **Localized:** Affect only one or two loops of small intestine.

الطب والجراحة لجنة

Functional Obstruction (Paralytic ileus)

Pathophysiology :

1. Stressful stimulation of the bowel → Activation of the sympathetic system
2. Inflammation release NO locally → Relaxation of the intestinal smooth muscle

} Decrease peristalsis will cause bowel distention which progress to mechanical obstruction.

Abdominal X-ray AP view :


1- Multiple dilated loop of bowel.

2- Several air fluid level.

3- Visible gas in rectum.

الطب والجراحة لجنة

Functional Obstruction (Paralytic ileus)



Multiple images of Paralytic ileus :

A: Multiple small bowel dilatation (black arrow) with multiple air fluid level (red arrow).

B: **Generalized adynamic ileus :** Dilated loop of large bowel (white arrows), Dilated loops of small bowel (dotted white arrows) with Gases in rectum (black arrows).


C: AXR – left lateral view : Significantly distended intestinal loops with air fluid levels are visible.

Special Case: Gallstone ileus

- A gallstone ileus is an uncommon cause of mechanical small bowel obstruction. Recurrent episodes of cholecystitis cause adhesion of the gallbladder to the bowel (usually duodenum) and eventually a fistula forms. A large gallstone then enters the bowel and causes obstruction, typically at the ileocecal valve.
- A gallstone ileus gives the classical Rigler's triad:
    1. Pneumobilia
    2. Small bowel obstruction
    3. Gallstone (usually in the right iliac fossa, but only seen in approximately 30% of cases)


الطب والجراحة لجنة

Special Case: Gallstone ileus




There are branching dark lines (gas) projected over the center of the liver, larger and more prominent towards the hilum, in keeping with pneumobilia. There are centrally located loops of bowel measuring >3cm in diameter with valvulae conniventes seen throughout, in keeping with dilated small bowel. There is a calcified opacity projected over the left sacrum in keeping with a large calcified gallstone. The right radiograph shows the pneumobilia marked in dark blue, dilated small bowel marked in blue and gallstone marked in yellow.

Example #1



Example #1: The bowel is visible as there is gas (black) within. You can tell that it is small bowel as it is centrally located and valvulae conniventes can be seen throughout. The loops measure >3cm in diameter therefore they are dilated. The right radiograph shows the dilated small bowel marked in blue.

Example #2



Example #2: The loop of bowel is visible as there is gas (black) within. You can tell that it is small bowel as valvulae conniventes can be seen throughout. The loop measures>3cm in diameter and is therefore dilated. When a single dilated loop is seen (as in this case) it is known as a **sentinel loop**. It is a feature that is occasionally due to a localised ileus from nearby inflammation causing local paralysis and accumulation of gas in the intestinal loop. The right radiograph shows the dilated small bowel marked in blue.

الطب والجراحة لجنة

Section 5: Large Bowel Obstruction

Introduction

- Large bowel distension is almost always due to **large bowel obstruction**. The bowel proximal to the obstruction is dilated and the bowel distal to the obstruction is usually collapsed.
- Causes of large bowel obstruction include the following:
  - **Malignancy** (colorectal carcinoma is the most common cause of large bowel obstruction in adults)
  - **Diverticular stricture**
  - **Faecal impaction** (most common cause in immobile elderly persons)
  - **Volvulus** (*examples in next few slides*)


الطب والجراحة لجنة

Notes to keep in mind

- **Haustra:** These are the small pouches in the wall of the large intestine. The taenia coli (ribbons of smooth muscle which run along the length of the colon)
- smooth muscle which run along the length of the colon) are shorter than the colon itself, therefore the colon becomes sacculated between the taenia coli forming the haustra. The lines between the haustra are called **haustral folds** and typically do not cross the entire width of the bowel (unlike valvulae conniventes).
- **Note:** If the bowel is grossly distended, then the haustra may not be seen.

الطب والجراحة لجنة

Sigmoid volvulus



- A volvulus is the **twisting of the bowel** on its **mesentery**. It causes partial or complete bowel obstruction. The two commonest types of volvulus in adults are **sigmoid volvulus** and **caecal volvulus**.
- A **volvulus** can give symptoms by **two** processes:

1. **Bowel obstruction:** The loop of twisted bowel causes a ‘closed-loop’ obstruction.
2. **Bowel ischemia:** In some cases the twisting of the bowel mesentery compromises the vascular supply to the bowel leading to ischemia and eventually necrosis, which can be fatal.


الطب والجراحة لجنة

Sigmoid volvulus

- A sigmoid volvulus is caused when the **sigmoid colon twists** on its **mesentery**. It is usually seen in the elderly or institutionalized patients.
- Radiological signs of a sigmoid volvulus:
- **Coffee bean sign:** The shape of the distended gas filled ‘closed loop’ of colon looks like a large coffee bean. (see figure below)
- **General lack of haustra:** Often the bowel is so distended that haustra flatten out and are no longer seen.
- **Distension of the ascending, transverse and descending colon:** The colon proximal to the obstruction (volvulus) is often distended, but not always.



Sigmoid volvulus




Multiple images Show a Sigmoid Volvulus:

A: Diagrammatic figure represent coffee bean sign.

B&C: loss of haustra & coffee bean sign.

Sigmoid volvulus



Shape of dilated displaced colon appears similar to coffee bean.
Northern exposure sign represents dilated sigmoid colon that extends cranial to the transverse colon(star).
Above two features on X-ray Abdomen are suggestive of sigmoid volvulus.

Section 6: Pneumoperitoneum

- It’s covered the lecture and the seminar.

Introduction


- Pneumoperitoneum literally means **free gas in the peritoneal cavity**. It usually indicates **bowel perforation**. Free gas may also be seen up to 3 weeks after abdominal surgery and in trauma (e.g. stabbing).
- **Main causes of pneumoperitoneum:**
  - Perforated peptic ulcer
  - Perforated appendix/bowel diverticulum
  - Post-surgery
  - Trauma
- **Note:** An abdominal radiograph and an erect chest radiograph are requested together when looking for a pneumoperitoneum. This is because an erect chest radiograph is very sensitive for detecting free abdominal gas since it can detect as little as 2–3 ml. On an erect chest X-ray the free gas is seen as a rim of blackness beneath and very closely opposed to the curve of the diaphragm.

الطب والجراحة لجنة

Radiological signs of Pneumoperitoneum

Figures A & B:

- Two identical erect radiographs of the lower chest. The lower radiograph shows the pneumoperitoneum marked in turquoise.
- Meaning that there is air under the diaphragm.



- **Radiological signs of pneumoperitoneum include:**
  1. Rigler's sign (also called the double wall sign)
  2. Gas outlining the liver
  3. Falciform ligament sign (also called silver sign)

Radiological signs of Pneumoperitoneum


- **Rigler's sign:** Also known as the **double-wall sign**, this is seen when gas is present on both sides of the intestinal wall (i.e., gas within the bowel *and* free gas in the peritoneal cavity). Normally the bowel wall is only just visible, outlined by the gas within the bowel and peritoneal fat outside of the bowel. With a pneumoperitoneum the bowel wall is easily seen as it is outlined by gas within the bowel and gas outside of the bowel.

الطب والجراحة لجنة


Radiological signs of Pneumoperitoneum

1. Diagrammatic representation of normal appearances of the bowel wall. The lumen of the bowel contains gas. You can see the bowel wall, but there is little contrast between the bowel wall and the peritoneal fat outside of the bowel.

2. Diagrammatic representation of **Rigler's sign** (double-wall sign). The lumen of the bowel contains gas, and there is also gas within the peritoneal cavity. The bowel wall is therefore clearly seen outlined by the gas either side.




Examples #1



Example #1: two identical abdominal radiographs showing a pneumoperitoneum. There are loops of bowel with gas outlining both sides of the bowel wall in keeping with Rigler's sign. The right radiograph shows in turquoise and brown the areas where Rigler's sign is most clearly seen. The lumen of the bowel is marked in brown and the free gas outlining the bowel wall marked in turquoise. The best example of Rigler's sign is marked with a white circle. An area of normal appearing bowel wall is marked with a white dashed circle for comparison. (You can also see dilated loops of large bowel.) الدائرة بخطوط متقطعة طبيعي بينما الدائرة الكاملة تبين العلامة

الطب والجراحة لجنة

Examples #2




Example #2: showing a large pneumoperitoneum. There are loops of bowel with gas outlining both sides of the bowel wall in keeping with Rigler's sign. The right radiograph shows in turquoise the areas where the pneumoperitoneum is most clearly seen. Where Rigler's sign is most clearly seen, the lumen of the bowel is marked in brown. The best example of Rigler's sign is marked with a white circle. You can also see **gas outlining the liver as shown by the white line**.

هون الدائرة البيضاء توضح اكثر مكان مبينة فيه علامة ريغلر بوضوح


الطب والجراحة لجنة

Examples #3




Example #3: similar to the previous examples

Examples #4




Example #4: There are loops of bowel with gas outlining both sides of the bowel wall in keeping with Rigler's sign, and there is **gas outlining the falciform ligament in keeping with the falciform ligament sign.** The right radiograph shows in turquoise and brown the areas where Rigler's sign is most clearly seen. **Notice the position of the falciform ligament is shown with white arrows.**

Examples #5



Example #5: gas outlining the falciform ligament in keeping with the falciform ligament sign and there is also gas outlining the liver. The right radiograph shows in turquoise the areas where the pneumoperitoneum is most clearly seen.

Examples #5



Example #6: free intraperitoneal gas is seen easily because it is contrasted against the liver, indicating pneumoperitoneum seen in a neonate in left lateral decubitus position.

Apple core sign

- The “**apple core sign**” was first described in barium enema examinations of the colon. The apple core appearance is the visual manifestation of an annular **lesion of the** bowel with irregular overhanging edges and shouldered margins. The apple core sign is **classically seen in cases of colon carcinoma.**



Chapter 5
Skeletal System Imaging

Done By: Laith Faiz Al-Najada

Sources:

Dr. Mahmoud Khasawneh slides; Mutah University

Table of Content

1. Skeletal system overview
2. Imaging modalities
3. Pathology
4. Systemic approach in cervical spine x-ray
5. Bone fractures

Writer’s notes

اللون الأحمر: مهم

اللون الأخضر: للقطاعات بس (مشروع تأويت)

اللون الأزرق: شرح الأشياء اللي مش واضحة

الوان الأسهم (لتوضيح الصور) متناسبة مع الوان الكلام، يعني الكلام باللون البنفسجي، السهم اللي اله علاقة بالكلام راح يكون بنفسجي كمان

الطب والجراحة لجنة

Section 1: Skeletal system overview

Skeletal system

- What is the skeleton?
  - The skeleton is the complete set of bones that make up a human.
  - The average adult human skeleton has about 206 bones.
  - The skeleton can be divided into axial skeleton and appendicular skeleton.

The skeletal system is composed of all the bones in the body and the tissues such as tendons, ligaments and cartilage that connect them.

These bones meet at joints, the majority of which are freely movable, making the skeleton flexible and mobile.

الطب والجراحة لجنة


Skeletal system

- What is the axial skeleton?
  - The axial skeleton is the central portion of the bony skeleton comprising the head, neck and trunk (80 bones in total). It has many functions including housing and protecting the central nervous system as well as the organs of the chest, abdomen and pelvis.



Skeletal system

- What is the appendicular skeleton?
  - The appendicular skeleton is the portion of the bony skeleton that includes and supports the limbs. It includes the shoulder girdle (clavicles and scapulae) and the bony pelvis, connected to the axial skeleton centrally and is composed of 126 bones in total.



Section 2: Imaging modalities

الطب والجراحة
لجنة

Imaging modalities

❖Plain film:

- Plain films still remain the mainstay of radiological investigation of the skeletal system.
- views should always be obtained in two projections (frontal and lateral view).

❖Ultrasound:

- neonatal hip for congenital dislocation.
- soft tissue lesions, abscesses and masses.
- joint effusions.

الطب والجراحة لجنة

Imaging modalities

- CT scan:
  - It’s very helpful for assessment of bone tumors before surgery.
  - Evaluation of certain fracture such as the acetabulum, spine and calcaneum.
  - Study of the spinal column.

- MRI:
  - Assists in the investigation of bone tumors and in soft tissue and joint masses.
  - It’s extremely sensitive in injuries to cartilage, muscle, ligament, menisci, and tendon.

الطب والجراحة لجنة

Section 3: Pathology

الطب والجراحة لجنة

ملاحظة:
الموضوع دايماً يحذفوه وما يدخل بالامتحان


Osteoporosis

- **Is a generalized decrease of bone mass.**
- **Radiological investigation:**
  - Bone densitometry.
  - Plain x-ray film:
- Detection of osteoporosis on plain film requires a reduction in bone mass of at least 30%.
- Osteoporosis result in a loss of bone density, decrease in the number of trabeculae and coarse striations.


الطب والجراحة لجنة

An osteoporosis case

60-year-old female complaining of repeated vertebral fractures and chronic back pain.




Frontal view



Lateral view

Osteoarthritis
(Degenerative joint disease)

- Is a degenerative condition affecting the articular cartilages and subjacent bone.
- It's part of normal ageing process ('wear and tear'), but it can also results from trauma or infection.
- Any joint may be affected, but the knees, hips, shoulders and hands are frequently involved.




Osteoarthritis
(Degenerative joint disease)


- **Radiological features:**
  - Joint space narrowing (the most important sign).
  - Subchondral sclerosis: dense line in bone due to necrosis
  - Osteophytes formation: reactive bony outgrowths at joint margin due to inflammation.
  - Loose bodies: osteophytes at joint space as a result from separation of it from cartilage.
  - Subchondral cyst.
  - Joint erosions.

الطب والجراحة لجنة

Osteoarthritis
(Degenerative joint disease)





Narrow joint space,
osteophytes, sclerosis.




Normal joint

Osteoarthritis
(Degenerative joint disease)





Osteoarthritis
(Degenerative joint disease)



Narrow joint space, osteophytes, sclerosis, loose body, also there is moderate dislocation.

ملاحظة:
مطلوب للفاينل بس


Osteomyelitis

- Infection of marrow and bone caused by *S. aureus* occurs usually in children.
- Sign and symptoms:
  - bone pain with fever and leukocytosis.
  - Soft tissue swelling due to edema, redness and pain.
- Radiological feature:
  - May be normal for up to 10 days.
  - Periosteal reaction: nonspecific radiographic finding that indicates periosteal irritation.
  - Bone destruction.

الطب والجراحة لجنة

ملاحظة:
مطلوب للفاينل بس

Osteomyelitis





Malignant bone tumors

- They are destructive lesions, often associated with periosteal reaction, and have a wide zone of transition between normal and abnormal bone.

- The most common malignant bone tumour is a metastasis and it's often solitary.

- The most common primary malignant bone tumors are:
  - Osteogenic sarcoma
  - Ewing's tumour
  - Chondrosarcoma
  - Fibro sarcoma
  - Giant cell tumor

الطب والجراحة لجنة

Malignant bone tumors

- CT and MRI are the best imaging modalities to evaluate tumours and determine bone and soft tissue involvement.

- Radiological feature:
  - Plain film shows an area of bone destruction.
  - CT / MRI show:
    - Tumor vascularity.
    - Infiltration of surrounding tissues.
    - Relationship to nerves or vessels.

الطب والجراحة لجنة

Bone metastasis

- Are the most common malignant bone tumors.
- Any primary tumor may metastasize to bone, but the most frequents are:
  - Breast cancer: usually lytic in nature but may be sclerotic (dense) or mixed.
  - Prostate cancer: the vast majority are sclerotic (90% dense).
  - Lung, kidney, and thyroid cancer: lytic lesion.
  - Adrenal cancer: predominantly lytic lesion

الطب والجراحة لجنة

Bone metastasis




Multiple well defined lytic lesion involve all the skull and decrease bone density with no pathological fracture

DDx:

1) Metastasis from lung cancer in male or breast cancer in female.
2) Multiple myeloma most commonly in old age (discussed later at this section).

الطب والجراحة لجنة

Bone metastasis




Normal iliac bone (left)
Large lytic lesion in iliac bone (right)

Frontal view of pelvis X-ray, showing pathological dislocation of Rt. hip joint Because of large lytic lesion in Rt. Iliac bone

DDx:
1) Malignant lesion.

Bone metastasis



Normal bowel gas (not lytic lesion)
Sclerotic lesion (hyperdense)

Frontal view of pelvic and abdomen region x-ray showing:
Diffuse metastasis involve all skeleton, hyperdense vertebral bodies and scleroting mets.

DDx:
1) Metastasis from prostate cancer.
2) Metastasis from breast cancer if female.

الطب والجراحة لجنة

Multiple myeloma

- Malignant proliferation of plasma cells in the bone marrow.
- Most common primary malignancy of bone.
- Signs and symptoms (Old CRAB):
  - Old age.
  - Calcium elevated (hypercalcemia).
  - Renal failure.
  - Anemia.
  - Bone lytic lesion.
- The most common bones involved are the skull, spine, pelvis, and ribs.
- The disease may occur in a disseminated form, or as a localized solitary mass (Plasmacytoma).

الطب والجراحة لجنة

Multiple myeloma

- At time of presentation, 80% have skeletal abnormalities.
- Radiological features on plain film:
  - Generalized osteoporosis (decrease bone density).
  - Compression fractures of the vertebral bodies.
  - Scattered multiple lytic lesions with well defined margins.

الطب والجراحة لجنة

Multiple myeloma



Generalized osteoporosis



Compression fracture




Scattered multiple lytic lesions


Section 4: Systemic approach in cervical spine x-ray

Systemic approach in cervical spine x-ray

- Standard views:



Frontal view



Lateral view

Systemic approach in cervical spine x-ray


- How to read cervical spine x-ray? (ABCD)
  - Alignment
  - Bodies
  - Cortical outline
  - Disk spaces & Density

الطب والجراحة الجنة

Systemic approach in cervical spine x-ray

❖Alignment

- All lines should be smooth and continuous.
- Needed to see if there is dislocation (it's significant in the spine).
- The vertebral bodies should be all intact between anterior spinal line and posterior spinal line.



Anterior spinal line

Posterior spinal line

Spinolaminar line

الطب والجراحة لجنة

Systemic approach in cervical spine x-ray

❖Bodies

- Height of vertebral body indicates compression fracture.
- Check that each vertebral body is correspond to it's spinal process.



Systemic approach in cervical spine x-ray

- **Cortical outlines**
  - Check for any fracture.



Systemic approach in cervical spine x-ray

❖Disk spaces & Density

- The intervertebral disk spaces should be the same.




Section 5: Bone fractures

Normal X-ray Review


ملاحظة:
للاطلاع فقط

- Normal x-ray of upper limb:



Shoulder AP view

the axillary view is **useful for detecting anterior or posterior shoulder dislocations that are not evident** in the AP view. The x-ray beam is directed into the axilla from below (inferior to superior), or down through the axilla from above the shoulder (superior to inferior)




Shoulder axillary view

الطب والجراحة لجنة

ملاحظة:
للاطلاع فقط

Normal X-ray Review

❖Normal x-ray of upper limb:



Forearm lateral view




Forearm AP view

الطب والجراحة الجنة


ملاحظة:
للاطلاع فقط

Normal X-ray Review

❖Normal x-ray of upper limb:



Elbow AP view



Elbow lateral view

الطب والجراحة لجنة


ملاحظة:
للاطلاع فقط

Normal X-ray Review

- Normal x-ray of upper limb:



Wrist PA view




Wrist lateral view


ملاحظة:
للاطلاع فقط

Normal X-ray Review

- Normal x-ray of lower limb:



Pelvis AP view




Leg AP view

الطب والجراحة الجنة


ملاحظة:
للاطلاع فقط

Normal X-ray Review

- Normal x-ray of lower limb:



Knee AP view




Knee lateral view

الطب والجراحة الجنة

ملاحظة:
للاطلاع فقط

Normal X-ray Review

❖Normal x-ray of lower limb:



Ankle AP view



Ankle lateral view



Foot AP view



Foot lateral view


Bone fractures

- Plain films are the initial evaluation of a patient with suspected skeletal trauma.
- At least two views (AP and lateral views) should always be obtained.
- In any significant head or spine injury, CT scan is the initial investigation.
- CT will detect fractures as well as underlying intracerebral hemorrhage or contusions.


الطب والجراحة لجنة

Bone fractures

- A fracture is a partial or complete break in the continuity of bone or cartilage tissue.
  - **Closed fracture:** is a break to the bone that does not damage surrounding tissue or tear through the skin.
  - **Open fracture:** is one that damages surrounding tissue and penetrates the skin, open fractures are generally more serious than simple closed fractures due to the risk of infection.



Closed fracture



Open fracture

Bone fractures

- Causes:
  - Physical trauma.
  - Overuse.
  - Healthy conditions that weaken the bone such as osteoporosis.

- Signs and symptoms:
  - Pain.
  - Swelling.
  - Bruising.
  - Bleeding if it’s an open fracture.
  - Dizziness and nausea in severe cases.

الطب والجراحة لجنة


Bone fractures

- Types of fracture:
  - **Linear Fracture.**
  - **Comminuted Fracture.**
  - **Avulsion Fracture.**
  - **Pathological Fracture.**
  - **Greenstick Fracture.**
  - **Compression Fracture.**
  - **Depressed Fracture.**
  - **Epiphyseal Plate Fracture.**
  - **Stress Fracture.**
  - **Spiral Fracture.**
  - **Oblique Displaced/Non-Displaced Fracture.**

الطب والجراحة لجنة

Bone fractures

❖**Linear Fracture:** break in the bone, but without fragments of dislocation.








الطب والجراحة لجنة

Bone fractures

❖**Comminuted Fracture:** the bone is broken or crushed into several pieces (more than 2 pieces), also known as multi-fragmentary fracture.








Bone fractures

❖**Avulsion Fracture:** when tendon or ligament pulls small fragment of the bone away from the main part of bone.









Bone fractures


❖**Pathological Fracture:** caused by an underlaying disease (anemia, osteoporosis, metastasis, ..) that weakens the bones.



Tumor mass



Osteolytic lesion in head of humerus
(Bone Cyst)




Osteolytic lesion in head of humerus
(Bone Cyst)

Bone fractures

❖**Greenstick Fracture:** it's an incomplete fracture of bone (fracture at one side of bone margin without separation of whole length), the bone may also buckle without an actual break, usually happen in children.










Bone fractures

❖**Compression Fracture:** force is applied in the longitudinal axis of bone, usually occurs in the spine (decrease in the height of the bone).








الطب والجراحة لجنة

Bone fractures

- **Depressed Fracture:** is a break in a cranial bone (or "crushed" portion of skull) with depression of the bone in toward the brain, usually due to blunt force trauma to the head, detected by CT scan.



Bone fractures

❖**Epiphyseal Plate Fracture:** it's a fracture in growth plate of a child.

Type I Fractures

These fractures break through the bone at the growth plate, separating the bone end from the bone shaft and completely disrupting the growth plate.

Type II Fractures

These fractures break through part of the bone at the growth plate and crack through the bone shaft, as well. This is the most common type of growth plate fracture.

Type III Fractures


These fractures cross through a portion of the growth plate and break off a piece of the bone end. This type of fracture is more common in older children.

Type IV Fractures

These fractures break through the bone shaft, the growth plate, and the end of the bone.

Type V Fractures

These fractures occur due to a crushing injury to the growth plate from a compression force. They are rare fractures.



Salter-Harris Epiphyseal Fracture Classification

*Physis (growth plate) is highlighted in blue. Fracture line is black or red.

fpnotebook.com

الطب والجراحة لجنة

Bone fractures

❖**Epiphyseal Plate Fracture:** it's a fracture in growth plate of a child.





Growth plate fracture with complete dislocation of fracture fragment

الطب والجراحة لجنة

Bone fractures

- **Stress Fracture:** fractures occurring in the bone due to a mismatch of bone strength and chronic mechanical stress placed upon the bone, they are common in athletes, especially runners and military recruits, the most common bone is the tibia.






Bone fractures

❖Stress Fracture:


- Plain radiographs have poor sensitivity (15-35%) in early-stage injuries, which increases in late-stage injuries (30-70%), due to possible callus formation.
- MRI is the most sensitive modality for diagnosis of a stress fracture



الطب والجراحة لجنة

Bone fractures

- **Spiral Fracture:** complete fracture of long bone that result from a rotational force applied to the bone and are likely to be associated with displacement.





Bone fractures


❖**Oblique Fracture:** the break is on an angle through the bone.






Bone fractures

❖**Additional Fractures in case of spinal injury:**



Burst Fracture (As comminuted fracture)



Fracture dislocation (Abnormal alignment)

Chapter 6
Urinary Tract Imaging

Done By: Salah Saed

Sources:

- DR. Hana Quidsieh slides 2021/2022; Mutah University

Writer’s notes

- الدكتور بالميني اوسكي بجيب أسئلة من الصور والاسئلة مش صعبة ,, ركزوا على الصور ومروا عالكتابة حتى تفرقوا بين كل صورة والأخرى

Table of Content

- **Urinary tract investigations:**

1. Plain film (KUB) : kidney , ureter, bladder.
2. Ultra Sound
3. CT of urinary tract
4. MRI
5. Intravenous Urography (IVU) // Intravenous pyelography (IVP)
6. Micturating cytogram (MC)
7. Renal Arteriography

- **Urinary tract pathology :**

1. Congenital Renal Anomalies
2. Urinary tract caliculi
3. Cystitis
4. Acute pyelonephritis
5. Chronic pyelonephritis
6. Urinary obstruction
7. Benign renal lesion
8. Polycystic kidney disease
9. Benign renal tumours
10. Malignant renal tumour

Plain film(KUB):

- **The kidneys, ureters, bladder (KUB) radiograph** is optimized for assessment of the urogenital system, and should not be confused with the AP supine abdomen view.

- **Patient position**
- the patient is supine, lying on their back, either on the x-ray table (preferred) or a trolley
- patients should be changed into a hospital gown, with radiopaque items removed (e.g. belts, zippers, buttons, ECG electrodes)
- the patient should be free from rotation; both shoulders and hips equidistant from the table/trolley
- the x-ray is taken on full inspiration
  - this causes the diaphragm to contract, hence compressing the abdominal organs, allowing all renal contents to be visualized on a single image

الطب والجراحة لجنة

Plain film(KUB):

Plain film useful in :

- Renal calculi or calcification
- Stones in the ureters
- Bladder calcifications and calculi
- Bone abnormality or metastasis

#normally ureter cant be seen on plain abdomen except in IVU.

#ureters can be seen above the transverse processes of lumber V.


Urinary tract calculi

Renal stones	Plain film
1. Stones composed **Calcium (majority about 90 %)**	(radio-opaque) .
2. Stones composed of **Uric acid**	not visible (radiolucent)
3. Stones composed of **Cystine**	minimally dense (Semi-opaque)

Urinary tract calculi / continuation

What is the initial imaging test usually ordered to find urinary tract stones ?

- ✓ **Plain radiograph (KUB)**, because the majority of stones are radio-opaque
- ✓ Other calcifications may be confused with urinary tract stones such as a phlebolith in the pelvis , which is a venous calcification , often with a lucent center .




KUB

Normal kidneys, ureters and bladder.

Renal calyces:

On hydronephrosis there is lost of cupping of the renal calyces.



KUB

Small radio-opaque stone in Right renal pelvis with multiple small phleboliths in left lower pelvis.

phlebolith:

Is a calcification in the pelvic veins and it's small look like stone but mostly it contain small black center due to presence of fat.

Its bellow the level of ischial spine.but stone mostly above the level of ischial spine.



KUB

Radio-opaque stone at the middle portion on left ureter



KUB

Semi-opaque stone at the site of insertion of left ureter to the bladder (vesico-uretric junction).

Stone in ureter:

Site of hidden area of stone in ureter are:

1. *ischial bone.*
2. *Pelvis near bladder.*

الطب والجراحة لجنة



KUB

Radio-opaque stone in the left renal pelvis with semi-opaque stone in left renal pelvis too, and small phlebolith.

Ultrasound of the urinary tract

- Ultrasound is one of the most valuable investigations of the urinary tract and the investigation of choice in children.

- non-invasive procedure useful in distinguishing the etiology for renal failure.
- It is critical for ruling out obstruction

- It is very effective in evaluating:


-renal size

-Masses

-renal obstruction

-Bladder residual volume

-prostatic size.



Normal kidney ultrasound

- Blood flow to kidneys can also be evaluated with **doppler US**

Ultrasound of the urinary tract

❖ **Preferred procedure in:**

1- pregnant women

2- patients allergic to IV contrast.


❖ Not dependent on the composition of stones and detects **uric-acid stones** as well as **calcium stones.**


❖ **Stones** are seen as **highly echogenic foci** and often **produce distal acoustic shadowing** .

❖ Detects hydronephrosis

❖ Generally good sensitivity .

❖ Ureteral stones are difficult because of overlying gas





U/S


Longitudinal section shows stone in renal pelvis.



U/S

Transverse section shows dilated renal pelvis (hydronephrosis)

Cortical thickness decrease in renal failure



U/S

Longitudinal section shows dilated renal pelvis and dilated ureter filled with fluid (hydronephrosis and hydroureter)




U/S


Longitudinal section shows hypoechoic cystic lesion in the renal cortex (renal cyst)



U/S


Mass in the renal cortex, well defined mass with heterogeneous density (hypo and hyperechoic), tumor





U/S

Shows full bladder with stone.





Prostate:

The size of the prostate measured on ultrasound by taking ant-post diameter and transverse diameter and upper-lower diameter.

CT of the urinary tract

Computed tomography of kidneys, ureters and bladder (CT KUB) is a quick non-invasive technique for diagnosis of Urolithiasis .

CT is excellent modality for assessment of:

- Renal masses.
- Obstruction.
- Retroperitoneal disease.
- Staging of renal and bladder neoplasms.
- Tumor invasion into the renal vein or IVC
- Evaluation after trauma or surgery.

Urolithiasis refers to the presence of calculi anywhere along the course of the urinary tract.

Urinary tract stones and CT

What is the most sensitive radiological test for urinary tract stone ?

CT , performed without contrast, is highly sensitive for detecting urinary tract stone


.

Are any urinary tract stones radiolucent on CT ?

No , virtually all urinary tract stones , regardless of their composition are visible on CT .

#stones seen on CT as **hyperdense** wither it is radio-opaque or radio-lucent.

#CT is the best modalities to detect stones.




CT

Stone in left renal pelvis.

Stone on CT:

All stones on CT are radio-opaque.



CT

Left ureteric stone with calcification on aortic wall.

Sensitivity, Specificity Of Procedures.

PROCEDURE	SENSITIVITY	SPECIFICITY
KUB	45-70 %	77 %
CT	95-97 %	96 %
ULTRASOUND	32-70%	70-97%

الطب والجراحة لجنة

MRI

- When a renal mass or abscess is suspected but intravenous contrast cannot be administered, because of either contrast allergy or abnormal renal function, in this case MRI can be performed.

- **Gadolinium**, the contrast agent for MRI, Can be safely administered in such circumstances.
- *#contrast contraindicated when there is renal failure or allergy to contrast.*
- *#gadolinium don't produce any kind of allergy except in advanced case of renal failure it produce systemic fibrosis ??????.*

Intravenous urography (IVU)
Intravenous pyelography (IVP)

Is a radiological procedure used to visualized abnormalities of the urinary system, including the kidneys, ureters, and bladder by using intravenous contrast.

Indication:

- Haematuria
- Renal colic or calculi
- Suspected stone in the ureters
- Renal trauma

#IVU *for kidney, ureter and bladder.*
#IVP *for kidney only.*

IVU /continuation

- After a preliminary control film of the abdomen, 50ml of contrast medium is injected intravenously.
- Contrast is excreted by glomerular filtration.
- Films after 5, 10, and 15 minutes are taken and reveal contrast in the pelvi-calyceal systems, ureters, and in the bladder.
- Post-micturition film is taken to assess bladder residual volume.
- Renal obstruction may require a delayed films.

#before doing IVU we give the patient castor oil and laxative and should avoid drinking in the last 6 hours.

#before injection of contrast antiallergic drug should be available.

#film taken at 5 min, 10 and 15 and during voiding.



IVU

Normal kidneys, ureters and bladder

Renal calyces:

On hydronephrosis there is lost of cupping of the renal calyces.



IVU

Radio-opaque stone at left ureter with filling defect and dilatation of left ureter proximal to the obstruction with hydronephrosis

Renal calyces:

in hydronephrosis there is lost of cupping of the renal calyces.



IVU

Dilatation of left ureter (more than right ureter) with stone at site of insertion of left ureter to the bladder.

الطب والجراحة لجنة




IVU

Stone on left renal pelvis and stone in lower part of left renal pelvis with dilation of left ureter.

Micturating cystogram

- Is the study of the urinary bladder and urethra with contrast medium.
- The bladder is filled with contrast via a urethral catheter. Films of the bladder are obtained.
- After removal of the catheter, patient is asked to void and films are taken during micturition to assess the bladder neck and urethra, as well as reflux.
- Examination of the urethra in oblique position is necessary, particularly in suspected posterior urethral valves in infants and small children, as they are usually only demonstrated during micturition.

الطب والجراحة لجنة




MC


MC(micturating cystogram) shows reflux of contrast on the left ureter during voiding.

Micturating cystogram

#**Vesicoureteral reflux** have five degrees:


1. to lower ureter.
2. to ureter and kidney without hydronephrosis.
3. mild hydronephrosis.
4. moderate hydronephrosis.
5. severe hydronephrosis.





Urethrogram

- The adult male urethra can be studied by ascending urethrogram.
- Contrast is injected through foley catheter inserted into the meatus, and its balloon inflated with 1 to 2ml of sterile water placed in the navicular fossa.
- Films are taken to the urethra in oblique position during contrast injection.
- The most common indication for urethrogam is **urethral strictures.**



Urethrogram

Normal urethra.

Urethra:

Divided into two parts


Ant urethra that divided into penile urethra and bulbous urethra.

Post urethra that divided to membrane urethra (1.5cm) and prostate urethra (2.5cm).



Urethrogram

Urethral stricture in ant urethra.



Urethrogram

Multiple Urethral strictures in ant urethra.

Renal arteriography

Renal arteriography is a special x-ray of the blood vessels of the kidneys.

Evaluation of the renal arterial circulation may be necessary for:

- Further investigation of equivocal renal masses.
- Arterio-venous malformation
- Renal artery stenosis
- Anatomical detail prior to renal transplantation





Congenital renal anomalies

1- **Unilateral renal agenesis**

2- **Renal hypoplasia**

the kidney is small but perfectly formed (*normal function*)

3- **Duplex kidney**

Is the commonest renal anomaly with a variable degree of duplication ranging from minor changes of duplication in the renal pelvis , to a total duplication (complete) of the renal pelvis and ureters.

#see the radiological pic to know the types of duplex kidney.



IVU

Small right kidney (renal hypoplasia)



IVU

Right unilateral renal agenesis



IVU

Bilateral Incomplete duplication of both kidneys, with right hydroureter and hydronephrosis.



IVU

Complete duplication of left kidney.

Duplex kidney:

Two types:

1. *incomplete duplication where there is 2 renal calyces and 2 ureters fused with each other and have one insertion site in u. bladder.*
2. *Complete duplication where there is 2 renal calyces and 2 ureters with different insertion sites.*

الطب والجراحة لجنة



IVU

Complete duplication with phliboliths on left side.

الدكتور حكى الصوره مش واضحة

Congenital renal anomalies/2_

4- Renal Ectopia

- ✓ Refers to a birth defect in which a kidney is located in an abnormal position usually in the pelvis
- ✓ The ectopic kidney is frequently malrotated and small in size.

5- Crossed fused renal ectopia

- ➢ One kidney is displaced across the midline and fused to the other normal kidney .
- ➢ The ureteric orifice lie in a normal position .



MC

Right and left ectopic kidney, both kidneys in the pelvis.



IVU


Crossed left fused ectopic kidney where the left kidney is ectopic it cross the midline and fuses with right kidney.

الطب والجراحة لجنة

Congenital renal anomalies / 3

6- Horse shoe kidney

- Is a fusion anomaly in which the lower poles of the kidneys fuse across the midline .
- The connective tissue may be functioning (*take the contrast*) or non functioning (*fibrous tissue doesn't take contrast*)
- In horseshoe kidney , there is increased incidence of infection and stone formation.





KUB

Horse shoe kidney where both kidney are fused at the lower poles, with multiple radio-opaque stones on left side, not in ureter.

Horse shoe kidney:

Look at lower lobes of both kidney more than the tissue in between to diagnose HSK.

In HSK the lower lobe tilted to the spine.



IVU

Horse shoe kidney.

Horse shoe kidney:

If the tissue in between takes the contrast then its renal tissue (functioning tissue), if not it's fibrous tissue (non-functioning tissue).

الطب والجراحة لجنة



KUB

Stag horn on right kidney, its calcification of renal pelvis and the renal pelvis in this condition is seen without giving the contrast (KUB).

Stag horn>> مثل قرن الأيل





KUB

Stag horn stones on both kidneys

الطب والجراحة لجنة



KUB

Multiple radio-opaque stones in gall bladder with semi-opaque stone in lower part of left ureter.


Gall bladder stone VS renal stone.

To differentiate between them ask for lat abd x-ray.



IVU

Multiple gall bladder stones with stones on left ureter causes hydronephrosis and hydroureter on the left side.



IVU

Semi-opaque stone on left renal pelvis with filling defect.



KUB

Semi-opaque stone on lower left side of pelvis.



IVU

Hydronephrosis and hydroureter on the right side with stone at site of insertion of right ureter with filling defect.



KUB

??????



IVU

Multiple radio-lucent stones in the bladder with thick irregular bladder wall



IVU

Multiple radio-lucent stones in the bladder with thick irregular bladder wall and bladder diverticulum.



KUB

Heavy calcification of the wall of the bladder with thickening, schistosomiasis

Schistosomiasis produce heavy calcification to bladder then ureter then kidney while TB produce calcification to kidney then ureter then bladder.



KUB


Heavy calcification and thickening to the wall of the bladder occur in schistosomiasis.

CYSTITIS

- Is usually due to bacterial organisms ascending through the urethra, although occasionally it may occur by hematogenous spread.
- The bladder outline is irregular due to mucosal edema.
- On ultrasound the bladder wall is thickened and hypoechoic.
- Emphysematous cystitis usually present in diabetic patients. Gas is seen in the bladder wall and sometimes within the bladder lumen.

Acute pyelonephritis

- Is much more commonly seen in women, especially during pregnancy.
- On ultrasound, the kidneys are enlarged, and a wedge shaped areas of decreased echogenicity may be seen.
- CT shows triangular areas of decreased enhancement.
- The infection may proceeds to abscess formation.



CT

Enlargement of right kidney with multiple hypodense areas in the cortex, those areas not enhanced after injection of contrast due to compression on vessels. Occur in acute pyelonephritis.

In infarction of kidney it produce hypodense areas in the cortex not enhanced, but the kidney will not enlarged as pyelonephritis.

Chronic pyelonephritis

- The kidney is reduced in size with irregular renal outline.
- Narrowing of the renal substance thickness.

The affected calyces are dilated and distorted (deformed calyces).

- Vesico- ureteric reflux is frequently associated with chronic pyelonephritis, especially in children.

#75% of children with chronic pyelonephritis have vesico-uretric reflux.



IVU

Distortion and deformity of right renal calyces with small right kidney. Occur in chronic pyelonephritis.

URINARY OBSTRUCTION

- Obstruction of the renal tract may occur at many sites.
- The most common causes are:

- Urinary tract stones.
- Urinary tract strictures.
- Urinary tract tumors.
- Prostatic hypertrophy or cancer.

Urinary obstruction / 2

Why is it important to recognize renal obstruction ?

- Because over time, obstructed kidneys may lose function permanently.

What is the best initial imaging test for suspected renal obstruction ?

- Ultrasound. It is relatively inexpensive, save, and effective. The cause of obstruction also may be identified.

Urinary obstruction / 3

Hydronephrosis and /or hydroureter is identified on ultrasound, but the cause is unclear, what should be done next ?

CT scan or an intravenous urogram.



IVU

Hydronephrosis on left kidney with filling defect at level of pelvi- uretric junction.

Radio-lucent Stone at pelvi-uretric junction.



IVU

Hydronephrosis on left kidney with filling defect at level of pelvi- uretric junction.

Semi-opaque Stone at pelvi-uretric junction.



KUB

Semi-opaque stone in the left ureter.



IVU

Hydronephrosis and hydroureter on left side due to stone.



KUB

??????? Radio-lucent stone on the left ureter.



IVU

No contrast pass to left ureter due to obstruction.



IVU

On voiding, hydroureter and hydronephrosis seen on left side.

Benign renal lesions

What is the most common renal mass ?

The most common mass is a **simple cyst.**

- They are more common in older patients and are found in approximately 50% of the population over 50 years of age.
- They are usually cortical in position and an incidental finding.

Benign renal lesions /2

- What is the best way to confirm that a renal mass is a simple cyst ?

  Ultrasound.

- The ultrasound appearance of a simple cyst is that of a well-defined round mass with very thin wall, smooth margin and no internal echoes.



U/S

Cortical renal cyst with thin wall and smooth margin and hypoechoic core, this cyst didn't cause distraction to kidney tissue.

Tumor in kidney cause destruction to surrounding kidney tissue.



CT

Renal cortical cyst on right kidney with smooth thin wall and hypodense core. And no destruction to renal tissue.

Polycystic kidneys disease


- Adult polycystic kidney disease is a congenital renal parenchymal disorder.
- Usually both kidneys are involved.
- In some cases, there is associated cysts in the liver and more rarely in the spleen and pancreas.

#PKD in newborn==we don't see cysts because the cysts are small and multiple and contain thin wall.

Polycystic kidney disease / 2


Radiological features on Ultrasound and CT:

- Kidneys are enlarged with lobulated contours.
- The renal parenchyma is replaced by multiple cysts of varying size, causing distortion of the collecting system.
- Spontaneous hemorrhage into some of the cysts may occur.



CT

Multiple hypodense cysts in liver and both kidney with enlargement of both kidney. Occur in Polycystic kidney disease.



CT

Multiple hypodense cysts in liver and both kidney with enlargement of both kidney. Occur in Polycystic kidney disease.

Polycystic kidneys / 3

- Polycystic kidney disease of the new born:

  Discovered in the first few days of life with renal failure and gross enlargement of both kidneys.

- Polycystic kidney disease of childhood:

  Usually present at 3 to 5 years.

Benign renal tumors

The most common benign renal tumors are:

- Angiomyolipoma *(most common)*
- Adenoma
- Oncocytoma

Malignant renal tumors


- Renal cell carcinomas (RCC)*(most common)* or Hypernephroma: account for 85% of renal tumors.
  - Are bilateral in 4% of cases.
  - Von Hippel- Lindau disease is associated with RCC in one third to one half of patients.
  - Patients with polycystic kidney disease and chronic renal failure are also associated with RCC.
- Transitional cell carcinoma:

are relatively rare and represent 7% of all renal tumors.




CT

Tumor in left kidney invade the left renal vein




CT

Bilateral renal cell carcinoma.



CT

Bilateral renal cell carcinoma.




CT

Tumor in right kidney.




IVU

Filling defect in the wall of the bladder, indicates tumor in bladder.



CT on abd trauma

Multiple hypodense areas in the abd, indicate Hemorrhage with laceration of spleen and left kidney.



CT on abd trauma

Multiple hypodense areas in the abd, indicate Hemorrhage with laceration of left kidney.

Chapter 7
Pediatrics

Done By: EL SOLO LOBO

Sources:

- DR. Hana Quidsieh slides 2021/2022; Mutah University
- radiopaedia.org
- radiologyassistant.nl

BEFORE WE START…

The aim of the Pediatric chapter is to :

- Be able to recognise the radiological findings and the differential diagnosis or even top diagnosis of the main pediatric cases.
- Be confident and “ never miss “ the pediatric emergency cases.
- Choose the best investigation for further assessment according to the clinical data in the main pediatric cases.
- Have brief clinical knowledge about selected cases in pediatric radiology

الطب والجراحة
الجنة

BEFORE WE START…

- Color code for this chapter :
  - White + blue : everything that is in the slides and you should read it for the exam.
  - Red **: extremely important information** that can help you answer and distinguish radiological cases
  - Green : extra information was put to help you understand better don't read it for the exam
  - purple : some notes and clarification I wrote (not required for exam)
- Test Yourself Cases = (اسئلة ارشيف)
- كل الصور في الشابتر مطلوبة
  + قسم النوت تحت كل سلايد ممكن يكون في توضيح او ملاحظه

الطب والجراحة لجنة

Table of Content

1. Respiratory Pediatric :
   - CDH (congenial diaphragmatic hernia) 845
   - CCAMA (congenital cystic adenomatoid malformation) 851
   - RDS (respiratory distress syndrom) 856
   - CLE (congenital lobar emphysema) 860
   - Round pneumonia 864
   - Esophagal atresia 867
   - Tracho esophagal fistula
2. Abdominal Pediatric :
   - **GIT :**
     - Hypertrophic pyloric stenosis. 876
     - Necrotizing Enterocolitis (NEC) 883
     - Midgut volvolus. 904
     - Midgut malrotation
     - Intussception 914
     - Hirschprung disease. 919
     - Doudenal atrasia. 923
     - Meconuim ileus 926
     - Meconuim plug syndrome. 929
   - **URINARY :**
     - VUR 932
     - Neurogenic bladder 941
3. Musculoskeletal Pediatric :
   - DDH 956
   - Ricket 967
   - Perthes disease 977
   - Child abuse 997
   - Slipped capital femoral epiphysis. 1010

Section1: Respiratory Pediatric

Congenital diaphragmatic hernias CDH

- There are 3 main types of congenital diaphragmatic hernia:
  - Posterolateral: Bochdalek hernia
  - Anterior: Morgagni hernia
  - Hiatus hernia
- **Left-sided Bochdalek** hernias are most common, accounting for approximately **85%** of cases. Left-sided hernias, lacking the 'protective' effect of the liver may have herniation of both the small and large bowel and intraabdominal solid organs into the thoracic cavity.
- When right-sided, only the liver and a portion of the large bowel tend to herniate. Bilateral hernias are uncommon and are usually fatal due to **pulmonary hypoplasia.**

الطب والجراحة لجنة

Congenital diaphragmatic hernias…

- Most congenital diaphragmatic hernias are detected either **soon after birth or on antenatal ultrasound**.
- Mortality is predominantly due to the development of **pulmonary hypoplasia**, which is thought to be due to mass effect on the developing lung.
- Such neonates are **hypoxic and have persistent fetal circulation due to pulmonary hypoplasia and pulmonary hypertension.**

الطب والجراحة لجنة

Congenital diaphragmatic hernias…


- A Bochdalek hernia is a form of congenital diaphragmatic hernia. They occur posteriorly and are due to **a defect in the posterior attachment of the diaphragm** when there is **a failure of pleuroperitoneal membrane closure in utero.** Retroperitoneal structures may prolapse through the defect, e.g. **retroperitoneal fat or left kidney.**
- Clinical presentation
- Large hernias typically present in infancy, are usually are said to be most frequently left-sided, presumably owing to the protective effects of the liver, Complications are usually due to pulmonary hypoplasia

الطب والجراحة لجنة


Congenital diaphragmatic hernias…

- On conventional radiographs, the hernia may appear as **a lung base soft-tissue opacity lesion** seen posteriorly on lateral images.
- Differential diagnosis
- Bochdalek herniae may mimic diaphragmatic rupture from trauma, but you would expect to find other supportive signs of chest and/or abdominal trauma

Multiple cystic structures in the Lt lung cavity with shift of mediastinal structure to the Rt , small bowel gas is not seen in the abdominal cavity, the diaphragm is not destinct





After giving contrast media orally the contrast filed the bowel loops in the left lung cavity.



الطب والجراحة
لجنة

Air fluid level seen in the retrocardiac area that appears in the posterior mediastinum in the lateral xray





Congenital pulmonary airway malformations (CPAM) Congenital cystic adenomatoid malformations (CCAM).

- Congenital cystic adenomatoid malformation (CCAM) is a benign lung lesion that appears before birth as a cyst or mass in the chest. It is made up of abnormal lung tissue that does not function properly, but continues to grow. CCAM is also frequently referred to as a congenital pulmonary airway malformation (CPAM).
- they are multicystic masses of segmental lung tissue with abnormal bronchial proliferation. CPAMs are considered part of the spectrum of bronchopulmonary foregut malformations.
- **Clinical presentation :** the diagnosis is usually either made on antenatal ultrasound, or in the neonatal period on **the investigation of progressive respiratory distress** If large, they may cause pulmonary hypoplasia, with resultant poor prognosis.
- In cases where the abnormality is small, the diagnosis may not be made for many years or even until adulthood. When it does become apparent, it is usually as **a result of recurrent chest infection** .

الطب والجراحة لجنة

Congenital pulmonary airway malformations (CPAM) Congenital cystic adenomatoid malformations (CCAM)…

- Five subtypes are currently classified, mainly according to cyst size:
- type I
  - most common: 70% of cases
  - large cysts
  - one or more dominant cysts: 2-10 cm in size
  - may be surrounded by smaller cysts
- type II
  - cysts are <2 cm in diameter
  - associated with other abnormalities
  - renal agenesis or dysgenesis
  - pulmonary sequestration
  - congenital cardiac anomalies

الطب والجراحة لجنة

Congenital pulmonary airway malformations (CPAM) Congenital cystic adenomatoid malformations (CCAM)…

- **type III**
  - ~10% of cases
  - microcysts: <5 mm in diameter
  - typically involves an entire lobe
  - has a poorer prognosis
- **type IV**
  - unlined cyst
  - typically affects a single lobe
  - indistinguishable from type I on imaging


Congenital pulmonary airway malformations (CPAM)
Congenital cystic adenomatoid malformations (CCAM)…

- Chest radiographs in type I and II CPAMs may demonstrate a **multicystic** **(air-filled)** **lesion**. Large lesions may cause a mass effect with resultant **mediastinal shift**, depression, and even inversion of the diaphragm. In the early neonatal period, the cysts may be completely or partially fluid-filled, in which case the lesion may appear solid or with air-fluid levels. Lesions may change in size on interval imaging . Type III lesions appear solid.


الطب والجراحة لجنة

CASE : Newborn with progressive fetal distress.

Multicystic structure in the Rt lung cavity with shift of mediastinal structure to the left , almost clear visible portion of diaphragm.



After giving contrast via feeding tube the cystic structures in the lung are not related to the bowel



Respiratory distress syndrome (RDS)

- **Respiratory distress syndrome** (**RDS**) is a relatively common condition resulting from insufficient production of surfactant that occurs **in preterm neonates.**
- On imaging, the condition generally presents as **bilateral and relatively symmetric diffuse ground glass lungs** with low volumes and a bell-shaped thorax.
- **Terminology :** RDS is also known as **hyaline membrane disease** (not favored as reflects non-specific histological findings), **neonatal respiratory distress syndrome**, **lung disease of prematurity** (both non-specific terms), or as some authors prefer **surfactant-deficiency disorder** .

الطب والجراحة لجنة


Respiratory distress syndrome (RDS)

- **Clinical presentation :** respiratory distress presents in **the first few hours of life in a premature baby**. Signs include tachypnea, expiratory grunting, and nasal flaring. The infant may or may not be cyanosed. Substernal and intercostal retractions may be evident.
- **Plain radiograph :** typically gives diffuse ground glass lungs with low volumes and a bell-shaped thorax , often bilateral and symmetrical ,air bronchograms may be evident
- **Hyperinflation (in a non-ventilated patient) excludes the diagnosis ,may show hyperinflation if the patient is intubated.**
- RDS can be safely excluded if the neonate has a normal chest radiograph **at six hours after birth.**
- If treated with surfactant therapy there may be asymmetric improvement.

الطب والجراحة لجنة

CASE: 35-week gestation neonate with No complications at birth. Oxygen requirement and increasing respiratory distress requiring mechanical ventilation


- **Bilateral diffuse bilateraly smmetrical ground glass infilterate with air bronchogram no hyperinflation**



Another CASE …

- Preterm neonate with low volume lungs the x-ray shows **complete opacification of both lung** so we think of RDS

العمر مهم ركز عليه



Congenital Lobar Overinflation
(Congenital Lobar Emphysema)

- Congenital lobar overinflation **(CLO),** previously called congenital lobar emphysema, is a congenital lung abnormality that results in progressive overinflation of one or more lobes of a neonate's lung.
- On imaging, it classically presents on chest radiographs as a **hyperlucent lung segment with overinflation and contralateral mediastinal shift**
- Clinical presentation : Patients typically present with **respiratory distress**, most commonly in the neonatal period, and usually within the first six months of life

الطب والجراحة لجنة

Radiographic Features

- Interestingly, there is a pronounced predilection for certain lobes:
  - left upper lobe: most common, 40-45%
  - right middle lobe: 30%
  - right upper lobe: 20%
  - involving more than a single lobe: 5%
  - much rarer in the lower lobes
- Therefore, although the left upper lobe is most commonly affected, **the right hemithorax is more commonly affected than the left**

الطب والجراحة لجنة

Radiographic Features…

❖Plain radiograph:

❖Immediate postpartum period: (مش مطلوب لكن موجود بالسلايد)
The affected lobe tends to appear opaque and homogeneous because of fetal lung fluid or it may show a diffuse reticular pattern that represents distended lymphatic channels filled with fetal lung fluid.

❖Later findings :

- Appears as an area of hyperlucency in the lung with oligaemia (i.e. paucity of vessels)


نقص كمية الدم والاوعية الدمويه الي ظاهرة اقل على الجهة الي فيها hyperlucency

- Mass effect with **mediastinal shift** and hemidiaphragmatic depression

الطب والجراحة لجنة

CASE :22 Day newborn with Respiratory distress.

- **Hyperlucent right middle lobe** with compressive atelectasis of the right upper and lower lobes.There is contralateral **mediastinal shift**.
- Findings suggest right middle lobe **congenital lobar emphysema**.



Round Pneumonia

- It is a type of pneumonia usually only seen in pediatric patients. They are **well defined, rounded opacities that represent regions of infected consolidation.**
- Epidemiology : the mean age of patients with **round pneumonia is 5 years** and 90% of patients who present with round pneumonia **are younger than twelve** Round pneumonia **is uncommon after the age of eight** because collateral airways tend to be well developed by this age
- Clinical presentation : Patients present with symptoms of **chest infection** including fever, sweats, and cough. History of infective symptoms is really helpful when excluding other imaging differentials.
- The infective agent in round pneumonia is **bacterial**. There is no specific bacterium that causes round pneumonia, but since **Streptococcus pneumoniae** is the most common cause of chest infection, it is little surprise that it is the leading cause of round pneumonia

الطب والجراحة لجنة


Radiographic Features

- **Plain radiograph :** Round pneumonias are round-ish and while they are **well-circumscribed parenchymal opacities**, they tend to have **irregular margins.** They most commonly occur in superior segments of lower lobes and in the majority of cases (98%), they are **solitary**.
- **Air-bronchograms** are often present, and helpful in clinching the diagnosis


الطب والجراحة لجنة

CASE : 4 years old girl with fever cough

Rounded opacity in the Rt middle zone contains airbronchogram




follow up after 3 months postreatment



Esophageal Atresia

- Esophagus atresia is an anomaly which arises in **the fourth week** of the embryogenesis, at a stadium in which the trachea and esophagus should separate from each other.
  In case of failure of complete separation esopaghus atresia can occur.
- **Clinically the neonate cannot swallow saliva, may blow bubbles and will aspirate on feeding.**
- **When a feeding tube is inserted it cannot be passed distally.**
- A radiograph with **a curled up feeding tube** will confirm the diagnosis. Contrast swallow studies **should not be** performed as they can cause severe aspiration.

الطب والجراحة لجنة


Distal tracheo-esophageal fistula
No fistula
Proximal tracheo-esophageal fistula
Proximal and distal fistula

Esophageal Atresia…

- **80% of cases a distal tracheo-esophageal fistula is present.**
- Less common is:
  - no fistula - no air in stomach
  - proximal fistula - no air in stomach
  - two fistulas - air in stomach

❖ Cases without a distal fistula can be suspected antenatally when there is a polyhydramnios and an empty stomach.

❖ Always screen the radiograph for other anomalies as esophagus atresia can be part of the **VACTERL association** (vertebral anomalies, anal atresia, cardiovascular malformations, tracheoesophageal fistula, renal and limb anomalies).


❖ **Feeding tube cannot be passed and lies in the proximal esophagus and Gasless abdomen This indicates that there is no distal trachea-esophageal fistula**

الطب والجراحة لجنة

CASE : Newborn with direct regurgitation of milk post feed


- Feeding tube cannot be passed and lies in a dilated proximal esophagus
- air in the abdomen.

Diagnosis: esophageal atresia with a distal tracheo-esophageal fistula



Another Case…


- No passage of NG tube to stomach with absent stomach gas
- Esophageal atresia with no distal trachea-esophageal fistula



Test yourself CASE 1

- **Preterm neonate without complications at birth , increasing O2 requirement in the first hours :**

1. Congenital diaphragmatic hernia
2. **Respiratory distress syndrome**
3. Normal lungs of preterm
4. Congenital cystic adenomatoid malformations




Test yourself CASE 2

- 32 premature baby ,findings ,diagnosis
- Chest x ray shows


bilateral diffuse ,bilateral symmetrical ground glass infiltrate with air bronchogram ,no hyperinflation

Diagnosis : RDS



Test yourself CASE 3

- **15 years old with fever and chest pain ,findings and diagnosis :**
- Right opacity in the right middle zone contains air bronchogram
- **Diagnosis : Round pneumonia**




Section2: Abdominal Pediatric

GIT : Hypertrophic Pyloric Stenosis (HPS)

- **Infants 2 -12 weeks most common** and more common in males
- Present with **NONbilious projectile vomiting**
  - If bilious (green) vomiting, obstruction is past ampulla of Vater (beyond pylorus)
- Can lead to **dehydration, hypochloremic metabolic alkalosis**
- Unknown cause ,thickening of muscle, unable to relax
- Lumen obstructed by thick redundant mucosa
- Other differential includes intestinal atresia, gastro esophageal reflux and gastroenteritis
- **Ultrasound of the pylorus should be the next imaging study.**
  - Some center request x ray as they have no experienced sonographic hands

الطب والجراحة لجنة

Pyloric stenosis



Normal values *


Length: <15mm
Single muscle thickness: <3mm
Pyloric width: <7mm

* values vary somewhat from publication to publication

F Gaillard 2010

Radiopaedia.org CC-NC-SA-BY

❑ Normal ultrasound of pylorus



بنشوف القيم وبنقارن مع القيم الطبيعية

NORMAL STUDY ON SEPARATE PATIENT FOR COMPARISON


الطب والجراحة لجنة

CASE : One month old baby with projectile vomiting and poor weight gain



Hypertrophied hypoechoic pylorus because the values are higher than normal

نفس الحالة



Hypertrophic pyloric stenosis (HPS)

Radiographs may show "caterpillar" sign (gastric distention with large contractions or peristaltic waves as stomach contracts against obstructed pylorus)



- **This sign indicates distal obstruction and the stomach is trying to overcome it it's not specific for Hypertrophic Pyloric Stenosis**

Hypertrophic pyloric stenosis (HPS)

Upper GI may show "string sign" of contrast obstruction at pylorus



- Treatment is surgical laparoscopic pyloromyotomy

الطب والجراحة لجنة

Necrotizing Enterocolitis (NEC)

- Necrotizing enterocolitis is a severe bowel inflammation.
- The etiology is not entirely clear and seems to be a combination of immature bowel mucosa, infection and ischemia.
- Initially radiographs are nonspecific and may only show bowel **dilatation.**
- Absence of a changing bowel pattern over time is worrisome.
- **Pneumatosis intestinalis** and portal venous air (**pneumoportogram**) can both be seen on radiographs and with ultrasound.
- **The most feared complication is perforation**.
- NEC occurs most often, but not exclusively, in prematures.
  Neonates with severe stress, for example with cardiac disease, are also at risk.
  Clinically, retentions and bloody stools can be a key to the diagnosis.

الطب والجراحة لجنة

Necrotizing enterocolitis (NEC)

- Most commonly occurs in premature infants <750g
  - Incidence inversely related to patient age
- Premature neonates have decreased motility, altered intestinal microbiome
  - Leads to stasis, inflammation, altered mucosal integrity
  - Bacteria invade mucosa into bowel wall
  - Gas forms in bowel wall
- May have distention, bloody stools, thrombocytopenia, acidosis, perforation

الطب والجراحة
الجنة

Necrotizing enterocolitis (NEC)

- Prevention
  - Breast milk?
- Treatment
  - Medical (no perforation): antibiotics, NPO with IV nutrition
  - Bowel resection if perforation (implies necrotic bowel)
- Complications
  - Strictures long term (more surgery)
  - Short bowel syndrome if severe

الطب والجراحة لجنة

Necrotizing Enterocolitis (NEC)

- **Different stages of necrotizing enterocolitis :**
- **STAGE 1 : Bowel dilatation** with/without bowel wall thickening
- **STAGE 2 : Pneumatosis intestinalis** with/without **portal venous air**
- **STAGE 3 : Perforation with pneumoperitoneum.**


Radiographic Features

- Supine abdominal x-rays are the mainstay of diagnosis. If perforation is suspected clinically, or there is a concern on supine films, an additional cross-table lateral or left-lateral decubitus view provides increased sensitivity
- Radiological Findings :
  - Dilated bowel loops.
  - Pneumatosis intestinalis. (هواء داخل جدار الامعاء)
  - Portal pneumatosis.
  - Complicated NEC:
    - **Perforation (acute) free gas** : (staging مهم تميز متى بصير عشان ال)
      - ✓Cresent shape
      - ✓Falciform ligament sign
      - ✓Riglers sign
    - Stricture (chronic)

الطب والجراحة لجنة


Necrotizing enterocolitis (NEC)

Pneumatosis intestinalis on radiographs




Necrotizing enterocolitis (NEC)

May see portal venous gas (air tracks along bowel wall to mesentery to SMV to portal vein



Necrotizing enterocolitis (NEC)

May see pneumoperitoneum (falciform ligament [black arrows] outlined by air if patient is supine)



Perforation free gas in peritoneum NEC stage 3


الطب والجراحة لجنة

Necrotizing enterocolitis (NEC)

Another sign of pneumoperitoneum on supine radiograph

Rigler's sign:

Both sides of bowel wall visible (normally you can only see air inside bowel)




Perforation free gas in peritoneum NEC stage 3

الطب والجراحة لجنة

Necrotizing enterocolitis (NEC)

Cross table lateral (left) or left lateral decubitus (right image) will confirm free air




Cresent sign ,Perforation free gas in peritoneum
NEC stage 3


الطب والجراحة لجنة


Necrotizing enterocolitis (NEC)

Contrast enema shows contrast fails to pass proximal to stricture





CASE : **Premature** girl baby was admitted to the NICU for supportive care. After 2 days she was noted to have increasing abdominal distention. Different abdominal radiographs were ordered.





SAME CASE…





THE CASE : Radiological Findings

- Mildly **dilated loops of bowel** with **pneumatosis intestinalis** and portal venous gas. **No free intraperitoneal air is identified.**

- X-ray (Fontal (tight windowing)): Tight windowing allows for easy visualisation of pneumatosis and portal venous gas

THE CASE :UPDATE

- The patient was managed conservatively and is eventually transferred out the ICU. While on the general pediatrics floor, the patient becomes distended once again and is inconsolable. Repeat abdominal X-ray shows multiple distended loops of large and small bowel.

contrast does not flow past the sigmoid colon. this finding is highly concerning for colonic stricture. a complication of nec





CASE 2 : the images shows a typical case of NEC with **pneumatosis intestinalis.**
on the horizontal beam image **there is no sign of free air. (STAGE 2)**






Here images of a neonate who developed NEC. At this early stage the radiograph only shows non-specific bowel dilatation ,At this stage you cannot make the diagnosis.







Here another typical case of NEC ,Notice the air in the portal vein (arrow) and peripheral portal branches.This is seen on the X-Ray and on ultrasound.



Pneumoperitoneum in severe NEC ,Air can be seen on both sides of the bowel wall.This is called the **Rigler sign**. **STAGE 3**



Here a long stricture in the colon transversum in a child who had a NEC.




Midgut Volvolus & Midgut Malrotation

- Malrotation
  - Normal rotation of midgut thru umbilicus and back into abdomen NOT completed in utero
  - Midgut is the bowel supplied by SMA (distal descending duodenum through distal transverse colon)
- With midgut volvulus, patients present with BILIOUS (GREEN) vomiting because obstruction is past the ampulla of Vater


❑**Normally ,The DJ flexure should be located to the left of midline at the level of the duodenal cap.**

الطب والجراحة لجنة


Malrotation with midgut volvulus



NORMAL



MALROTATION (no volvulus yet)




MALROTATION WITH MIDGUT VOLVULUS

الطب والجراحة لجنة


Malrotation with midgut volvulus

- Left image
  Normal situation with the duodenum retroperitoneal.
  Treitz ligament is on the left side of the spine.
  The small intestine is predominantly on the left.
  The cecum is in the right lower quadrant
  There is a long mesentery.



- Middle image
  Displacement of Treitz inferiorly and rightward.
  The small intestine is found predominantly on the right.
  Fibrous bands course over the vertical portion of the duodenum causing obstruction.

- Right image
  Volvulus due to short mesentery. Ischemic bowel.



Malrotation with midgut volvulus

- 60-80% of patients with volvulus present in first month of life
- Infant is initially well, then sudden onset of bilious vomiting
- Volvulus causes vascular compromise
  - Failure to quickly recognize and diagnose can lead to bowel ischemia, shock, and death
- Any child with true bilious vomiting must have an emergent upper GI evaluation to exclude volvulus


الطب والجراحة
لجنة

Malrotation with midgut volvulus


- Treatment
  - Correct electrolyte abnormalities
  - Emergent surgery with detorsion of bowel and Ladd procedure (tack down the bowel so it cannot twist on itself)

الطب والجراحة
لجنة

Malrotation with midgut volvulus



MALROTATION WITH
MIDGUT VOLVULUS




Normal upper GI, normal C-loop of duodenum. Duodenojejunal junction is left of pedicle, at height of pylorus

- The DJ flexure should usually be located to the left of midline at the level of the duodenal cap. Here it is not, indicating that there is malrotation.
- The lumen is narrowed and passes directly inferiorly in a corkscrew pattern. Findings are of midgut volvulus secondary to intestinal malrotation.

- Contrast demonstrates a misplaced DJ flexure and a corkscrew appearance of the jejunum. This confirms both malrotation and volvulus



corkscrew



Here a neonate with a malrotation. The abdominal radiograph is non-specific (image on the left)
The upper GI study clearly demonstrates that the small bowel projects to the right of the spine



MALROTATION NO VOLVOLUS




Malrotation with midgut volvulus

- The malrotation will become symptomatic only when a volvulus occurs due to the short mesentery or when the Ladd's band obstruct the duodenum
- Both presentations are most common in the neonatal period.
- However sometimes it can also present later in life,
- Acute volvulus is a life-threatening presentation and requires prompt surgical intervention.

الطب والجراحة لجنة

The upper GI-study shows a malrotation complicated by a volvulus ,This results in the typical corkscrew or reversed 3 sign. An overfilled stomach could hide the corkscrew on the AP projection so the stomach should first be aspirated by use of a nasogastric tube and the volume of injected contrast should be small.




ع

Intussusception


- It is **invagination of a segment of bowel (the intussusceptum) into another, usually more distal, segment of bowel (the intussuscipiens)**
- **If not reduced, edema, ischemia and bowel obstruction (usually partial) ensue with necrosis of bowel**
- Age of presentation is usually **3 months to 24 months.**
- Pathologic intussusception is associated with a lead point such as a tumor, inspissated feces (cystic fibrosis) or lymphoma, often in older child greater than age 2.
- Symptoms include **crampy abdominal pain, bloody stools, and vomiting**.
- Treatment is **fluoroscopically guided reduction with air or fluid enema** or surgery if unreducible.

الطب والجراحة لجنة

CASE : 16 months old girl with abdominal pain and distension, vomiting and red jelly stool.



Target sign in transverse image



THE CASE

Claw sign with no passage of contrast to proximal transverse colon



Air reduction as treatment




THE CASE…


- X-rays demonstrate a large soft tissue opacity in the right upper quadrant which appears to bulge into what is presumably the transverse colon.
- Ultrasound confirms an intussusception with a nice example of **a target sign**. The intussusception was reduced with an air enema.

الطب والجراحة
لجنة

TARGET SIGN



PSEUDO KIDNEY SIGN



Hirschprung disease.

- In Hirschsprung disease ganglion cells are absent in the distal part of the colon. Because the intestinal ganglion cells migrate in a craniocaudal direction, the area of aganglionosis always involves the rectum.
- More extensive disease extends orally in a contiguous fashion. The involved bowel has a small diameter and the bowel proximal to the affected segment is dilated.
- Different age presentation varies **between failure to pass meconium in neonate period to constipation and bowel motion disturbances.**
- In Hirschsprung disease the ratio between the denervated and the non-affected bowel is <1.
- It is important to describe the length of the affected segment.
- Most cases are short-segment and total aganglionosis is rare.
- In case of total aganglionosis the diagnosis is difficult, because the entire colon has a small caliber and resembles a microcolon.


الطب والجراحة لجنة

Hirschprung disease.

- Start the enema in the lateral position to evaluate the rectum. Save cine images from the first contrast injection, as with progressive filling signs can become obscured by too much bowel distention.
- Normally the rectum should be wider than the sigmoid. The image shows an abnormal recto-sigmoid index <1.
- The definitive diagnosis of Hirschsprung disease is confirmed with biopsy.
- About 90% of Hirschsprung disease is diagnosed in the neonatal age, but some cases are discovered later in life.


الطب والجراحة لجنة

- Supine abdominal film demonstrates numerous dilated loops of bowel, most likely down to the descending colon. Some gas is seen in the pelvis, most likely in the rectosigmoid.
- Further investigations is required.



Narrow rectum with a short narrow transition zone between in sigmoid with dilated descending colon






Duodenal Atresia

- In duodenal atresia the duodenum fails to canalize properly late in the first trimester and a web or several webs occur.
  In cases of atresia the web is complete.
  Most often the atresia occurs distal to Vater's ampulla.
  The obstruction causes the duodenum to expand and this creates **the double bubble sign (dilated stomach and duodenum).**
  When **a polyhydramnios and a double bubble are present antenatally, the diagnosis can be suspected before birth..**
- In extreme prematures the diagnosis can be more challenging when the duodenum has not had enough time to dilate.
- Note that the nasogastral tube may decompress the stomach and duodenum limiting the interpretation of the film.
  One may inject some air through the tube prior to the film.
- About 30% of the patients with duodenal atresia have **Down syndrome and there is an association with VACTERL malformations, malrotation and biliary tree abnormalities.**

الطب والجراحة لجنة

Radiological Findings


- Double bubble without distal bowel gas.
  This confirms the diagnosis of duodenal atresia and no further imaging is needed.



Double bubble without distal bowel gas.

- Feeding tube cannot be passed and lies in a dilated proximal esophagus
- air in the abdomen.

Diagnosis: esophagus atresia with a distal tracheo-esophageal fistula



Meconium ileus

- Meconium ileus occurs nearly **exclusively in patients with cystic fibrosis**.
- In 10% of patients it is the first presentation of the disease.
  Due to the exocrine dysfunction of the pancreas and abnormal intestinal
- secretions, the meconium is abnormally thick and becomes impacted in the ileum.
- Sometimes radiographs demonstrate **typical 'soap bubbles', which represent captured air between meconium pellets.**
  The tenacious meconium prevents the occurrence **of air fluid levels** on the decubitus image.
  Bowel loops are usually of different caliber and not all loops are dilated. **Colon enema shows a micro colon with stacked meconium pellets in the ileum.**


الطب والجراحة لجنة

Meconium ileus

- Once you have made the diagnosis of a meconium ileus, you can opt to set in moderately **hyperosmolar contrast** for therapeutic purposes as this can help to dissolve the meconium and act as an effective enema.
  Since the hyperosmolar contrast will create a fluid shift and thereby may cause dehydration, it is of importance to clearly communicate with the clinician to administer extra fluids and secure continuous careful surveillance.

CASE : Neonate with abdominal distension no passage of meconium

- **Small colon with multiple meconium pellets in the distal small bowel (arrows).**





Meconium plug syndrome

- is also known as small left colon syndrome.
- Meconium plugging in the left colon occurs when the colon is functionally immature with little motility.
- There is an association with **maternal diabetes** and **drug use in pregnancy.**
- The condition is **temporarily** and when the meconium plugs resolve, the colon distends normally and functions normally.
- The neonate is **otherwise healthy** and there is no association with cystic fibrosis.

الطب والجراحة لجنة


Meconium plug syndrome

- There is no air in the rectum on the radiograph. Colon enema shows a normal rectal diameter which excludes Hirschsprung disease.
- A microcolon is absent.
- Meconium is found throughout the colon, but **most typically found in the left colon**, which may also have a smaller diameter.
- Just as with a meconium ileus, you may now opt to give a hyperosmolar contrast enema to help resolve the meconium

الطب والجراحة لجنة

Case 2 : neonate with no passage of meconuim

- **Left microcolon but normal rectum, but plain film shows dilated loops**






Vesicoureteral reflux (VUR)

- ❖ Abnormal ureterovesicular junction with too large or abnormal position of orifice.
- ❖ Allows urine (and bacteria) to **travel backwards toward kidney**
- ❖ Increased risk of **upper UTI and pyelonephritis** with worse severity of VUR
- ❖ We do kidney ultrasound for Patients with recurrent UTI and we look for a **dilated pelvicalcyceal system (hydronephrosis)**

الطب والجراحة لجنة

Normal ultrasound of kidney

- We see two layers the innermost is hyperechoic and the outermost is hypoechoic



NOT IMPORTANT

Ultrasound Terminology



Anechoic
Echolucent


Hypoechoic

Hyperechoic
Echogenic

Echogenecity is relative

CASE : 5 Year Old Girl With Recurrent Uti


- **Dilated pelvicalyseal system (hydronephrosis)**



VUR


Febrile UTI first evaluated with renal and bladder ultrasound.

Notice urinary tract dilatation, an abnormal finding




VUR


- If renal and bladder ultrasound is abnormal, VUR is diagnosed with **voiding cystourethrogram (VCUG).**
- Contrast instilled via bladder catheter
  - **With VUR, contrast seen in collecting system**
  - **Normally (no VUR), contrast stays in bladder and not seen in ureters**



Grading of VUR

Grade	Description
I	Into a nondilated ureter
II	Into the pelvis and calyces without dilatation
III	Mild to moderate dilatation of the ureter, renal pelvis, and calyces with minimal blunting of the fornices
IV	Moderate ureteral tortuosity and dilatation of the pelvis and calyces
V	Gross dilatation of the ureter, pelvis, and calyces; loss of papillary impressions; and ureteral tortuosity





- Pre-void contrast filled bladder demonstrated bilateral vesico-ureteral reflux (VUR) with mildly tortuous and moderately dilated ureters, with contrast reaching blunted dilated calyces (low pressure VUR)
- After voiding, the collecting systems fill even further, compatible with a component of high pressure VUR.

Findings are keeping with bilateral type 4 vesico-ureteral reflux.

الطب والجراحة لجنة

VUR

- Treatment depends on severity
  - Antibiotic prophylaxis
  - Deflux procedure (surgical “mounds” made at ureterovesicular junction to close a big orifice)
  - Ureteral reimplantation


الطب والجراحة
الجنة

Neurogenic bladder

- ❖ is a term applied to a dysfunctional urinary bladder that results from an injury to the central or peripheral nerves that control and regulate urination. Injury to the brain, brainstem, spinal cord or peripheral nerves from various causes including infection, trauma, malignancy or vascular insult can result in a dysfunctional bladder
- ❖Elongated appearance to the bladder, with trabeculation.The appearances are those of a pine or Christmas tree.

الطب والجراحة لجنة


8 year old child with urine control problem





Test yourself CASE 1


- **32 weeks baby presented with abdominal distention , his x-ray shows :**
  1. **Pneumatosis intestinalis**
  2. Football sign
  3. Free gas
  4. Regler’s sign



الطب والجراحة لجنة


Test yourself CASE 2

- **One month old baby with non-bilious vomiting , the ultrasound demonstrate : (note A: 11mm , B:7 mm)**
  1. Gallbladder stones
  2. **Normal pylorus**
  3. Hypertrophic pylroic stenosis



Test yourself CASE 3

- 6 months old male complaining from chronic constipation:
  1.Name of study?
  Barium enema
  2. Findings and diagnosis?
  There is narrowing in the rectum
  Dilatation in sigmoid colon
  Recto sigmoid index <1



Test yourself CASE 4

- Catrabiller sign
- AXR
- hypertrophic pyloric stenosis




Test yourself CASE 5

- Claw sign
- barium enema
- intussusception




Test yourself CASE 6

- **Target sign**
- **Ultrasound**
- **intussusception**



Test yourself CASE 7

- **Barium meal**
- **inverted 3 sign**
- **Malrotation with volvolus**



Test yourself CASE 8

- What is the condition and the appearing sign ?

  Intussusception , claw sign



Test yourself CASE 9

- **Spot diagnosis:** intussusception
- **Main sign:** claw sign




Test yourself CASE 10

- **Spot diagnosis:** Duodenal atresia
- **Main sign:** double bubble sign




Test yourself CASE 11

- **Spot diagnosis: Hydronephrosis**
- **Main sign: branching hypoechoic**
  Dilated pelvicalyseal system



Test yourself CASE 12

- **Name of study :**
  ultrasound of pyloric region
- **Findings :**
  Increase in the single muscle thickness and length
- **Diagnosis :**
  hypertrophic pyloric stenosis




Section3: Musculoskeletal Pediatric


DDH :Developmental dysplasia of the hip

- In older texts **Congenital Dislocation Of The Hip (CDH)**, denotes aberrant development of the hip joint and results from **an abnormal relationship of the femoral head to the acetabulum**.
- Unlike CDH, developmental dysplasia of the hip is not confined to congenital malformations and includes perturbations in development .
- **There is a clear female predominance**, and it usually occurs from ligamentous **laxity and abnormal position in utero**. Therefore, it is more common with **oligohydramnios pregnancies**.


NORMAL HIP





H. Line (Hilgenreiner's line) join two triradiate cartilage
Perkin : lateral margin of acetabulum and perpendicular over h line




Shenton’s line :
continuous line from lesser trochenter to lower pubic bone (normally)







Perkin line
Acetabular index
Hilgenreiner line
Shenton line


Dislocated hip
Perkin's Line
Perkin's Line
H Line
Shenton's Line

ACETABULAR INDEX (ANGEL)

Newborns	6 months	2 years
27.5° (mean)	23.5°	20°

If it is more than these values think
About DDH





Developmental dysplasia of the hip (DDH)






Developmental dysplasia of the hip (DDH)





OTHER SIGNS

- Lateral displacement of femoral head
- Medial gap
- Capital epiphyseal dysplasia
- Pelvic (Ilium) hypoplasia
- Delayed fusion
- Adducted attitude



الطب والجراحة
لجنة

One year old girl with lower limb instability ,(delay motoric) with positive family history her aunt with DDH.



Rickets

- **Rickets** refers to deficient mineralization of the growth plate in the pediatric population.
- The onset and presentation of rickets depend on the etiology and degree of deficiency. Typically, in severe cases, rickets becomes apparent in the second year of life.
- The presentation is usually with skeletal changes and bone pain
- In the growing skeleton, the deficiency of normal mineralization is most evident at **metaphyseal zones** of provisional calcification where there is an excess of non-mineralized osteoid resulting **in growth plate widening and abnormal configuration of the metaphysis**

Rickets

- As osteomalacia co-occurs with rickets, it is important to remember that even bones that appear mineralized are weak and result in bowing, most commonly seen in the lower limbs once the child is walking.
- Other bone deformities are also noted such as genu valga and vara as well as protrusio acetabuli
- Treatment requires correction of the metabolic imbalance. Only rarely is the orthopedic surgical intervention necessary to correct skeletal deformities

الطب والجراحة لجنة

RADIOLOGICAL FINDINGS

- Metaphysis :
  - Fraying: Indistinct Margins Of The Metaphysis
  - Splaying: Widening Of Metaphysial Ends
  - Cupping: Concavity Of Metaphysis
- Epiphyseal Widening
- Ribs Richet Rosary (Beeded Appearance On Posterior Rib)
- General Bone :
  - Demineralisation
  - Bowing Of Weight Bearing Bones.
  - **Protrusio Acetabuli**
  - Coarse trabeculation.
  - Fracture

الطب والجراحة لجنة

CAUSES


- Vitamin D Deficiency (Most Common Cause) Due To Diet, Insufficient Sun Exposure, And/Or Malabsorption
- Dietary Calcium Deficiency
- Defective Vitamin D Metabolism:
- Hereditary Hypophosphatemic Rickets (Vitamin D Resistant Rickets)
- Tumor-induced (Oncogenic) Rickets
- Fanconi Syndrome

الطب والجراحة لجنة

Fraying ,splaying and cupping of metaphysis




Normal hand xray



One year old baby presented with failure to thrive, bowed legs and a thick wrist.




Rachitic Rosary





Bowing of both femure,tibia and fibula




- Bowing of the femurs, metaphyseal cupping and fraying.
- Coarsening of the trabecular pattern.
- Increase in distance between end of shaft and epiphyseal center.
- poorly ossified epiphyseal centers.



الطب والجراحة
لجنة

Protrusion of hip and spine into the soft pelvis with protrussio acetabuli



Legg calve perthes

- Legg-Calve-Perthes disease is **idiopathic avascular necrosis of the femoral head.**
- It occurs most commonly in children between the **ages of 5-8 who present with knee or hip pain or a limp.**
- **Boys are more commonly affected than girls** (5:1).
- The disease is bilateral in approximately 15% of cases.
- A good prognostic indicator in Legg-Calve-Perthes disease is the age of onset because if it occurs by age six, then restoration of the spherical femoral head is likely and degenerative osteoarthritis can be avoided

الطب والجراحة لجنة

The radiographic changes

- The radiographic changes to the femoral epiphyses depend on the severity of osteonecrosis and the amount of time that there has been an alteration of blood supply:
- early: there may be no appreciable change
- established: reduction in epiphysis size, lucency
- late: fragmentation, destruction
- As changes progress, the width of the femoral neck increases (coxa magna) in order to increase weight-bearing support.

الطب والجراحة لجنة

Early signs

- joint effusion: widening of the medial joint space
- asymmetrical femoral epiphyseal size (smaller on the affected side)
- apparent increased density of the femoral head epiphysis
- blurring of the physeal plate (stage 1)
- radiolucency of the proximal metaphysis

الطب والجراحة لجنة

Late signs

- Eventually, the femoral head begins to fragment (stage 2), with subchondral lucency (crescent sign) and redistribution of weight-bearing stresses leading to thickening of some trabeculae which become more prominent.
- The typical findings of advanced burnt out (stage 4) Perthes disease are:
  - femoral head deformity with widening and flattening (coxa plana)
  - proximal femoral neck deformity: coxa magna
  - "sagging rope sign" (thin sclerotic line running across the femoral neck)

الطب والجراحة لجنة

Legg Calve Perthes

❖**Four phases of Legg Calve Perthes Disease Waldenstrom stages:**

1) Initial stage
2) Fragmentation stage
3) Reossification stage
4) Healed stage

الطب والجراحة الجنة

Initial Stage

- Early radiographic signs:
  - Failure of femoral ossific nucleus to grow
  - **Widening of medial joint space**
  - **"Crescent sign"**
  - **Irregular physeal plate**
  - **Blurry/ radiolucent metaphysis**

Fragmentation Stage

- Bony epiphysis begins **to fragment**
- Areas of increased lucency and density
- Evidence of repair aspects of disease

الطب والجراحة الجنة

Reossification Stage

- Normal bone density returns
- **Alterations in shape of femoral head and neck evident**

Healed Stage

- Left with residual deformity from disease and repair process

Catterall Criteria


Stage I — Histologic and clinical diagnosis without radiographic findings

Stage II — Sclerosis with or without cystic changes with preservation of the contour and surface of femoral head

Stage III — Loss of structural integrity of the femoral head


Stage IV — Loss of structural integrity of the acetabulum in addition

4 year old BOY with painless limping then became painful on LT knee and hip



Mild reduction of epiphyseal height on left side


- NOT CONVENCING
- ASK FOR FROG HIP XRAY
- ALWYS ASK FOR FROG POSITION







Left side: reduction of the femoral head height, wide epiphyseal plate ,subchondral lucency (subchondral fracture)



6 month follow up later….. collapsed LT femoral head with sclerosis

الطب والجراحة لجنة

- Irregularity in contour of the right femoral head and widening of the right joint space (ascompared with the normal findings on the left) with right femoral head epiphyseal height reduction




- The femoral head has become markedly flattened (coxa plana) and enlarged (coxa magna).




- Crescentic subchondral lucency consistent with a subchondral fracture through necrotic bone. **“Crescent sign”**





الطب والجراحة
لجنة





الطب والجراحة
لجنة





SUBCHONDRAL FRACTURE LUCENCY

الطب والجراحة
لجنة



COXA PLANA : Collapsed both femoral head epiphysis

الطب والجراحة لجنة

CHILD ABUSE (non-accidental trauma)

- Child abuse, or non-accidental trauma, is an all-too-common entity. Estimates have suggested that over one million children (most under one year of age) are seriously injured and up to 5000 children killed each year in the United States secondary to physical abuse. When suspicions of potential abuse are raised due to either clinical or radiographic findings, a **skeletal survey** must be obtained.
- A **skeletal survey** is a series of radiographs, performed to cover the entire skeleton or the anatomic regions appropriate for the clinical indications.

الطب والجراحة لجنة

CHILD ABUSE…

- A skeletal survey consists of :
  - two views of the skull
  - lateral thoracic and lumbar spine
  - AP views of both upper and both lower extremities
  - AP views of both hands and both feet
  - AP view of the Pelvis
  - may require CT scan of head
  - may require repeat skeletal survey in two weeks to look for healing injuries not seen on initial survey

الطب والجراحة لجنة

Fractures

- **Fractures highly Specific for Child Abuse :**
  - **Metaphyseal fracture (Bucket handle or Corner fractures)**
  - Ribs (especlally posterior)
  - Acromion
  - Spinous processes
  - Sternum
  - Occipital impression fractures

Fractures…

- ❖Diaphyseal fractures are non-specific as they do occur in both accidental and non-accidental injury. However in these cases **the age of the child** and **the history** become very important.
- ❖**Skull fractures are common child abuse injuries but they are also common in accidental trauma.**
- Patterns of skull fracture that **suggest child abuse** are:
  - Multiple 'eggshell' fractures
  - Occipital impression fractures
  - Fractures crossing sutures

  The infant's skull is very resistent to trauma, so any fracture that is inconsistent with a history should raise the question **of non-accidental injury.**

الطب والجراحة لجنة

Fractures…

- Callus in Diaphyseal fractures generally forms **no earlier than 5 days after a fracture**, but will usually form by 14 days.
  Thus, fractures without visible callus may be up to 14 days old, and fractures which demonstrate a little bit of callus are at least 5 days old. Large amounts of callus indicate at least 2 weeks old.
  These signs can be used to correlate with the history. For instance **a child that fell out of bed the day before cannot have a fracture with callus formation.**
- Metaphyseal fractures **do not** typically heal with callus as there is no periosteal disruption, so dating of metaphyseal fractures is difficult.

الطب والجراحة لجنة

CASE : One year old male presented with downfall from the couch.


- a spiral fracture in the femur (diaphyseal fracture)
- It's suspicious of child abuse because of the age of the child and an inconsistent history given by the parents (the given history can't (normally) cause a spiral fracture).



Corner Fracture

- Corner Fracture a small piece of bone is avulsed due to shearing forces on the fragile growth plate it is seen as the typical corner fracture.
- In a circumstance of abuse, an adult may **grab, twist, or shake a child's limb in frustration**, causing this type of injury. Shaking the child, causing the limbs to move back and forth, can also cause a bucket-handle fracture







الطب والجراحة لجنة

Bucket-Handle Fracture

- Bucket handle fractures These fractures are essentially the same as corner fractures. The avulsed bone fragment is larger and seen 'en face' as a **disc or bucket handle.**



They are identified in the healing stage as a result of associated callus.




Skull Fracture

Patterns of skull fracture that suggest child abuse are:

- Multiple 'eggshell' fractures
- Occipital impression fractures
- Fractures crossing sutures





This 2 month old baby girl was brought into the local ER with scalp swelling
There are multiple skull fracture –eggshell appearance (arrow)


الطب والجراحة
لجنة

Diaphyseal femur fracture with a lot of callus is at least 2 weeks old.



OTHER FINDINGS…

T1wi shows bilateral fluid collections as a result of chronic bilateral subdural hematomas and
New subdural hematomas in the right frontal and posterior
Interhemispheric region
(bilateral subdural hematoma with different ages)




Slipped Capital Femoral Epiphysis


- It should be suspected in any adolescent who **complains of hip or knee pain.**
- **It is twice as common in males** (12-15 years) than females (10-13 years).
- Predisposed risk factors include **obesity, renal osteodystrophy, and endocrine disorders** including hypothyroidism and hypopituitarism.
- Bilateral involvement of the hips can be seen in 20-30% of patients, but it is unusual to present at the same time.


الطب والجراحة لجنة


Slipped Capital Femoral Epiphysis

- **Displacement of the femoral head is posteromedial** and often difficult to see on a standard AP film.
- Findings on AP film can include **asymmetric physeal widening and/or an indistinct metaphyseal border at the level of the physis.**
- Frogleg lateral views are often essential for diagnosis as **minimal slippage can be better appreciated.**
- SCFE is typically treated surgically with **pin fixation (done at current location) to prevent further slippage.**
- Potential complications include **avascular necrosis and chondrolysis.**




Mild
0–1/3
Moderate
1/3–2/3
Severe
2/3–complete
Change in apposition, AP projection
Mild
0–30°
Moderate
30°–60°
Severe
60°–90°
Slip angle, true lateral projection







- CASE : 13 years old boy Lt hip pain
  AP and frogleg views demonstrate widening of the left proximal femoral physis,there is medial and dorsal displacement of the capital femoral epiphysis. **On the frogleg view**, the line drawn tangential to the lateral cortex of the left metaphysis does not bisect any portion of the ossified left epiphysis (compare with a similar tangential line on the contralateral side). Note the **asymmetric appearance of the left acetabulum and the irregularity of the femoral neck adjacent to the physis.**

الطب والجراحة لجنة


SCFE CASES



SCFE CASES



SCFE CASES



Test yourself CASE 1

- Sign and Diagnosis :

1- Widening of growth plate space (alarming sign)
2- Slipped capital femoral epiphysis.



Test yourself CASE 2

- Regardless the Acetabular angle , Is there DDH and Why ?

Answer :


Yes There Is , The Shenton's Line Isn't continues curved and There is a medial Gap .



Test yourself CASE 3

- Is this a DDH or not ,and why ?
  no its normal , and the reason by excluding these signs , also if we draw shintons and helgenrines lines the capitel epiphysis is not in the upper lateral quadrant .
  (the question also gived acetabular index)





Test yourself CASE 4

- What are the radiological signs, the name of the condition, mention one cause ?


  bowing of the femurs, metaphyseal, cupping and fraying, coarsening of the trabecular pattern, increase in distance between end of shaft and epiphyseal center, poorly ossified epiphyseal centers. Its rickets

CAUSES :
vitamin D deficiency (most common cause)



Test yourself CASE 5

- **Spot diagnosis**: SCFE
- **Main sign**: increased growth plate space (alarming feature)



Test yourself CASE 6

- **Spot diagnosis** : Normal adult pelvis
- **Main sign:** non



Test yourself CASE 7

- AP and frogleg views demonstrate widening of the left proximal femoral physis.
- There is medial and dorsal displacement of the capital femoral epiphysis.
- On the frogleg view, the line drawn tangential to the lateral cortex of the left metaphysis does not bisect any portion of the ossified left epiphysis
- (compare with a similar tangential line on the contralateral side). Note the asymmetric appearance of the left acetabulum and the irregularity of the femoral neck adjacent to the physis
- Diagnosis : Slipped Capital Femoral Epiphysis






"That's all Folks!"