

Musculoskeletal System

THE SKULL II

Dr. Aiman Qais Afar

Surgical Anatomist


College of Medicine /University of Mutah

2025-2026

Monday 23 February 2026

Inferior View of the Skull

- ❖ The palatal processes of the **maxillae** and the horizontal plates of the **palatine bones** can be identified.
- ❖ In the midline anteriorly is the **incisive fossa and foramen**.
- ❖ Posterolaterally are the **greater and lesser palatine foramina**
- ❖ The **Choanae (posterior nasal apertures)** are separated from each other by the posterior margin of **the Vomer**




Inferior View of the Skull

❑ The greater wing of the sphenoid is pierced by the **large foramen ovale** and the **small foramen spinosum**.

❑ In the interval between the **greater wing of the sphenoid** and the **petrous part of the temporal bone**, is a groove for the **cartilaginous part of the auditory tube (anterior 2/3 of the tube)**


❑ The opening of the **bony part of the tube** can be identified (**posterior 1/3 of the tube**)



Inferior View of the Skull


✓ The **styloid process** of the **temporal bone** projects downward and forward from its inferior aspect.

✓ The **opening of the carotid canal** can be seen on the inferior surface of the **petrous part of the temporal bone**.





Inferior View of the Skull

✓ The medial end of the **petrous part of the temporal bone** is irregular and, together with the **basilar part of the occipital bone** and the **greater wing of the sphenoid**, forms **the foramen lacerum**.



Inferior View of the Skull


✓ In the interval between **the styloid and mastoid processes**, the stylomastoid foramen can be seen.



Dr. Aiman Al -Maathidy
Monday 23 February 2026


Inferior View of the Skull

✓ Medial to **the styloid process**, the **petrous part of the temporal bone** has a deep notch, which, together with a shallower notch on **the occipital bone**, forms **the jugular foramen**.



Inferior View of the Skull

- ✓ **The pharyngeal tubercle** is a small prominence on the under surface of the basilar part of **the occipital bone** in the midline.
- ✓ **The occipital condyles** they articulate with the superior aspect of the lateral mass of the first cervical vertebra, **the atlas**.
- ✓ Superior to **the occipital condyle** is the **hypoglossal canal** for transmission of the hypoglossal N.




- ✓ Posterior to **the foramen magnum** in the midline is the **external occipital protuberance**.

Base of the skull

The interior of the base of the skull is divided into three cranial fossae:

- **ANTERIOR, MIDDLE, AND POSTERIOR.**
- ❖ The anterior cranial fossa is separated from the middle cranial fossa by **the lesser wing of the sphenoid,**

❖ **and the middle cranial fossa is separated from the posterior cranial fossa by the petrous part of the temporal bone**




Anterior Cranial Fossa

❑ The anterior cranial fossa lodges **the frontal lobes** of the cerebral hemispheres

✓ It is bounded anteriorly by the inner surface of **the frontal bone**, and in the midline is a **crest** for the attachment of **the falx cerebri**.

✓ Its posterior boundary is **the sharp lesser wing of the sphenoid**

✓ The medial end of the lesser wing of the sphenoid forms **the anterior clinoid process** on each side, which gives attachment to **the tentorium cerebelli**



Anterior Cranial Fossa

11


Dr. Aiman Al-Maathidy
Monday 23 February 2026

✓ The floor of the fossa is formed by the orbital plates of the frontal bone laterally and by the cribriform plate of the ethmoid medially

✓ The crista galli is a sharp upward projection of the ethmoid bone in the midline for the attachment of the falx cerebri.

✓ There is slit in the cribriform plate for the passage of the anterior ethmoid nerve into the nasal cavity.

✓ The upper surface of the cribriform plate supports the olfactory bulbs, and the small perforations in the cribriform plate are for the olfactory nerves.




Middle Cranial Fossa

□ The median raised part is formed by **the body of the sphenoid**, and the expanded lateral parts form concavities on either side, which lodge the **temporal lobes of the cerebral hemispheres**.

It is bounded

anteriorly by the lesser wings of the sphenoid
posteriorly by the superior borders of the petrous parts of the temporal bones.

Laterally lie the squamous parts of the temporal bones, the greater wings of the sphenoid, and the parietal bones




□ **The floor** of each lateral part of the middle cranial fossa is formed by **the greater wing of the sphenoid and the squamous and petrous parts of the temporal bone**.

Posterior Cranial Fossa

It is deep and lodges the parts of the cerebellum, pons, and medulla oblongata.

- ❖ **Anteriorly**, the fossa is bounded by the superior border of the petrous part of the temporal bone,
- ❖ **posteriorly** it is bounded by the internal surface of the squamous part of the occipital bone
- ❖ **The floor** of the posterior fossa is formed by the basilar, condylar, and squamous parts of the occipital bone and the mastoid part of the temporal bone.



- ❖ **The roof** of the fossa is formed by a fold of dura, the tentorium cerebelli

Summary of the More Important Openings in the Base of the Skull and the Structures That Pass Through Them

Opening in Skull

Bone of Skull


Structures Transmitted

Anterior Cranial Fossa

Perforations in cribriform plate

Ethmoid

Olfactory nerves



Dr. Aiman Al -Maathidy
Monday 23 February 2026

Summary of the More Important Openings in the Base of the Skull and the Structures That Pass Through Them

Opening in Skull

Bone of Skull

Structures Transmitted

Middle Cranial Fossa

Optic canal



Lesser wing of sphenoid

Optic N, **ophthalmic A**

Superior orbital fissure

Between lesser and greater

Lacrimal, frontal, trochlear nerves




Dr. Aiman Al -Maathidy
Monday 23 February 2026

Summary of the More Important Openings in the Base of the Skull and the Structures That Pass Through Them

Opening in Skull	Bone of Skull	Structures Transmitted
Foramen rotundum	Greater wing of sphenoid	Maxillary division of the trigeminal N.
Foramen ovale	Greater wing of sphenoid	Mandibular division of the trigeminal N., lesser petrosal N.
Foramen spinosum	Greater wing of sphenoid	Middle meningeal artery
Foramen lacerum	Between petrous part of temporal and sphenoid	obliterated

Monday 23 February 2026
Dr. Aiman Al -Maathidy



Summary of the More Important Openings in the Base of the Skull and the Structures That Pass Through Them

Opening in Skull	Bone of Skull	Structures Transmitted
------------------	---------------	------------------------

Posterior Cranial Fossa

Foramen magnum

Occipital

Medulla oblongata

Hypoglossal canal

Occipital

Hypoglossal nerve

Jugular foramen

Between temporal and occipital

Glossopharyngeal, vagus, and accessory nerves; sigmoid sinus becomes internal jugular vein

Internal acoustic

Petrous part of temporal

Vestibulocochlear and facial Nerves



Dr. Aiman Al -Maathidy
Monday 23 February 2026

Neonatal Skull



Dr. Aiman Al -Maathidy

Monday 23 February 2026

The bones of the vault of neonate are separated by **unossified membranous intervals** called **fontanelles**





Clinically, the anterior and posterior fontanelles are most important and are easily examined in the midline of the vault.

- ✓ The **anterior fontanelle** is diamond shaped and lies between **the two halves of the frontal bone** in front and **the two parietal bones** behind
- ✓ The fibrous membrane forming the floor of the anterior fontanelle is replaced by bone and is closed **by 18 months of age**.
- ✓ The **posterior fontanelle** is triangular and lies between **the two parietal bones** in front and **the occipital bone** behind. **By the end of the 1st year**, the fontanelle is usually closed and can no longer be palpated.



Types of Skull Fracture

- Linear fracture
 - a/w EDH, SDH
- Depressed fracture
 - a/w focal parenchymal lesions
- Skull base fracture
- Open head injuries
 - Knife, firearm
 - Laceration of dura



Monday 23 February 2026
Dr. Aiman Al -Maathidy

Basilar Skull Fracture


- The most common type involves the temporal bone
- Commonly associated with a tear in the dura leading to a CSF leak
- Classic signs and symptoms are often absent on initial presentation but will develop gradually over the first hours of evaluation
- Raccoon Eyes are caused by bleeding from a fracture site in the anterior portion of the skull base



Monday 23 February
2026


Dr. Aiman Al -Maathidy

Basilar Skull fractures can cause leakage of CSF from meninges



"RACCOON EYES"

Periorbital ecchymosis is a sign of a basal skull fracture. Blood tracks along the periosteum and can collect in soft tissues of the orbital lid.



Dr. Aiman Al -Maathidy
Monday 23 February 2026

