

- 1- case of panic with agoraphobia
 - 2- ADHD: methylphenidate
 - 3- ADHD drugs work on: dopamine, NE
 - 4- medication induced delirium
 - 5- mismatch: trazadone 5HTC2..
 - 6- wernicke's mortality 20%
 - 7- substance use criteria 2 for 12 months
 - 8- ASD in monozygotic twins: 70-90%
 - 9- schizophrenia in first degree relative: 10%
 - 10- for severe and refractory cases: ECT
 - 11- safe and effective during pregnancy : ECT
 - 12- case of countertransference
 - 13- mens rea
 - 14- case of a patient afraid of dogs, best therapy method: systematic desensitization
 - 15- ASD criteria all except: intellectual disability
 - 16- DSM5 criteria for LD,choose the correct answer:
 - A)dyslaxia with normal IQ..3months
 - B)dyscalculia..low IQ
 - C)none of the above
 - D) dyslaxia with low IQ.more than 12months
- Answer: C
- 17- gender dysphoria
 - 18- sleep walking occurs in which stage: 3
 - 19- depressive symptom with diurnal variation: mood

20- case of postpartum psychosis, what's your first step: immediate hospitalization

****21- PTSD most important risk factor: nature of trigger.

22- Which of the following is considered a poor prognostic sign in patients with OCD:

A) Early onset

B) Good insight

C) Hoarding behavior

D) Response to treatment

Answer: C

23- FDA approved drug for resistant OCD: clomipramine

24- A 28-year-old woman reports recurrent episodes of eating very large amounts of food in a short period, feeling out of control. She does not vomit, use laxatives, or exercise excessively afterward. She feels distressed and ashamed about overeating not about weight gaining.

Answer: binge eating disorder

25- Q3 Which condition is considered a poor prognosis in depression?

A) Single episode

B) Postpartum onset

C) Seasonal pattern

D) Double depression

Answer: D

26- atomoxetine: NE reuptake inhibitor

27- malingering is external, factitious is internal

28- waxy flexibility, grimace..

29- patient reports sudden loss of muscle tone in the face when laughing, but remains fully conscious and able to interact.

Which of the following is the most likely diagnosis?

- A) Cataplexy
- B) Sleep Paralysis
- C) catalepsy
- D) Stroke
- E) Bell's Palsy

Answer: A

30- case of separation anxiety disorder

31- A person has persistent, well-systematized false beliefs but does not have hallucinations or No disorganized speech. What is the most likely diagnosis :

- A) Schizophrenia
- B) Delusional Disorder
- C) Schizoaffective Disorder
- D) Brief Psychotic Disorder
- E) Mood Disorder with psychotic features

Answer: B

32- schizoaffective: 2 weeks without mood symptoms

33- patient brought to psychiatric clinic due to behavioral abnormalities,he starts moving his hands toward and away from a glass of water repeatedly, as if unsure whether to pick it up,When asked about what he sees, he says: "I see tiny people, about this small (shows with fingers), walking on the table ,he said also "when the moon is knitting my wall , the clock rythum rum),she takes also all something personally, when the TV is open :

A)Ambitendency,liliputin hallucination , clang association, persecutory

delusion

B)Ambitendency,liliputin hallucination ,derailment,perceptual delusion

Answer: B

34- tertiary TCA except: protriptyline

35- lowest sedation:

36- antipsychotic least weight gain and causes QT prolongation: arpiprazole

37- intermittent explosive?

38-Which of the following is a characteristic of Dependent Personality

Disorder :

A)Perfectionism that prevents task completion

B)difficulty expressing disagreement with others because of fear of loss of support or approval

C)requires four or more specific criteria to be met for diagnosis

D) Excessive devotion to work at the expense of leisure

Answer: B

39-Narcolepsy Central hormone ? Hypocretin