

surgery files



surgery files

** اول اربع اسئلة تكررنا تقريبا لكل الدفع #ارشيف

#تعديلات_الارشيف

اخر سوالين مكتوبين بدون خيارات هيك بالملف
الكبير

BUT

Intraabdominal Compartment syndrome—
increase intraabdominal pressure —
elevate of diaphragm —reduce Tidal
volume —hypoxia and hypercarbia

والسوال الثاني

Compartment syndrome is one of the
complication of electrical burn 🙄

بالتالي مفروض الخيارات صحيحات not except

👁 371 4:45 PM

Reply



Copy



Copy Link



Forward



Report



healing Fibroblasts

fibroblast. Fibroblasts
for initiating angiogenesis

هيك مكتوب السؤال بالارشيف بدون خيارات
وفي مصادر بتحكي انها most imp cell is
fibroblast

* لكن الدكتور بالمحاضرة حكي انها
macrophage واكد عليها 🙋



The most important cell is **the fibroblast**. Fibroblasts peak approximately day 7 from injury and are responsible for initiating angiogenesis, epithelialization, and collagen formation. Aug 27, 2021

29. most important cell in wound healing **Fibroblasts**

surgery files

#تعديلات_الارشيف

All the following conditions are known to have a familial pattern associated with a predisposition to cancer except one:

- a. Colonic polyposis
- b. Breast cancer.
- c. Peutz-Jeghers syndrome.
- d. Gardner's syndrome.
- e. retinoblastoma.

الجواب الصحيح E



4

537 12:24 AM

Dr. Jameel Sawaqid Q for Other lectures

38-All of the following are true regarding the management of anal fissure except

A-Glyceryl trinitate[0.2%] 2-3 times as topical application is used for acute and chronic fissures with a success rate of 50%-85%

B-Botulinum toxin injection into the internal sphincter is used for the chronic fissure with a success rate of 65-85%

C-Topical 2% diltiazem [Ca channel blocker] is effective in 50% of cases

* D-Surgical closed lateral internal sphincterotomy for the failed medical treatment is the treatment of choice

فرع D صحيح

Partial Lateral internal sphincterotomy either with open or closed method is the treatment of choice for chronic anal fissure

** السؤال لدكتور ما بدرسنا لكن وجب

التنبیه



2



1

572 5:52 PM

16.The cause of death in established tetanus is:

- A. Septic shock
- B. Heart failure
- C. Renal failure
- D. Respiratory failure
- E. Pulmonary embolism

#تعديلات_الأرشيف

الجواب الصحيح D



3



1

550 3:41 PM



closure

What intra-abdominal injury is associated with seatbelt use?

Small bowel injuries (L2 fracture, pancreatic injury)

#تعديلات_الأرشيف

In a patient with a seat belt injury.
The most likely organ to be injured
is:

- A. Spleen
- B. Liver
- A. Kidney
- B. Intestine

الجواب الصحيح small bowel



4



1

548 2:51 PM

Reply



Copy



Save Image



Copy Link



What surgery is done for insulinoma?

Surgery is the treatment of choice for insulinomas. Enucleation is indicated in smaller tumors without contact with the main pancreatic duct. The other possibility is a left-side pancreatectomy, with or without splenectomy. A central resection of the pancreas can be performed in some specific cases.

Aug 4, 2023

#تعديلات_الارشيف

A 43 year-old male presented with Whipple triad, by computed tomography he had one cm lesion at the head of pancreas, the best surgical management would be :

Select one:

- a. Enucleation
- b. Pancreaticoduodenectomy
- c. Subtotal pancreatectomy
- d. Distal pancreatectomy

الجواب A



3



1

572 edited 6:36 PM



surgery files

Concerning chronic duodenal ulcer all are TRUE, EXCEPT:

- A. It is more common in males than females
- B. Pain usually occurs two hours after meals
- C. Vomiting is rare unless stenosis has occurred
- D. Increased high fasting gastric secretion is usual
- E. Malignant change occurs in 5-10% of this ulcer

Ans is E

النوت المكتوبه بالملف عند السؤال تابعه لل
gastric ulcer سقطت سهوا

#تعديلات_الأرشيف



490 edited 10:06 PM

Reply



Copy



Copy Link



Forward



Report

