




ARCHIVE

"ENT"



Q1. spot diagnosis ?

- ethmoid polyp
- maxillary polyp
- Allergic fungal sinusitis



Q2. presentation ?

- Bilateral obstruction
- Unilateral obstruction
- Bilateral nasal polyp

Q3. treatment ?

- Antibiotic
- steroid
- steroid + FESS
- steroid + antibiotic, if failed FESS

. presented with vertigo, tinnitus.

Q1. type of hearing loss ?

left sensory neural hearing loss



Q2. vertigo ?

last for 20 minute or more

Q3. Tuning fork test ?

- left Rinne positive



- one of the following artery in post. epistaxis  
sphenopalatine  
ant. ethmoidal  
pre auricular



- All of the following used in post. epistaxis except :-  
ant. packing  
post. packing  
silver nitrate  
vasoconstrictor packing  
surgical ligation

- 5 years old child has fever, ear pain, bulging ear, post. auricular enlarge  
Acute mastoiditis  
cholesteatoma  
otitis externa



- complication of bacterial tonsillitis if doesn't treated  
rheumatic fever  
epiglottitis

- sudden loss of hear all of the following except  
caused by herpes zoster  
hearing loss for 3 days or less  
hearing loss on 30DB or more  
medical emergency  
treated by steroid.

- treatment of stage 1,2 laryngeal cancer  
chemotherapy  
surgery only  
radiotherapy  
Radiotherapy or conservative surgery




- recurrent respiratory papillomatosis

\* Cause ?

HPV infection

\* treatment ?

surgical excision



- 5 years boy

\* ventilation tube in ant. inf quadrant ?

- Chalazion appear ?



- 52 years old man -

- diagnosis ?

mostly malignant

septal deviation

antrochoanal polyp.






\* most common age of adenoiditis?

↳ adenoid become symptomatic.

- 3 - 6 year

below 1 year



\* laryngeal cancer.

CT neck + chest (lung mets).



\* all are true except :

iatrogenic

Trauma

Cocaine

surgical repair has high

success rate.

nasal septal perforation

