

Lung Volumes & Capacities

By

Dr. Nour A. Mohammed


Associate professor of physiology

Faculty of medicine, Muthah University

2025-2026


Lung volumes

- **Tidal volume (TV) = 500 ml**
Vol. of air inspired or expired with each normal breath of normal quiet breathing (**eupnea**)
- **Inspiratory reserve volume (IRV) = 3000 ml** اد ال tidal اللى 500 ملى محسوس منهم
Vol. of air which can be inspired by **maximum forced inspiration** **AFTER** normal inspiration.
- **Expiratory reserve volume (ERV) = 1100 ml**
Vol. of air which can be expired by **maximum expiration** **AFTER** normal expiration.
- **Residual volume (RV) = 1200 ml**
Vol. of air remaining in the lung after maximal expiration.
Can't be tested by spirometry.



Lung capacities → More than one volume

* Si capacity داخل مياں ال Residual مارج اقدر احسبنا بال spirometry



1- Inspiratory capacity (IC):

↳ IRV

- It is the volume of air that can be inspired by maximal inspiratory effort *After* the end of normal resting expiration

* بحز ال tidal
بدنا نحسبه

- $IC = TV + IRV = 500 + 3000 = 3500 \text{ ml.}$

2- Expiratory capacity (EC):

↳ ERV

- It is the volume of air that can be expired by maximal expiratory effort *After* the end of normal resting inspiration

* بحز ال tidal
بدنا نحسبه

- $EC = TV + ERV = 500 + 1100 = 1600 \text{ ml.}$

3- Functional residual capacity (FRC):

- It is volume of air remaining in lungs after normal expiration.

- $FRC = ERV + RV = 1100 + 1200 = 2300 \text{ ml.}$

Can't be tested by spirometry.

4- Vital capacity (VC):

- Volume of air expired maximally after maximal inspiration.

- $VC = IRV + TV + ERV = 3000 + 500 + 1100 = 4600 \text{ ml.}$

5- Total lung capacity (TLC):

- Volume of air present in the lung at end of maximal inspiration.

- $TLC = VC + RV = 4600 + 1200 = 5800 \text{ ml}$

Can't be tested by spirometry.

Static pulmonary function tests

ما ربح أقدر
أعيسى RV
باد spirometer
طبي كيف حسيه؟؟


1. Residual volume:

Measured by **Helium dilution method**, using the dilution principle


$$C_1 \times V_1 = C_2 \times V_2 \rightarrow V_1 + RV$$

Helium is used as an inert gas & not diffuse to blood from alveolar air

بنقدر نقيس ال functional residual capacity بنفس الطريقة
بس بنطلب من المريض قبل يعمل normal expiration



Before equilibration



After equilibration

* أول اشي
forced expiration

عشانه بس يظل ال

RV

$n_1 = n_2$
اد amount
تأبته

$$C_1 V_1 = C_2 V_2$$

كيف تأبته :-

closed circulation

ما ياشي بالجسم بطرح helium
exchange عند ال alveolar wall بالتالي ما ربح يفرق ال blood

Importance of Residual volume

- 1) Provides air in alveoli to oxygenate the blood between breaths : *Cause oxygenation of blood during Respiratory pause*
- 2) Prevents lung collapse & Keeps the lung distended
- 3) Prevents marked changes in PO₂ & PCO₂ in the blood with each respiration
- 4) Prevents marked changes in inspired air temperature & humidity
- 5) RV / TLC Less than 30% (increase in **bronchial asthma** & **emphysema** due to **insufficient expiration**)

to prevent damage of the epithelial lining of alveoli

*Substance allergy
IgE
Histamine
Leukotrienes
Mast cell*


6) Medico legal importance

↳ obstructive lung disease

It determines cause of death of baby after birth

If baby is born alive, he will respire, so contain RV → lung float in water while If baby is born dead, he will not respire, so no RV → lung sink in water

loss of elastic element of lung because of high activity of elastase enzyme → No elastic recoil → So no sufficient expiration



Minimal air: Few air remain in lung even after lung collapse
(150 ml)

2. Total lung capacity (TLC)

- Definition: the volume of air present in the lung at the end of maximal inspiration
- Measurement:


$$\text{TLC} = \text{IRV} + \text{TV} + \text{ERV} + \text{RV}$$

$$\text{TLC} = \text{VC} + \text{RV}$$

Normal value: 5800 ml

- Significance:

Decreases in pneumothorax → $\frac{\text{air}}{\text{IP}}$ وجود في



3. Vital capacity (VC)

Definition: It is the amount of air expired maximally after maximal inspiration

Measurement: by spirometer

Value: $VC = IRV + TV + ERV = 4600 \text{ ml}$

Significance:

It indicates the [★]strength of respiratory muscles and [★]lung elasticity

Factors affecting Vital Capacity


	Increase	Decrease
		<p>Handwritten notes in the 'Decrease' header:</p> <ul style="list-style-type: none"> Female: 20-25 (with Arabic text: 20-25 نسوة) male: 25-30 (with Arabic text: 25-30 رجال) muscle power (with Arabic text: قوة العضلات) Limit the descending of diaphragm (with an arrow pointing to the right)
Physiological	Athletes	Females, old age, pregnancy and <u>recumbent position</u> due to return of more blood to the lung. ←
Pathological		<p>a- Chest wall diseases:</p> <ul style="list-style-type: none"> - Paralysis of respiratory muscles & myasthenia gravis - Fracture ribs or kyphosis (limit expansion of thorax) <p>b- Lung diseases:</p> <ul style="list-style-type: none"> - Decreased <u>compliance</u> (stretchability) as (fibrosis, hydrothorax, pneumothorax) <small>↳ indicator of stretchability</small> - Decreased elasticity as (emphysema) - Obstructive conditions like bronchial asthma as resistance to air flow <u>mainly during expiration</u> <p>c- Increased blood volume in the lung:</p> <p>as in pulmonary congestion by left side heart failure.</p> <p>d- Presence of intra-abdominal masses: as tumour and ascites. So, prevent free descent of diaphragm.</p>

Dynamic pulmonary function tests

❖ Respiratory minute volume (RMV) (Minute ventilation):

It is the volume of air respired/min.

At rest = TV x respiratory rate = 0.5 x 12 = 6 L/min.



$$\text{Minute ventilation} = \text{respiratory rate (RR)} \times \text{tidal volume (V}_T\text{)}$$

Dead space (DS)

➤ **Def.:** Volume of air which does not undergo gas exchange in respiratory system

➤ **Types:**

1. **Anatomical DS:** thick respiratory passages (from nose to terminal bronchioles).

2. **Alveolar DS:** non functioning alveoli (normally absent)

* pathological



* ممكن يكونه بسبب thrombosis clot مسكوه
الطريقه مع الدم وبالتالي ما يصير gas exchange

3. **Physiological DS:** = anatomical + alveolar DS.

Normally, DS = anatomical = **150 ml**

N.B.: Inspiration through a tube → **increases DS**

زود ال space

Significance of dead space


- 1) Protective functions
- 2) Prevents marked changes in **PO₂** & **PCO₂** in the blood with each respiration.
- 3) Prevents marked changes in inspired air temperature & humidity.
- 4) It is responsible for difference between Respiratory minute volume (**RMV**) & Effective ventilation volume (**EVV**)

❖ Effective ventilation volume (EVV):

It is the volume of air that enters in gas exchange/ min.

At rest = $(TV - DS) \times \text{respiratory rate} = 0.35 \times 12 = 4.2 \text{ L/min.}$

Dead space



❖ **Maximum breathing capacity (MBC) or maximum voluntary ventilation:**

Maximal volume of air that can be inspired or expired using the deepest and fastest respiratory movements.

Measured in 15 seconds then multiplied by 4.

MBC= 80 to 160 L/min in **males**, 60 to 120 L/min in **females**.

~100 L/min

★ Sustained maximal breathing (>1 minute) → syncopal attack → ↓ CO₂ → respiratory alkalosis → tetany

Dysapnea Painful breath with awareness

❖ Breathing reserve:

- The ^{↳ (spare)} difference between the MBC and RMV
- $BR = 100 - 6 = 94 \text{ L.}$

❖ Dyspneic index (DI):

- The percentage between the **breathing reserve** and the **MBC**.
- Normally $DI > 90\%$
- If $DI < 70\%$ Dyspnea

$$\frac{BR}{MBC} = \frac{94}{100} > 90\%$$


إذا كان أقل من 90%
Dysapnea

❖ Timed vital capacity:

- ❑ **FEV1**: The fraction of vital capacity expired maximally and rapidly in the first second. **FEV1 = 83% of VC**, and reaches **97% in three seconds** (good test for airway resistance so, it is helpful in **obstructive lung diseases** diagnosis & prognosis (e.g. asthma & emphysema))

4 sec - لیس جی

Healthy



Obstructive lung disease

- E.g. Asthma & Emphysema
- VC decreased
- FEV1 decreased markedly
- FEV1/ VC is reduced
- TLC is almost normal
- RV is increased

Restrictive lung diseases

- E.g. Lung fibrosis
 - VC is decreased
 - FEV1 is decreased
 - FEV1/ VC *may be normal*
- As both decreased equally*
- TLC reduced

THANK YOU

