

SOLUTIONS AND BUFFERS

Professor Sameeh Al-Sarayreh

Professor of Medical Biochemistry

Department of Biochemistry and Molecular Biology

Faculty of Medicine, Mutah University

Solutions

A solution is a homogeneous mixture of at least two substances (one is called a solute and the other is called a solvent)

Solutes and solvents can be gases, liquids or solids

1- Liquid solutions



Liquids dissolve gases, other liquids, and solids.

An example of a dissolved gas is oxygen in water, which allows fish to breathe under water.

An examples of a dissolved liquid is ethanol in water, as found in alcoholic beverages.

An example of a dissolved solid is salt in water

eg: NaCl solution



2- Gaseous mixtures

If the solvent is a gas, gases (non-condensable) or vapors (condensable) are dissolved under a given set of conditions.

An example of a gaseous solution is air (oxygen and other gases dissolved in nitrogen).

3-Solid solutions

If the solvent is a solid, then gases, liquids, and solids can be dissolved.

Gas in solids:

Hydrogen dissolves rather well in metals, especially in palladium

Liquid in solid:

Mercury in gold, forming an amalgam

Solid in solid:

Radium sulfate dissolved in barium sulfate

Concentration Expressions:

Molarity = number of moles / Volume in Litre

Number of moles = mass (gm) / molar mass

Example:

Prepare 1 liter of 1.00 M NaCl solution.

First calculate the molar mass of NaCl which is the mass of a mole of Na plus the mass of a mole of Cl (or $22.99 + 35.45 = 58.44$ g/mol)

Molarity = number of moles/ volume (L)

Number of moles = molarity X Volume = $1 \times 1 = 1$

Number of moles = mass/ molar mass

Mass = molar mass X number of moles

Mass = $58.44 \times 1 = 58.44$ gm

Weigh out 58.44 g NaCl.

Place the NaCl in a 1 liter volumetric flask

Add a small volume of distilled, deionized water to dissolve the salt.
Fill the flask **upto** the 1 L line.



Normality = Molarity * n

n: is number of H⁺ or OH⁻

eg: for 1M HCl solution the normality of this solution is equal to 1



For 1M solution of H₂SO₄ the normality of this solution is equal to 2



For 3N solution of H₂SO₄, the Molarity of this solution is equal to ?



Normality = Molarity * n
in this case n is equal to 2

$$3 = M * 2$$

$$M = 3/2 = 1.5$$

For 6M solution of $\text{Al}(\text{OH})_3$, the Normality of this solution is equal to ?



Normality = Molarity * n
in this case n is equal to 3

$$\text{Normality} = 6 * 3 = 18$$

Dilution Formula

$$C_1 V_1 = C_2 V_2$$

C1 is the initial concentration
V1 is the initial volume

C2 is the final concentration
V2 is the final volume

Example:

A laboratory procedure calls for 250 ml of an approximately 0.10 M solution of NH_3 . Describe how you would prepare this solution using a stock solution of concentrated NH_3 (14.8 M).

Substituting known volumes in equation

$$\begin{array}{ccccccc} \mathbf{C1} & & \mathbf{V1} = & \mathbf{C2} & & \mathbf{V2} & \\ 14.8 \text{ M} \times V_1 & = & 0.10 \text{ M} \times & 250 \text{ mL} & & & \end{array}$$



and solving for V_1 gives 1.7 mL.

Since we are making a solution that is approximately 0.10 M NH_3 we can use a micropipette to measure the 1.7 mL of concentrated NH_3 , transfer the NH_3 to a volumetric flask (250 ml), and add sufficient water to give a total volume of approximately 250 mL.

BUFFERS

A buffer is a solution that resists the change in pH, a buffer can be a weak acid and its conjugate base (or a weak base with its conjugate acid). The weak acid is able to donate H^+ ions to neutralize incoming basic ions while the conjugate base is able to accept H^+ .

Practically a buffer is prepared from mixing weak acid and its strong salt, example:




The pH scale is a logarithmic scale representing the concentration of H⁺ ions in a solution which equal the negative log of the H⁺ ions concentration.

$$\text{pH} = -\log [\text{H}^+]$$

[H⁺] = 10⁻⁷ mol/L neutral solution pH = 7

[H⁺] > 10⁻⁷ mol/L acidic solution pH < 7

[H⁺] < 10⁻⁷ mol/L basic solution pH > 7



The pH of a buffer solution can be determined using the Henderson-Hasselbalch equation.



$$\text{pH} = \text{pK}_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$$

K_a : dissociation constant for the weak acid

pK_a (- log K_a) for weak acid.

$[\text{HA}]$ = concentration of the buffer weak acid

$[\text{A}^-]$ = concentration of conjugate base for the buffer weak acid.



pH test strips



pH meter

Types of buffers

1-Synthetic: Can be prepared in the lab.

2.Physiological buffers (natural) :

The main physiological buffers are the bicarbonate, proteins (example haemoglobin), and the phosphate buffers .

Bicarbonate buffer

This is the most important buffer in blood . It is made from equilibrium between carbonic acid and its conjugate base bicarbonate.



Phosphate buffer

The phosphate buffer consists of dihydrogen phosphate ion (H_2PO_4^-) in equilibrium with monohydrogen phosphate ion (HPO_4^{2-}) and H^+



The weak acid, H_2PO_4^- , and its conjugate base, HPO_4^{2-} , are in equilibrium