



CVS MODULE – 4

RHEUMATIC HEART DISEASE

DR. EMAN KREISHAN, M.D.

25-11-2025


RHEUMATIC HEART DISEASE

- Rheumatic heart disease is the cardiac manifestation of rheumatic fever.
- Rheumatic fever is an acute, immunologically mediated, multisystem inflammatory disease that occurs after group A β -hemolytic streptococcal infections (usually pharyngitis, but also occasionally infections at other sites, such as skin).
- valvular inflammation and scarring produce the most important clinical features, rheumatic heart disease is essentially the only cause of acquired mitral stenosis.

PATHOGENESIS


- The pathogenic mechanism involves autoantibodies and T cells directed against group A streptococcal cell wall components that cross-react with heart or brain.
- Predisposing factors include genetic and environmental factors such as poverty and overcrowding.

Pathogenesis





GROSSING

MITRAL VALVULITIS



MITRAL SCARRING






The pericardium may exhibit a fibrinous exudate, which generally resolves without sequelae.




HISTOLOGY


- Acute rheumatic valvulitis manifests with :
 - ✓ active inflammation characterized by lymphocytic infiltration, Aschoff bodies, and Anitschkow cells (have abundant cytoplasm and nuclei with chromatin that is centrally condensed).
- Chronic rheumatic valvulitis is associated with:
 - ✓ neovascularization, valvular fibrosis and/or dystrophic calcification.
- During acute rheumatic fever, Aschoff bodies can be found in any of the three layers of the heart—pericardium, myocardium, or endocardium (including valves).



Aschoff nodules






Anitschkow cells.



HISTOLOGY CONT.:

- Valve involvement results in fibrinoid necrosis and fibrin deposition along the lines of closure forming 1- to 2-mm vegetations—verrucae—that cause little disturbance in cardiac function.



CLINICAL FEATURES

- Acute rheumatic fever occurs most often in children; the principal clinical manifestation is carditis.
- However, about 20% of first attacks occur in adults, with arthritis being the predominant feature.
- Symptoms in all age groups typically begin 2 to 3 weeks after streptococcal infection.
- The clinical signs of carditis include pericardial friction rubs and arrhythmias; myocarditis may be sufficiently severe to cause cardiac dilation and resultant functional mitral insufficiency and CHF.

RHEUMATIC FEVER

DUCKETT-JONES DIAGNOSTIC CRITERIA

MAJOR CRITERIA


“CASES” is the Mnemonic

- C**arditis
- A**rthritis
- S**ubcutaneous nodules
- E**rythema marginatum
- S**ydenham's chorea

MINOR CRITERIA

“FRAPP” is the Mnemonic

- F**ever
- R**aised ESR/CRP
- A**rthralgia
- P**rolonged PR Interval
- P**revious RF



There must be evidence of streptococcal infection plus:

2 major or 1 major + 2 minor

Erythema Marginatum



LAB

- ESR, CRP.
- Elevated serum titers of antibodies against one or more streptococcal antigens (e.g., streptolysin O or DNAase) .
- Cultures are negative for streptococci at the time of symptom onset



Latex agglutination
Slide test
ASO

TREATMENT

- Treatment strategies for acute rheumatic fever (ARF) can be divided into the following:
 - ✓ Management of the acute attack
 - ✓ Management of the current infection
 - ✓ Prevention of further infection and attacks