

Surveillance and Outbreak investigation



Dr. Israa Al-Rawashdeh MD, MPH ,PhD
Faculty of Medicine/Mutah University
2025

Learning Objectives:

- Describe the purpose of public health surveillance
- Differentiate between types of surveillance systems
- Explain the framework and steps for outbreak investigation



578



1K



5K



Introduction to Surveillance

- التقصي الوبائي مستمر
- ▶ **Definition:** ▶ Surveillance is the ongoing, systematic collection, analysis, interpretation, and dissemination of **health-related (already available) data** for planning, implementing, and evaluating public health practice.



Information for Action

Introduction to Surveillance

- Surveillance provides the information for descriptive epidemiology, which is
 - Person (age, sex, description)
 - Place (where)
 - Time (dates, hours, days, months, years)

Most countries mandate reporting of selected **notifiable diseases** to ensure early detection and response:

- Vaccine-preventable diseases (e.g., measles, polio)
- Epidemic-prone infections (e.g., cholera, meningitis)
- High-impact communicable diseases (e.g., tuberculosis, hepatitis)
- Emerging and re-emerging infections (e.g., COVID-19)

Jordan eradicated this disease

↳ So one case is emergency.

If one case can threaten the community, it becomes notifiable

Purposes of surveillance:

Surveillance is the backbone of public health

5

- ▶ **Detect outbreaks** early to initiate a response
- ▶ **Monitoring Diseases' trends:** Routine observation of diseases with predictable patterns and Forecasting changes
- ▶ **Evaluate the effectiveness of interventions** (e.g., vaccination, hygiene practices).
- ▶ **Guide allocation of resources** (basic for costing studies).
- ▶ **Global Health Security:** Contributing to preparedness and response in international outbreaks (e.g., COVID-19).



عزيمر الخابرات
(Epidemiological Intelligence!)

Example: Surveillance of influenza cases during seasonal outbreaks to identify peaks, forecast outbreaks and plan vaccination campaigns.

Key Components of Surveillance

Data Collection:

- Systematic collection of data from hospitals, clinics, laboratories, and community health systems.
- Includes case counts and essential demographic and clinical information.
- Example: Collection of COVID-19 cases from hospitals worldwide

Data Analysis:


- Identification of patterns and trends by person, place, and time.
- Detection of unusual increases that is warning sign for an outbreak.
- Use of basic epidemiological tools and statistical methods.

Dissemination:

- Sharing analyzed data with public health authorities and stakeholders.
- Ensures surveillance data are translated into public health action.
- Formats: Reports, announcement, and press releases.
- Example: Weekly WHO reports on global disease surveillance.

Data that are not analyzed and disseminated are useless for public health

(Case Surveillance)



Types of surveillance

الأشكال المهمة



دورك هياحت تطلع على التقارير الي تسجلها المستشفيات بشكل روتيني

1. Passive Surveillance

- ▶ Passive surveillance depend on routine reporting of cases by healthcare providers to public health authorities.
- ▶ Physicians, clinics, laboratories, and hospitals that are required to report disease are given the appropriate forms and instructions, with the expectation that they will record all cases of reportable disease that come to their attention.
- ▶ It is cost-effective but can lead to underreporting.
- ▶ Most public health surveillance systems are passive
- ▶ Example: Tracking tuberculosis cases through mandatory hospital notifications.

لجزء تليفو المسؤوليين

Advantages	Disadvantages:
Simple	Under-reporting
cost-effective	incomplete data
Suitable for routine data collection	Delays in reporting

تأخر

Types of surveillance

متى استخدم هذا النوع؟!

ما قبل حالات * مرفحة عملنا الـ eradication وبيننا تتأكد

* بالأشخاص الي عدد الحالات فيهم قليل

* بالمناطق الي فيها weak reporting

ACTIVE SURVEILLANCE



PSA



BIOPSY



MRI

2. Active Surveillance → هون انت كبايت تقوم وتدور على الحالات

مثال عليها :- المسوح الطبية

PRIMR

- Proactively searching for cases through direct contact, field visits, or surveys.
- It is commonly used in elimination or eradication programs.
- (Smallpox).
- Negative reporting is also **necessary** (report of no cases).
- Requires periodic telephone calls, electronic contact or personal visits to the reporting individuals and institutions to obtain the required data.
- Example:** Door-to-door polio case identification campaigns

انك تبليخ انه ما في حالات
مهم زي زي تبليخك
انه في حالات

Advantages:	Disadvantages:
Accurate (higher quality data) and timely.	Expensive
Identifies unreported or missed cases.	resource-intensive

ستقوم عمل
وادوار

Types of surveillance

تكاليف مؤسسات أوسرالر طبية معينة لتبليغك عن أي حالة تجديهم

3. (Sentinel Surveillance)

- ▶ Monitors selected sites that represent a larger population ("sentinel sites"-e.g. GP Clinics). ٥٥
- ▶ Can be active or passive.
- ▶ Provides high-quality, detailed data for specific diseases.
- ▶ **Example:** HIV surveillance in sentinel antenatal clinics in sub-Saharan Africa to estimate population infection rates.

Types of surveillance

* بیعتمد علی clinical manifestation یعنی ما یستنبی لحتی نتایج اللاب تطلع

4. Syndromic Surveillance

- ▶ Monitoring symptoms and health indicators (cases that meet a clinical case definition for disease under surveillance) before diagnosis is confirmed (before laboratory confirmation)
- ▶ Helps in Rapid/Early detection of unusual health events.
- ▶ **Example:**
 - Monitoring acute fever/rash to monitor measles or rubella.
 - STDs
 - Using emergency room visits to detect potential bioterrorism events in the 2001 U.S. anthrax attacks.

Investigation of outbreaks

Why Investigate Outbreaks?

1. • **Control the current outbreak**
2. • **Prevent future outbreaks (Small Outbreaks May Uncover Widespread Problems)**
A localized outbreak can reveal larger systemic issues (e.g., contaminated water or unsafe practices).
3. • **Improve understanding of disease :**
 - Can be First Clue in Discovery of New Diseases or Organisms
Outbreak investigations often lead to identifying **new diseases** or previously unknown infectious agents. Example: HIV, SARS, or COVID-19 were initially identified through outbreak analysis.
 - Understanding the Disease and Organisms
Investigations provide insights into:
 1. **Incubation period**
 2. **Modes of transmission**
 3. **Natural history of the disease**



Investigation of outbreaks

Ten principles of investigation

Solving mystery action

* 5 دلو 5 دلو




1. Determine the existence of an outbreak
2. Confirm the diagnosis
3. Define and identify cases
4. Collect information
5. Describe and Analyze data
6. Develop a hypothesis
7. Test the hypothesis
8. Implement control and prevention measures
9. Communicate findings
10. Prepare a report



1. Determine the existence of an outbreak

تحديد تفشي المرض

- ▶ Verify whether the observed number of cases exceeds what is expected (using surveillance data and comparison to historical trends considering factors like seasonal variations).



In July 2020, a sudden increase in food poisoning cases was reported in Amman, Jordan, with over 800 individuals affected and one child tragically dying, encouraging further investigation into the outbreak.



Remember:

Compare the Observed with the Expected


- ▶ Increases in case numbers can be due to factors other than a true outbreak, including:
 - ▶ Changes in case definitions, → حالات تُعتبر outbreak
 - ▶ Adjustments to the surveillance system, → طرق وسائل الرصد هي تتغير
 - ▶ Seasonal trends, → في الأوقات الموسمية
 - ▶ Improvements in laboratory testing procedures,
 - ▶ Increased public awareness, or
 - ▶ Laboratory errors.

A true outbreak cannot be confirmed or ruled out until all these possibilities have been considered and investigated. →

لازم نجاوب على كل

النقاط الضوئ (8) علينا نحكى انه الأعداد وخلق زادت و صار في outbreak

Number of Reported Dysentery Cases by Month




Suspicious but not certain, we need historic data to rule out seasonal variation

* يعني بيانا Data

للمنوت الماوية علينا نتأكد انه الموضوع من موسمي

2. Confirm the diagnosis

- ▶ Confirm the correct diagnosed **through clinical assessment and laboratory testing**. This step ensures that the outbreak is not due to misdiagnosis or laboratory error.
- ▶ Samples not only just those affected (blood, stool, vomit..etc), but also possible sources (such as food or water) or environment (kitchen surfaces and utensils).
- ▶ Example: during the 2019 Campylobacter outbreak in Norway, laboratory tests confirmed the presence of Campylobacter bacteria in patients, validating the clinical diagnosis and confirming the outbreak's etiology.



3. Define and identify cases

- ▶ Establish a **Case Definition**: A standard set of criteria used to classify individuals as having the disease under investigation, based on person, place, time, and clinical features.
- ▶ Includes **four components**:
 - Information about the event,
 - Characteristics about the people who are affected,
 - Information about the location or place,
 - A specification of time during which the outbreak occurred.

*البيوتوزة بتجيب لنا أكثر من تعريف للحالة

واما بنحدد مين ادقه فيهم

(لازم يكون في التعريف ال 4 عناصر)


- ▶ Example 1: A ^{التعريف كامل} group of students studying at ^{People who affected} university X, who ate at the same cafeteria experienced ^{event} diarrheal illness between the months of ^{Place} January and ^{time} February.
- ▶ Example 2: ^{التعريف كامل} In the 2015 Legionnaires' disease outbreak in New York City, the case definition included individuals with pneumonia onset after a specific date, residing or working in a particular area, and with laboratory confirmation of Legionella bacteria.

3. Define and identify cases

- ▶ An ideal case definition includes most true cases while minimizing false positives (people without the disease who meet the definition).
- ▶ Due to diagnostic uncertainty, **Cases** are often classified as: → *صوت تاكدنا*
- ▶ **Confirmed:** A case usually must have laboratory verification
- ▶ **Probable:** A case has typical clinical features of the disease without laboratory confirmation
- ▶ **Possible:** A case has fewer of the typical clinical features, epidemiological link not confirmed.

→ *حسا حدوده كاتم cases ال outbreak لب نصيفه حسب تاكدنا*

Classifying a COVID-19 case using the CSTE definition



4. Collect information:

▶ “Line listing”

A line listing is a **table** that summarizes key information for each identified case during an outbreak investigation.

The columns represent specific patient information and the rows represent each case.

Information to be included in the line list:

- Identifying information: name, address, contact information.
- Demographic information: Age, sex, date of birth, current status (dead or alive)
- Clinical information: date of report, date of onset of symptoms, presenting symptoms, diagnosis, laboratory findings.
- Relevant risk factor information: last meal eaten (what and where), contact with animal, ingestion of undercooked meat, ingestion of water from contaminated source.

Line listing:

					Diagnostic						Lab			
					Signs and Symptoms									
Case#	Initials	Date of Report	Date of Onset	Physician Diagnosis	N	V	A	F	DU	J	HAIGM	Other	Age	Sex
1	JG	10/12	12/6	Hep A	+	+	+	+	+	+	+	SGOT ↓	37	M
2	BC	10/12	10/5	Hep A	+	-	+	+	+	+	+	Alt ↓	62	F
3	HP	10/13	10/4	Hep A	±	-	+	+	+	S*	+	SGOT ↓	30	F
4	MC	10/15	10/4	Hep A	-	-	+	+	?	-	+	Hbs/Ag-	17	F
5	NG	10/15	10/9	NA	-	-	+	-	+	+	NA	NA	32	F
6	RD	10/15	10/8	Hep A	+	+	+	+	+	+	+		38	M
7	KR	10/16	10/13	Hep A	±	-	+	+	+	+	+	SGOT = 240	43	M

منطقه برات
المنطقه



S* = Sclera; N = Nausea; V = Vomiting; A = Anorexia; F = Fever; DU = Dark urine; J = Jaundice;
 HAIGM = Hepatitis AIgM antibody test

5. Describe and Analyze data

Descriptive analysis: Time, Place, person

Place

- ▶ Place data can be included in line listing (text).
- ▶ Or a map to visualize the outbreak. (mapping) *Bebber*
- ▶ Two types:
 - **A spot map:** A map that marks individual cases at specific locations based on certain characteristics, such as **where they live, work, or were exposed**
 - **An area map:** A map that displays the number of cases aggregated by **geographical areas**, such as districts, neighborhoods, or regions.




5. Describe and Analyze data

Descriptive analysis: Time, Place, person


Person:

- Age, sex, occupation, education, social class, economic class (income), marital status, race and ethnicity, underlying medications, and others. (Table or visually)

Spot Chart: Cases by Age and Sex



Bar Chart: Cases by Age Group



مثال على سؤال: "شو أكثر فئة عمرية أخصيبت بالوبئة؟"

5. Describe and Analyze data

Descriptive analysis: Time, Place, person

Time

Epidemic curve

- ▶ An epidemic curve is a histogram showing the number of cases over time, providing clues about the source and mode of transmission.
- ▶ An epidemic curve (epi curve) is a graphical display of the number of disease cases (y axis) over time unit (x axis) ①
- ▶ The shape of the curve can give clues to the transmission route and source of the epidemic. ②
- ▶ Reports on the time course of an epidemic in a defined population

Time can be weeks, days, hours according to the incubation period. E.g. Bacillus cereus
incubation period 1-6 hours


↓
hours

Epi curve: A histogram


Column histogram

↳ most common

Cases of an Unknown Disease by Month of Onset



Stacks of boxes histogram



Steps to Create an Epidemic Curve

Collect Data: Gather information on symptom onset dates.

Choose an Appropriate Time Interval:

Based on disease incubation period (e.g., hours, days, weeks..etc).

Plot the Data:

Use bars to represent case counts over time.

Analyze the Shape:

Look for patterns, peaks, and outliers.

Patterns of Epi Curves:


Three common types of outbreak patterns:

1. Common Source Outbreak

a. Point Source Outbreak

• انما يبرأ بنفس الوقت وبنفس المصدر

- ▶ A single exposure event, all cases occur within one incubation period.
- ▶ **Shape:** A sharp peak, followed by a decline.
- ▶ **Example:** Food poisoning at a wedding.



Patterns of Epi Curves:

Three common types of outbreak patterns:



▶ 1. Common Source Outbreak

Source: *مشروع يظلمه لحاله لازم تفتح ال*

b) Continuous Common Source → لاينتهي

- ▶ Ongoing exposure to the source over a period. Multiple incubation periods.
- ▶ **Shape:** A plateau or gradual rise and fall.
- ▶ **Example:** Contaminated water supply.

Example of common source outbreak with continuous exposure:
Diarrheal illness in city residents by date of onset and character of stool,
Cabool, Missouri, December 1989-January 1990



Patterns of Epi Curves:

Three common types of outbreak patterns:

▶ 2. Intermittent Common Source


▶ Exposure to the source occurs intermittently (not continuous or one-time).

▶ **Shape:** Irregular peaks separated by periods with no cases.

▶ **Example:** Sporadic contamination of food at a restaurant.

zero cases →

من صون نادر نغرفه عن
continuous source





Patterns of Epi Curves:

Three common types of outbreak patterns:


▶ **3. Propagated (Person-to-Person) Outbreak**

- ▶ Disease spreads from person to person.
- ▶ a single common source cannot be identified → مافى مصدر واحد
- ▶ **Shape:** Multiple peaks (waves) occur, separated by incubation periods.
- ▶ Each wave represents new generations of transmission.
- ▶ **Example:** Measles, COVID-19, or influenza.






Point source




Continuous source

دقيق



Propagated source



Intermittent source

Zero cases

Source
الفرق بين


Pro → different sources
Inter → same source

Mixed Epidemic

- ▶ Initial Common-Source Exposure:
- ▶ Many individuals are exposed to a single source of infection (e.g., contaminated food or water).
- ▶ Followed by secondary cases caused by (propagated) person-to-person transmission.

Example: Cholera: People exposed to contaminated water initially (common source) may later spread the infection within households (propagated).

- ▶ Shape of the Epidemic Curve:
- ▶ The curve typically shows a sharp peak (indicating the common-source exposure), followed by successive smaller peaks as the disease spreads person-to-person. The smaller peaks represent the incubation period and multiple waves of transmission




Estimating the Time of Exposure: Steps to Determine the Likely Exposure Period:

- ▶ Identify the Peak of the Outbreak: (Find the date with the highest number of cases on the epidemic curve.) This represents **the median** case.
- ▶ Count Back the Average Incubation Period: Look up the average incubation period for the disease.
- ▶ Count backward from the peak date on the x-axis to estimate the likely exposure date.
- ▶ **Adjust for Early Cases:**
- ▶ Identify the **earliest case** on the curve.
- ▶ Count backward using the **minimum incubation period** to estimate the earliest exposure date.

▶ Example


- ▶ **Disease:** Hepatitis A
- ▶ **Average Incubation Period:** 28 days
- ▶ **Minimum Incubation Period:** 15 days
- ▶ Peak Date (Median Case): June 30
- ▶ Earliest Case: June 25
- ▶ **Step 1: Peak = June 30 → Count back 28 days → Likely exposure: June 2.**
- ▶ **Step 2: Earliest case = June 25 → Count back 15 days → Earliest exposure: June 10.**
- ▶ **Conclusion: The likely exposure period is between June 2 and June 10.**

Frequency distribution of the incubation period



- Use the **Date** as the X-axis (time).
- Use the **Number of Cases** as the Y-axis (frequency of cases).
- Plot each date as a **bar**, where the height of the bar equals the number of cases.

Date	Number of Cases
June 1	1
June 2	2
June 3	4
June 4	8
June 5	12
June 6	9
June 7	6
June 8	3
June 9	1



- **What is the peak date?**
- June 5.
- **If the incubation period is 3 days, when might the exposure have occurred?**
- Count back 3 days from the peak (June 5 → June 2)
- **What type of outbreak is this?**
- Point source outbreak (sharp rise and fall of cases).

Steps 6, 7, 8, 9 and 10

▶ Develop Hypotheses

- ▶ Generate theories about the **source of the outbreak, mode of transmission, and risk factors**.
- ▶ Example: "The outbreak was caused by contaminated water from Reservoir X."

▶ Evaluate Hypotheses

- ▶ Test hypotheses using **analytic studies** like:
 - ◆ **Cohort Study**: Compare exposed and unexposed groups.
 - ◆ **Case-Control Study**: Compare cases (sick individuals) with controls (healthy individuals).
- ▶ Calculate measures like **Relative Risk (RR)** and **Odds Ratio (OR)**.

▶ Implement Control and Prevention Measures

- ▶ Apply measures to stop the outbreak:
 - ◆ Isolation, vaccination, sanitation, or public health messaging.

▶ Communicate Findings

- ▶ Prepare and share reports with stakeholders, public health authorities, and the community.
- ▶ Use visual tools like epidemic curves, maps, and tables.

▶ Follow Up and Monitor

- ▶ Ensure control measures are effective.
- ▶ Monitor for additional cases to prevent recurrence.

▸ Thank you

Outbreak Investigations

