

MCQ – Seizures Subtypes 1

❖ All the following are generalized seizure, except:

فاينل (1)



- a. Tonic clonic
- b. Temporal lobe seizure
- c. Atonic seizure
- d. Absence seizure
- e. Myoclonic seizure

❖ Partial epilepsy is characterized by all of the following except:

فاينل معدل (1)



- a. Temporal lobe epilepsy can be associated with aura
- b. Temporal lobe epilepsy can be associated with automatism
- c. Partial epilepsy may progress to loss of consciousness
- d. Partial epilepsy can be associated with temporary paralysis of the limb (Todd's paralysis)
- e. Frontal lobe is associated with oroalimentary automatism (associated with temporal not frontal)

MCQ – Seizures Subtypes 2

فاينل (1)

❖ Partial epilepsy is characterized by all of the following except:

- * a. Temporal lobe epilepsy can be associated with aura
- b. Temporal lobe epilepsy can be associated with automatism
- c. Partial epilepsy may progress to loss of consciousness
- d. Partial epilepsy can be associated with temporary paralysis of the limb (Todd's paralysis)
- e. It does not progress to secondary generalization

فاينل (1)

❖ Which of the following statements is FALSE regarding absence epilepsy (petit mal)?

- * a. Typically starts in childhood
- b. Females are more affected
- c. Child becomes fatigue and tired for several hours after the attack (post ictal state takes hours)
- d. Characterized by daily attacks each lasting for seconds
- e. Associated with 3 H-z generalized symmetrical spike-wave complexes

MCQ – Seizures Subtypes 3

فاينل (2)

❖ **Elementary auditory seizures such as humming or buzzing sound arise from which area ?**

- a. frontal lobe
- b. frontorbital area
- c. lateral temporal lobe
- d. mesial temporal lobe
- e. parietal lobe

فاينل (1)

❖ **A transient neuro deficit, contralateral to seizure site is called:**

- a. Quadriplegia
- b. Hemiplegia
- c. Todd's Paralysis
- d. Dystonia
- e. Myoclonus

MCQ – Seizures Subtypes 4

فاينل (1)

❖ Regarding Primary generalized epilepsy, which of the following statements is FALSE ?

- a. Typically, tonic clonic
- b. Associated with loss of consciousness
- c. Incontinence does not occur
- d. Patient might remain tired for several hours after the attack
- e. Patient might bite their tongue

فاينل (1)

❖ All of the following about generalized epilepsy are true, EXCEPT:



- a. Composed of tonic and clonic stages.
- b. There is loss of consciousness.
- c. Post ictal state lasting for few minutes.
- d. Urinary incontinence occurs during clonic stage.
- e. EEG may confirm the diagnosis